

Appendix 1. List of plant diversity in the study area

Family name	Species name	Location, Habitat, Altitude, GPS Location, Collected Date, Collector Number, Chorotypes, Life Forms
PTERIDOPHYTA		
Aspleniaceae		
	<i>Asplenium ceterach</i> L.	İbradı, Suçikan Location way, rock, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2753; İbradı, around of Yirme Hill, rock in <i>Cedrus libani</i> forest, 1783 m, N 37°14'667" – E 31°23'097", 18.v.2008, Çinbilgel 3188; Manavgat, Akpınar Plateau (Deller Location), rock in mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4828, (H).
	<i>Asplenium onopteris</i> L.	Manavgat, between Dermebeli and Sürüçkan Mountains, rocky slopes, 1917 m, N 37°12'891" – E 31°19'890", 10.vi.2008, Çinbilgel 3766, (H).
	<i>Asplenium septentrionale</i> (L.) Hoffm.	Manavgat, Kaldırım Mountain, rocky places, 1647 m, N 37°09'558" – E 31°20'297", 01.vi.2009, Çinbilgel 4919; Manavgat, Kovacık Mountain, rock, 1952 m, N 37°13'631" – E 31°20'057", 29.vi.2009, Çinbilgel 5488; İbradı, Melik Mountain (Orta Hill), rock, 2077 m, N 37°16'553" – E 31°23'091", 17.vi.2010, Çinbilgel 6414, (H).
	<i>Asplenium trichomanes</i> L.	İbradı, Suçikan Location way, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2752; Manavgat, Kızılpınar Plateau (Nurboğazı Location), rocky places, 1907 m, N 37°14'167" – E 31°20'459", 11.vi.2008, Çinbilgel 3828; İbradı, Toka Plateau way (Gevenli Location), rock, 1143 m, N 37°12'008" – E 31°24'420", 28.iii.2010, Çinbilgel 5942, (H).
Cystopteridaceae		
	<i>Cystopteris fragilis</i> (L.) Bernh.	Manavgat, Burunçalı Location, 1891 m, N 37°09'118" – E 31°19'014", 02.vi.2009, Çinbilgel 5062; İbradı, Çeşgar Plateau, rock, 1618 m, N 37°17'457" – E 31°23'710", 01.vii.2010, Çinbilgel 6881; İbradı, between Boğazyurt Location and Yaylacık Mountain foothills, rock, 965 m, N 37°10'610" – E 31°25'419", 06.iii.2011, Çinbilgel 7377, (H).
Dryopteridaceae		
	<i>Dryopteris pallida</i> (Bory) Fomin subsp. <i>pallida</i>	İbradı, Toka Plateau way, Belova Location, rocky places, 1695 m, N 37°13'638" – E 31°23'243", 19.v.2008, Çinbilgel 3436; Manavgat, Kaldırım Mountain, rocky places, 1647 m, N 37°09'558" – E 31°20'297", 01.vi.2009, Çinbilgel 4920; İbradı, around of Çeşgar Plateau Breach, stony places, 1641 m, N 37°17'441" – E 31°24'212", 01.vii.2010, Çinbilgel 6816, Medit., (H).
Equisetaceae		
	<i>Equisetum giganteum</i> L.	İbradı, Melik Plateau, streamsides, 1537 m, N 37°15'549" – E 31°24'633", 02.vii.2010, Çinbilgel 6945; İbradı, Melik Plateau, streamsides, 1388 m, N 37°15'606" – E 31°24'669", 05.x.2010, Çinbilgel 7319, (H).
	<i>Equisetum telmateia</i> Ehrh.	Manavgat, Burmahan Neighbourhood, around of Erenli Location, streamsides, 753 m, N 37°09'844" – E 31°16'991", 26.iii.2011, Çinbilgel 7432; Manavgat, Burmahan Neighbourhood, around of Erenli Location, streamsides, 773 m, N 37°09'851" – E 31°16'989", 06.vi.2011, Çinbilgel 7844, (H).
Pteridaceae		
	<i>Chelianthes persica</i> (Bory) Mett. ex Kuhn	Manavgat, Derme Plateau (around of Kalbur Gözenekleri Location), rock, 1813 m, N 37°13'337" – E 31°20'464", 05.viii.2009, Çinbilgel 5731; İbradı, Yukarıtoka Plateau, rock, 1733 m, N 37°15'405" – E 31°22'545", 18.vi.2010, Çinbilgel 6476; İbradı, Çeşgar Plateau, rock, 1618 m, N 37°17'456" – E 31°23'710", 01.vii.2010, Çinbilgel 6848, (H).
	<i>Chelianthes pteridioides</i> (Reich.) C. Chr	İbradı, Aşağıtoka Plateau, rock, 1547 m, N 37°13'903" – E 31°22'326", 19.v.2008, Çinbilgel 3379; Manavgat, Kızılpınar Plateau surroundings (between Nurboğazı Location and Toka Plateau way), stony places, 1962 m, N 37°14'256" – E 31°20'649", 11.vi.2008, Çinbilgel 3850, (H).
MAGNOLIOPHYTA Subdivisio I: PINOPHYTINA		
Cupressaceae		
	<i>Cupressus sempervirens</i> L.	Manavgat, between Burmahan Neighbourhood and Çay Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°11'284" – E 31°16'291", 11.x.2009, Çinbilgel 5791, (Ph).
	<i>Juniperus drupacea</i> Labill.	İbradı, Suçikan Location way, <i>Pinus nigra</i> forest, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2737; İbradı, around of Güzle Plateau, mixed forest, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2805; İbradı, Çukuryurt Plateau, <i>Pinus nigra</i> forest, 1248 m, N 37°12'076" – E 31°24'011", 19.v.2008, Çinbilgel 3500, (Ph).
	<i>Juniperus excelsa</i> M.Bieb. subsp. <i>excelsa</i>	İbradı, Güzle Plateau surroundings, forest, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2804; Manavgat, around of Susuzkır Hill, forest, 1666 m, N 37°16'797" – E 31°21'755", 12.vi.2008, Çinbilgel 3989; İbradı, Çeşgar Plateau, rock, 1636 m, N 37°17'445" – E 31°23'646", 01.vii.2010, Çinbilgel 6867; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 974 m, N 37°10'600" – E 31°25'718", 26.ix.2010, Çinbilgel 7203, (Ph).
	<i>Juniperus foetidissima</i> Willd.	Manavgat, Çiğdem Hill surroundings (Çandır Hill), rocky slopes, 2007 m, N 37°14'458" – E 31°20'939", 11.vi.2008, Çinbilgel 3862; Manavgat, Kepezbeli Location, slopes, 1641 m, N 37°10'300" – E 31°21'722", 19.v.2009, Çinbilgel, 4767; İbradı, Melik Mountain (Orta Hill), debris, 1830 m, N 37°15'886" – E 31°22'533", 17.vi.2010, Çinbilgel 6294, (Ph).

	<i>Juniperus oxycedrus</i> L. subsp. <i>oxycedrus</i> var. <i>oxycedrus</i>	İbradı, Around of Güzle Plateau, Juniper forest, 1306 m, N 37°09'817" – E 31°23'980", 20.iv.2008, Çinbilgel 2837; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4638, İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1470 m, N 37°15'607" – E 31°24'429", 30.vi.2010, Çinbilgel 6637, (Ph).
Pinaceae		
	* <i>Abies cilicica</i> (Antoine & Kotschy) Carrière subsp. <i>isaurica</i> Coode & Cullen	İbradı, Around of Güzle Plateau, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2819; İbradı, Yaylacık Mountain (Alıç Location), 1486 m, N 37°10'112" – E 31°24'832", 19.iv.2009, Çinbilgel 4614, Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4635, End., E. Medit., (mt.), (Ph).
	<i>Cedrus libani</i> A. Rich. var. <i>libani</i>	İbradı, Around of Güzle Plateau, mixed forest, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2807; Manavgat, Kızılpınar Plateau, <i>Pinus nigra</i> forest, 1638 m, N 37°14'170" – E 31°19'484", 25.v.2008, Çinbilgel 3635; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4634, Medit. (mt.), (Ph).
	<i>Pinus brutia</i> Ten. var. <i>brutia</i>	İbradı, Gavur Breach – Etülgen boğazi, <i>Pinus brutia</i> forest, 1186 m, N 37°09'074" – E 31°25'116", 05.iii.2011, Çinbilgel 2961, E. Medit., (Ph)
	<i>Pinus nigra</i> J.F. Arnorl subsp. <i>pallasiana</i> (Lamb.) Holmboe var. <i>pallasiana</i>	İbradı, Suçikan Location way, <i>Pinus nigra</i> forest, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2724; Manavgat, Kızılpınar Plateau, <i>Pinus nigra</i> forest, 1638 m, N 37°14'170" – E 31°19'484", 25.v.2008, Çinbilgel 3634; İbradı, around of Çukuryurt Plateau, mixed forest, 1240 m, N 37°12'009" – E 31°24'104", 05.iii.2011, Çinbilgel 7352, (Ph).
MAGNOLIOPHYTA Subdivisio 2: MAGNOLIOPHYTINA		
Acanthaceae		
	<i>Acanthus hirsutus</i> Boiss. subsp. <i>hirsutus</i>	İbradı, Toka Plateau, dwarf bush formation, 1528 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3331; İbradı, between Melik Mountain foothills and Yukarıtoka Plateau, dwarf bush formation, 1599 m, N 37°15'290" – E 31°22'419", 18.vi.2010, Çinbilgel 6468; İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1223 m, N 37°14'468" – E 31°24'259", 02.vii.2010, Çinbilgel 7056, (H).
Adoxaceae		
	<i>Sambucus ebulus</i> L.	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings – roadsides, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3484; İbradı, Suçikan Location – Taşkesiği Location, <i>Pinus nigra</i> forest clearings – roadsides, 1308 m, N 37°15'191" – E 31°24'495", 18.vi.2008, Çinbilgel 4371; İbradı, Ekinlik Location surroundings, <i>Cedrus libani</i> forest (mixed with <i>Pinus nigra</i>), 1532 m, N 37°13'668" – E 31°23'510", 09.v.2010, Çinbilgel 6142, Euro-Sib., (Ph).
Amaranthaceae		
	<i>Chenopodium album</i> L. subsp. <i>album</i> var. <i>album</i>	Manavgat, around of Çandır Hill, dwarf bush formation, 1862 m, N 37°13'965" – E 31°20'869", 04.viii.2009, Çinbilgel 5662; İbradı, Melik Plateau, forest clearings, 1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6967; Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7738, (Th).
	<i>Chenopodium foliosum</i> Asch.	Manavgat, Kızılpınar Plateau, stony places, 1750 m, N 37°14'071" – E 31°19'981", 19.vii.2008, Çinbilgel 4452; Manavgat, Kürtağacıboğazi Location, rock, 1974 m, N 37°12'819" – E 31°18'917", 10.vi.2009, Çinbilgel 5314; İbradı, Melik Plateau, openings – stony places, 1387 m, N 37°15'533" – E 31°24'528", 30.vi.2010, Çinbilgel 6723, (Th).
Amaryllidaceae		
	<i>Allium amethystinum</i> Tausch	İbradı, Akmuar Location, Juniper Forest clearings, 1073 m, N 37°11'976" – E 31°24'786", 02.vii.2010, Çinbilgel 7149, Medit., (G).
	<i>Allium cassium</i> Boiss. var. <i>hirtellum</i> Boiss.	İbradı, Toka Plateau, <i>Pinus nigra</i> forest clearing – stony places, 1544 m, N 37°14'435" – E 31°22'650", 18.v.2008, Çinbilgel 3162; Manavgat, Çağlan Breach, stony places, 1680 m, N 37° 15' 751" – E 31° 20' 812", 12.vi.2008, Çinbilgel 4005; Manavgat, Semai Plateau, stony places, 1557 m, N 37°14'567" – E 31°19'654", 07.vi.2011, Çinbilgel 7916, E. Medit., (mt.), (G).
	* <i>Allium enginii</i> Ozhatay & B. Mathew	İbradı, Aşağıtoka Plateau, stony places, 1517 m, N 37°13'486" – E 31°22'482", 02.vii.2011, Çinbilgel 7990, End., E. Medit., (G), (Conf. Duman).
	<i>Allium guttatum</i> Steven subsp. <i>sardoum</i> (Moris) Stearn (Könem, Gönem)	Manavgat, Kepez Plateau, wet meadows, 1624 m, N 37°11'085" – E 31°21'009", 19.v.2009, Çinbilgel 4796; Manavgat, Kepez Plateau, meadows, 1628 m, N 37°11'110" – E 31°21'021", 03.vi.2009, Çinbilgel 5175; İbradı, Suçikan Location, streamsides, 1251 m, N 37°14'475" – E 31°24'186", 02.vii.2010, Çinbilgel 7107, Medit., (G).
	<i>Allium hirtovaginatatum</i> Kunth	Manavgat, Kızılpınar Plateau, dwarf bush formation, 1782 m, N 37°14'186" – E 31°20'257", 03.viii.2009, Çinbilgel 5604; Kızılpınar Plateau, thorny cushion formation – stony places, 1788 m, N 37°14'441" – E 31°20'293", 03.viii.2009, Çinbilgel 5645; İbradı, Hillsides of Melik Mountain, dwarf bush formation, 2106 m, N 37°16'066" – E 31°23'466", 18.vi.2010, Çinbilgel 6495, Medit., (G).
	<i>Allium myrianthum</i> Boiss.	Manavgat, Argıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4133, Ir. – Tur., (G).
	<i>Allium orientale</i> Boiss.	İbradı, Hillsides of Melik Mountain, rocky slopes, 1880 m, N 37°15'258" – E 31°23'363", 18.v.2008, Çinbilgel 3294; Manavgat, Akpınar Plateau, <i>Cedrus</i>

		<i>libani</i> forest clearings, 1547 m, N 37°10'036" – E 31°21'232", 19.v.2009, Çinbilgel 4808; İbradı, Çukuryurt Plateau, stony places, 1299 m, N 37°12'181" – E 31°23'938", 09.v.2010, Çinbilgel 6162, E. Medit., (G).
	<i>Allium pallens</i> L. subsp. <i>pallens</i>	İbradı, Çeşgar Plateau, rock, 1590 m, N 37°17'623" – E 31°23'753", 01.vii.2010, Çinbilgel 6827, (G).
	<i>Allium paniculatum</i> L. subsp. <i>paniculatum</i>	İbradı, around of Çukuryurt Plateau, <i>Pinus nigra</i> forest, 1227 m, N 37°12'049" – E 31°24'169", 03.vii.2011, Çinbilgel 8002, Medit., (G).
	<i>Allium phanerantherum</i> Boiss. & Hausskn. subsp. <i>phanerantherum</i>	İbradı, Çeşgar Plateau, rock, 1625 m, N 37°17'470" – E 31°23'695", 01.vii.2010, Çinbilgel 6842; İbradı, Suçikan Location, streamsides, 1259 m, N 37°14'503" – E 31°24'173", 02.vii.2010, Çinbilgel 7068, E. Medit., (G).
	<i>Allium scorodoprasum</i> L. subsp. <i>rotundum</i> (L.) Stearn	İbradı, Sürüçikan Mountain, screes, 1979 m, N 37°12'744" – E 31°19'789", 10.vi.2008, Çinbilgel 3789; Manavgat, Derme Plateau, stony places, 1803 m, N 37°13'054" – E 31°20'219", 30.vi.2009, Çinbilgel 5583; İbradı, Melik Mountain, screes, 2148 m, N 37°16'119" – E 31°23'536", 18.vi.2010, Çinbilgel 6502, (G).
	<i>Allium sphaerocephalon</i> L. subsp. <i>sphaerocephalon</i>	Manavgat, Dermebeli Location, rock, 1920 m, N 37°13'702" – E 31°20'304", 29.vi.2009, Çinbilgel 5524; Manavgat, Kalbur Gözenekleri, rock, 1788 m, N 37°13'259" – E 31°20'585", 05.viii.2009, Çinbilgel 5700; Manavgat, Çiğdem Hill, damp places, 1800 m, Çinbilgel 6467, Euro-Sib., (G).
	* <i>Allium stylosum</i> O.Schwarz	Kovacık Mountain foothills (Hebil Plateau Location), thorny cushion formation, 1771 m, N 37°13'824" – E 31°19'246", 09.vi.2009, Çinbilgel 5271, End., E. Medit., (G).
	* <i>Allium tauricola</i> Boiss.	Manavgat, Kaldırım Mountain (Kızlaroturağı Location), dwarf bush formation, 1710 m, N 37°10'890" – E 31°20'893", 03.vi.2009, Çinbilgel 5136; Manavgat, Kovacık Mountain (Gökçebel Location), thorny cushion formation, 1709 m, N 37°14'032" – E 31°19'421", 09.vi.2009, Çinbilgel 5222; İbradı, Melik Mountain (Hasan Hill), stony slopes, 2079 m, N 37°16'594" – E 31°23'035", 17.vi.2010, Çinbilgel 6417, End., Ir. – Tur., (G).
	<i>Allium vineale</i> L.	İbradı, Toka Plateau (Göçebe Location), rocky places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çinbilgel 3115; Manavgat, Burunçalı Location, slopes, 1846 m, N 37°09'124" – E 31°18'924", 02.vi.2009, Çinbilgel 4981; Manavgat, Burunçalı Location (Aslanboğazi), rocky hills, 1907 m, N 37°08'951" – E 31°19'501", 02.vi.2009, Çinbilgel 5051, (G).
	<i>Galanthus elwesii</i> Hook.f. var. <i>elwesii</i>	Manavgat, Akpınar Plateau, under the <i>Cedrus libani</i> forest, 1731 m, N 37°09'303" – E 31°20'650", 26.iii.2009, Çinbilgel 4536; İbradı, Toka Plateau Breach, <i>Cedrus libani</i> forest clearings – stony places, 1674 m, N 37°13'777" – E 23°38'198", 27.iii.2010, Çinbilgel 5864; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 990 m, N 37°10'768" – E 31°25'396", 05.iii.2011, Çinbilgel 7327, E. Medit., (mt.), (G).
Anacardiaceae		
	<i>Cotinus coggygria</i> Scop.	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7720, (Ph).
	<i>Pistacia palaestina</i> Boiss.	Manavgat, Kızılıpınar Plateau way, Akin – Karahan Location, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çinbilgel 4142; Manavgat, Akpınar Plateau way, forest, 1306 m, N 37°08'484" – E 31°19'182", 11.x.2009, Çinbilgel 5818, E. Medit., (Ph).
	<i>Rhus coriaria</i> L.	İbradı, Söbüova Location (around of Akpınar Location), under the forest, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3534; İbradı, Suçikan Location surroundings, <i>Cedrus libani</i> – <i>Pinus nigra</i> forest, 1271 m, N 37°14'863" – E 31°24'369", 02.vii.2010, Çinbilgel 7041, (Ph).
Apiaceae		
	<i>Anthriscus nemorosa</i> (M. Bieb.) Spreng.	İbradı, Suçikan Location way, under the forest, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2732; İbradı, Ekinlik Location surroundings, <i>Pinus nigra</i> forest (Mixed with <i>Cedrus libani</i>), 1480 m, N 37°13'532" – E 31°23'607", 09.v.2010, Çinbilgel 6153, (H).
	<i>Bunium microcarpum</i> (Boiss.) Freyn & Bornm. ex Freyn subsp. <i>bourgaei</i> (Boiss.) Hedge & Lamond	Manavgat, Kocatop Mountain, windy hills, 2174 m, N 37°12'467" – E 31°18'805", 10.vi.2009, Çinbilgel 5330; İbradı, Melik Mountain (Orta Hill), dwarf bush formation, 2153 m, N 37°16'186" – E 31°22'903", 17.vi.2010, Çinbilgel 6342; İbradı, Suçikan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7078, Ir. – Tur., (G).
	* <i>Bupleurum davisii</i> Snogerup	Manavgat, Akpınar Plateau (Hıncırak Location), rocky debris, 1468 m, N 37°09'337" – E 31°20'412", 01.vi.2009, Çinbilgel 4867; Manavgat, Yalıncaadağ, rocky hills, 2068 m, N 37°11'428" – E 31°18'822", 11.vi.2009, Çinbilgel 5367; İbradı, Melik Mountain (Orta Hill yamacı), debris, 1830 m, N 37°15'886" – E 31°22'533", 17.vi.2010, Çinbilgel 6292, End., E. Medit., (Th).
	<i>Bupleurum falcatum</i> L. subsp. <i>persicum</i> (Boiss.) Koso – Pol.	Manavgat, Çandır Hill, stony hills, 2037 m, N 37°16'626" – E 31°21'029", 11.vi.2008, Çinbilgel 3881; Manavgat, Çandır Hill, windy hills, 1987 m, N 37°14'492" – E 31°21'151", 04.viii.2009, Çinbilgel 5684, (H).
	* <i>Bupleurum pulchellum</i> Boiss. & Heldr.	Manavgat, Dermebeli surroundings, stony places, N 37°13'667" – E 31°20'270", 10.vi.2008, Çinbilgel 3712; Kovacık Mountain foothills (Hebil Plateau Location), thorny cushion formation, 1771 m, N 37°13'824" – E 31°19'246", 09.vi.2009, Çinbilgel 5267; İbradı, Çeşgar Plateau, rock, 1629 m, N 37°17'444" – E 31°23'680", 01.vii.2010, Çinbilgel 6859, End., E. Medit., (Th).

	<i>*Bupleurum subuniflorum</i> Boiss. & Heldr.	Manavgat, Burunçalı Location, rocky places, 1660 m, N 37°08'970" – E 31°18'852", 02.vi.2009, Çinbilgel 4967; İbradı, Melik Mountain (Orta Hill), windy hills, 1948 m, N 37°16'096" – E 31°22'564", 17.vi.2010, Çinbilgel 6308; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest clearings – stony places, 1001 m, N 37°10'801" – E 31°25'318", 02.vii.2010, Çinbilgel 7210, End., E. Medit., (Th).
	<i>Caucalis platycarpus</i> L.	Manavgat, Southern hillsides of Kaldırım Mountain, under the <i>Cedrus libani</i> forest, 1551 m, N 37°09'461" – E 31°20'374", 01.vi.2009, Çinbilgel 4879; İbradı, Çukuryurt Plateau, stony places, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6173; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7032, (Th).
	<i>Eryngium campestre</i> L. var. <i>virens</i> Link	İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest, 1344 m, N 37°15'376" – E 31°24'546", 02.vii.2010, Çinbilgel 7019; İbradı, Eynif Plain, arid meadows, 945 m, N 37°08'943" – E 31°26'971", 26.ix.2010, Çinbilgel 7194, (H).
	<i>Eryngium falcatum</i> F.Delaroche	İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest, 1344 m, N 37°15'376" – E 31°24'546", 02.vii.2010, Çinbilgel 7020, E. Medit., (H).
	<i>*Eryngium trisetum</i> Wörz & H. Duman	Manavgat, Sartaş Location, stony hillsides, 2074 m, N 37°15'015" – E 31°20'984", 11.vi.2008, Çinbilgel 3916; Manavgat, Kızılpınar Plateau, dwarf bush formation, 1782 m, N 37°14'186" – E 31°20'257", 03.viii.2009, Çinbilgel 5599; İbradı, Melik Plateau, meadows, 1446 m, N 37°15'954" – E 31°24'591", 01.vii.2010, Çinbilgel 6764, End., (H).
	<i>Falcaria vulgaris</i> Bernh.	İbradı, Güzle Plateau, moist stony places, 1294 m, N 37°10'219" – E 31°23'663", 20.iv.2008, Çinbilgel 2907, Euro-Sib., (H).
	<i>Ferula orientalis</i> L.	İbradı, Gevenli Location, streams'nin kenarları – stony places, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3526; İbradı, Cevizliedere Location, stony places, 1131 m, N 37°13'469" – E 31°24'205", 08.vi.2011, Çinbilgel 7963, Ir. – Tur., (H).
	<i>Ferulago trachycarpa</i> Boiss.	Manavgat, Kovacık Mountain (Uzunçaşır Location), screes, 1939 m, N 37°13'639" – E 31°20'047", 29.vi.2009, Çinbilgel 5499; İbradı, Çeşgar Plateau, rock, 1625 m, N 37°17'470" – E 31°23'695", 01.vii.2010, Çinbilgel 6843, E. Medit., (H).
	<i>Laser trilobum</i> (L.) Borkh.	İbradı, Suçikan Location way, under the forest, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2779; İbradı, Suçikan Location way, under the forest – stony rocky places, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4419; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7038, (H).
	<i>Opopanax hispidus</i> (Friv.) Griseb.	İbradı, Melik Mountain (Hasan Hill), stony hillsides, 2000 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4344, (H).
	<i>Orlaya daucoides</i> (L.) Greuter	İbradı, Tahtalı Location, stony places, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4397, Medit., (Th).
	<i>Peucedanum alpinum</i> (Sieb. ex Schult.) B.L.Burt & P.H.Davis	Manavgat, Burunçalı Location (Aslanboğazi), stony places, 1687 m, N 37°08'939" – E 31°19'500", 02.vi.2009, Çinbilgel 5021, E. Medit., (H).
	<i>*Pimpinella ibradiensis</i> Çinbilgel, Eren, H.Duman & Gökçeoğlu	İbradı, Toka Plateau, moist stony places, 1500 m, N 37°14'349" – E 31°22'627", 08.v.2010, Çinbilgel 6047 & Eren; İbradı, Aşağıtoka Plateau, moist stony places, 1527 m, N 37°12'804" – E 31°22'579", 16.vi.2010, Çinbilgel 6233; İbradı, Aşağıtoka Plateau, stony places, 1527 m, N 37°13'253" – E 31°22'503", 02.vii.2011, Çinbilgel 7975 & Eren, End., E. Medit., (H).
	<i>Pimpinella tragiium</i> Vill. subsp. <i>lithophila</i> (Schischk.) Tutin	Manavgat, Çiğdem Hill, dwarf bush formation, 1904 m, N 37°14'197" – E 31°20'828", 04.viii.2009, Çinbilgel 5659; İbradı, Aşağıtoka Plateau, stony places, 1531 m, N 37°12'882" – E 31°22'682", 16.vi.2010, Çinbilgel 6245; İbradı, Çeşgar Plateau Breach, stony places, 1664 m, N 37°17'389" – E 31°24'270", 01.vii.2010, Çinbilgel 6810, (H).
	<i>Scandix australis</i> L. subsp. <i>grandiflora</i> (L.) Thell.	İbradı, Akmuar Location, stony places, 1044 m, N 37°11'687" – E 31°24'880", 28.iii.2010, Çinbilgel 5952, (Th).
	<i>Scandix iberica</i> M.Bieb.	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest – stony places, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3461; Manavgat, Kızılpınar Plateau, rocky places, 1811 m, N 37°13'847" – E 31°19'648", 25.v.2008, Çinbilgel 3472/a; Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, Putraklı Location, stony places, 734 m, N 37°12'249" – E 31°16'454", 27.iii.2011, Çinbilgel 7563, (Th).
	<i>Scandix pecten-veneris</i> L.	Manavgat, Akpınar Plateau, under the <i>Cedrus libani</i> forest, 1591 m, N 37°10'194" – E 31°21'483", 19.v.2009, Çinbilgel 4745; İbradı, Toka Plateau Breach, stony slopes, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5976; İbradı, Around of Toka Plateau Breach, <i>Cedrus libani</i> – <i>Pinus nigra</i> forest, 1679 m, N 37°13'808" – E 31°23'202", 21.v.2011, Çinbilgel 7670, (Th).
	<i>*Seseli gummiferum</i> Pall. ex Sm. subsp. <i>corymbosum</i> (Boiss. & Heldr.) P.H.Davis	Manavgat, Burunçalı Location, rock, 1693 m, N 37°09'006" – E 31°18'835", 02.vi.2009, Çinbilgel 4962; Manavgat, Dermebelı Location, rock, 1944 m, N 37°13'711" – E 31°20'308", 29.vi.2009, Çinbilgel 5525; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1502 m, N 37°15'548" – E 31°24'365", 30.vi.2010, Çinbilgel 6658, End., E. Medit., (Ch).

	<i>Smyrniium connatum</i> Boiss. & Kotschy	İbradı, Suçikan Location way, stony places, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2783; İbradı, Melik Plateau, stony slopes, 1644 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4169; İbradı, Çeşgar Plateau, stony slopes, 1615 m, N 37°17'577" – E 31°23'727", 17.vi.2008, Çinbilgel 4214, E. Medit., (H).
	* <i>Tordylium ketenoglui</i> H. Duman & A. Duran	Manavgat, Around of Burmahan Neighbourhood, Öküztaşı Location, rock, 700 m, N 37°10'312" – E 31°16'923", 06.vi.2011, Çinbilgel 7819; Manavgat, Çay Neighbourhood, Taşhyatak Location, Pinus brutia forest, 960 m, N 37°12'764" – E 31°16'610", 07.vi.2011, Çinbilgel 7930, End., E. Medit., (Th).
	<i>Torilis arvensis</i> (Huds.) Link subsp. <i>elongata</i> (Hoffmanns. & Link) Cannon	Manavgat, Arıçca Location (cutting area), Pinus brutia forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4107; İbradı, Melik Plateau, forest clearings, 1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6959; İbradı, Melik Plateau surroundings, Pinus nigra – Cedrus libani forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7037; Medit., (Th).
	<i>Torilis leptophylla</i> (L.) Rchb.f.	Manavgat, Dügünük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4067; İbradı, Around of Yirme Hill, Pinus nigra forest, 1600 m, N 37°14'248" – E 31°22'862", 08.v.2010, Çinbilgel 6016; İbradı, Akpınar Location, stony places, 1073 m, N 37°11'976" – E 31°24'786", 02.vii.2010, Çinbilgel 7152, (Th).
Apocynaceae		
	<i>Cionura erecta</i> (L.) Griseb.	İbradı, Akmuar Location, roadsides, 1018 m, N 37°11'513" – E 31°24'932", 18.vi.2008, Çinbilgel 4442; Manavgat, Burmahan Neighbourhood – Akpınar Plateau way, roadsides, 798 m, N 37°09'483" – E 31°17'167", 11.x.2009, Çinbilgel 5812, E. Medit., (Ph).
	<i>Nerium oleander</i> L.	Manavgat, Kızılıpınar Plateau way, Göcek Location, roadsides, 600 m, 13.vi.2008, Çinbilgel 4152; Manavgat, Burmahan Neighbourhood surroundings, forest opennings, 699 m, N 37°10'800" – E 31°16'736", 26.iii.2011, Çinbilgel 7425; Manavgat, Burmahan Neighbourhood, around of Erenli Location, Pinus brutia forest, 723 m, N 37°09'738" – E 31°16'983", 26.iii.2011, Çinbilgel 7461, Medit., (Ph).
	<i>Vinca herbacea</i> Waldst. & Kit.	İbradı, Güzle Plateau – Hillsides of Yaylacık Mountain, moist stony places, 1430 m, N 37°10'227" – E 31°23'913", 20.iv.2008, Çinbilgel 2880; Manavgat, Kaldırım Mountain, Cedrus libani forest clearings, 1753 m, N 37°09'809" – E 31°20'339", 01.vi.2009, Çinbilgel 4853; İbradı, Melik Plateau, forest clearings – stony places, 1406 m, N 37°15'466" – E 31°24'469", 30.vi.2010, Çinbilgel 6688, (H).
	<i>Vincetoxicum fuscatum</i> Rchb.f. subsp. <i>fuscatum</i>	İbradı, Melik Plateau Location, stony slopes, 1644 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4163; İbradı, Toka Plateau way, Belova Location, Cedrus libani forest clearings, 1568 m, N 37°13'549" – E 31°23'433", 03.vii.2011, Çinbilgel 7999, (H).
	<i>Vincetoxicum tmoleum</i> Boiss.	Manavgat, Kızılıpınar Plateau way, Kavaklı Location, stony places, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4040; Manavgat, Sürüçikan Mountain, dwarf bush formation, 2107 m, N 37°12'335" – E 31°19'690", 30.vi.2009, Çinbilgel 5556; İbradı, Melik Mountain foothills, rock crevices, 1809 m, N 37°15'464" – E 31°23'088", 18.vi.2010, Çinbilgel 6483, İr. – Tur., (H).
Araceae		
	<i>Arum dioscoridis</i> Sm. var. <i>dioscoridis</i>	İbradı, Söbüova Location (around of Akpınar Location), under the forest – stony places, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3561, Medit., (G).
Araliaceae		
	<i>Hedera helix</i> L.	Manavgat, Dutluca Plateau, under the rock – stony places, 1282 m, N 37°13'785" – E 31°18'213", 13.vi.2008, Çinbilgel 4056; Manavgat, Burmahan Neighbourhood, Erenli Location, Pinus brutia forest – watersides, 729 m, N 37°09'926" – E 31°17'026", 11.x.2009, Çinbilgel 5799; İbradı, Cevizlidere Location, under the rock, 1139 m, N 37°13'357" – E 31°24'239", 05.x.2010, Çinbilgel 7272, (Ph).
Aristolochiaceae		
	* <i>Aristolochia lycica</i> P.H.Davis & M.S.Khan	İbradı, Between Gavur Breach and Etülgenboğazı, Pinus brutia forest, 1186 m, N 37°09'074" – E 31°25'116", 20.iv.2008, Çinbilgel 2971, End., E. Medit., (H).
	<i>Aristolochia paecilantha</i> Boiss.	İbradı, Tahtalı Location, under the Pinus brutia forest, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4404; İbradı, Akmuar Location, stony places, 1073 m, N 37°11'976" – E 31°24'786", 02.vii.2010, Çinbilgel 7154; Manavgat, Genger Location, Pinus brutia forest, 1150 m, N 37°09'242" – E 31°17'835", 06.vi.2011, Çinbilgel 7870, E. Medit., (H).
Asparagaceae		
	<i>Asparagus acutifolius</i> L.	Manavgat, Çay Neighbourhood, Pinus brutia forest, 688 m, N 37°12'133" – E 31°16'452", 27.iii.2011, Çinbilgel 7478, Medit., (Ch).
	* <i>Bellevalia tauri</i> Feinbrun	İbradı, Suçikan Location way, damp places, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2780; Manavgat, Kepez Plateau, wet places, 1648 m, N 37°10'838" – E 31°20'012", 03.vi.2009, Çinbilgel 5145; İbradı, Toka Plateau, meadows, 1514 m, N 37°13'924" – E 31°22'564", 09.v.2010, Çinbilgel 6091, End., E. Medit., (G).

	<i>*Hyacinthella heldreichii</i> (Boiss.) Chouard	İbradı, Güzle Plateau, stony places, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2818; İbradı, Güzle Plateau (Mezarlı Location), under the forest, 1287 m, N 37°10'193" – E 31°23'650", 18.iv.2009, Çinbilgel 4568; İbradı, Boğazyurt Location, stony places, 1048 m, N 37°11'404" – E 31°25'002", 28.iii.2010, Çinbilgel 5960, End., E. Medit., (G).
	<i>*Muscari anatolicum</i> Cowley & Özhatay	İbradı, Güzle Plateau (Mezarlı Location), stony places, 1287 m, N 37°10'193" – E 31°23'650", 18.iv.2009, Çinbilgel 4558; Manavgat, Kepezbeli Location, stony hillsides, 1641 m, N 37°10'300" – E 31°21'722", 19.v.2009, Çinbilgel 4720; İbradı, Aşağıtoka Plateau, stony places, 1519 m, N 37°14'106" – E 31°22'468", 28.iii.2010, Çinbilgel 5901, End., (G).
	<i>Muscari armeniacum</i> Leichtlin ex Baker	Manavgat, Kepez Plateau (İncayırı Location), snow patches, 1674 m, N 37°10'569" – E 31°21'292", 03.v.2009, Çinbilgel 4724; İbradı, Toka Plateau, meadows, 1514 m, N 37°13'924" – E 31°22'564", 09.v.2010, Çinbilgel 6090; İbradı, Gevenli Location, moist stony places, 1100 m, N 37°11'993" – E 31°24'440", 05.iii.2011, Çinbilgel 7355, (G).
	<i>*Muscari aucheri</i> (Boiss.) Baker	İbradı, Melik Mountain, stony places, 2136 m, N 37°15'764" – E 31°23'547", 18.v.2008, Çinbilgel 3318, End., (G).
	<i>*Muscari bourgaei</i> Baker	İbradı, Yirme Hill, rocky hills, 1820 m, N 37°14'717" – E 31°23'178", 18.v.2008, Çinbilgel 3242; Manavgat, Kocatop Mountain, snow patches, 2080 m, N 37°12'599" – E 31°18'884", 10.vi.2009, Çinbilgel 5280; İbradı, Toka Plateau, moist stony places, 1521 m, N 37°14'154" – E 31°22'670", 28.iii.2010, Çinbilgel 5889, End., E. Medit. (mt.), (G).
	<i>Muscari comosum</i> (L.) Mill.	İbradı, Akmuar Location, rock, 1024 m, N 37°11'720" – E 31°24'844", 19.v.2011, Çinbilgel 7604; İbradı, Toka Plateau Breach, meadows, 1676 m, N 37°13'774" – E 31°23'168", 21.v.2011, Çinbilgel 7667, Medit., (G).
	<i>Muscari neglectum</i> Guss. ex Ten.	İbradı, Güzle Plateau (Mezarlı Location), stony places, 1287 m, N 37°10'193" – E 31°23'650", 18.iv.2009, Çinbilgel 4564; İbradı, Boğazyurt Location, stony places, 1048 m, N 37°11'404" – E 31°25'002", 28.iii.2010, Çinbilgel 5959; Manavgat, Çay Neighbourhood, Kızılpınar Plateau way (Düğnük Plateau), stony places, 1163 m, N 37°13'385" – E 31°17'716", 27.iii.2011, Çinbilgel 7507, (G).
	<i>Muscari tenuiflorum</i> Tausch	İbradı, Gevenli Location, <i>Under the Pinus nigra forest</i> , 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3521; Manavgat, Kızılpınar Plateau way, Kavaklı Location, stony places, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4053; Manavgat, Burunçalı Location, rocky places, 1810 m, N 37°09'072" – E 31°18'873", 02.vi.2009, Çinbilgel 5066, (G).
	<i>Muscari weissii</i> Freyn	İbradı, Akmuğar Location, damp places, 1032 m, N 37°11'838" – E 31°24'840", 19.v.2008, Çinbilgel 3583, E. Medit., (G).
	<i>*Ornithogalum alpigenum</i> Stapf	İbradı, Melik Mountain, snow patches, 2264 m, N 37°16'041" – E 31°23'664", 18.vi.2010, Çinbilgel 6530, End., E. Medit., (G).
	<i>Ornithogalum armeniacum</i> Baker	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings, 1701 m, N 37°13'682" – E 31°23'231", 19.v.2008, Çinbilgel 3408; İbradı, Çukuryurt Plateau, stony places, 1299 m, N 37°12'181" – E 31°23'938", 09.v.2010, Çinbilgel 6164; İbradı, Aşağıtoka Plateau, wet meadows, 1514 m, N 37°13'772" – E 31°22'587", 08.vi.2011, Çinbilgel 7954, E. Medit., (G).
	<i>Ornithogalum neurostegium</i> Boiss. & C.I.Blanche ex Boiss.	Manavgat, Susuzkır Hill surroundings, stony places, 1666 m, N 37°16'797" – E 31°21'755", 12.vi.2008, Çinbilgel 3984; İbradı, Toka Plateau, meadows, 1500 m, N 37°14'367" – E 31°22'605", 08.v.2010, Çinbilgel 6038; İbradı, Aşağıtoka Plateau, stony places, 1508 m, N 37°13'377" – E 31°22'478", 02.vii.2011, Çinbilgel 7970, (G).
	<i>Ornithogalum oligophyllum</i> E.D.Clarke	İbradı, Melik Mountain (Hillsides of Çeşgar Hill), stony places, 2000 m, N 37°15'593" – E 31°23'351", 18.v.2008, Çinbilgel 3310; İbradı, Yaylacık Mountain, 1481 m, N 37°10'109" – E 31°24'830", 19.iv.2009, Çinbilgel 4601; İbradı, Aşağıtoka Plateau, meadows, 1544 m, N 37°13'888" – E 31°22'352", 16.vi.2010, Çinbilgel 6258, (G).
	<i>Ornithogalum orthophyllum</i> Ten. subsp. <i>kochii</i> (Parl.) Maire & Weiller	İbradı, Aşağıtoka Plateau, stony places, 1555 m, N 37°13'162" – E 31°22'666", 20.v.2011, Çinbilgel 7649, (G).
	<i>Ornithogalum pyrenaicum</i> L.	Manavgat, Düğnük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4078, (G).
	<i>Ornithogalum wiedemannii</i> Boiss. var. <i>wiedemannii</i>	Manavgat, Akpınar Plateau (Yumrutaş Location), under the mixed <i>Cedrus libani</i> forest, 1514 m, N 37°09'738" – E 31°21'024", 19.v.2009, Çinbilgel 4764, (G).
	<i>Prospero autumnale</i> (L.) Speta	İbradı, Eynif Plain, arid meadows, 945 m, N 37°08'943" – E 31°26'971", 26.ix.2010, Çinbilgel 7191, Medit., (G).
	<i>Ruscus aculeatus</i> L.	(Ph), (Observation).
	<i>Scilla bifolia</i> L.	Manavgat, Kepezbeli Location, stony slopes, 1674 m, N 37°10'329" – E 31°21'582", 03.v.2009, Çinbilgel 4709; İbradı, Aşağıtoka Plateau, meadows – stony places, 1521 m, N 37°13'187" – E 31°22'678", 27.iii.2010, Çinbilgel 5869; Manavgat, Akpınar Plateau, under the mixed <i>Cedrus libani</i> forest, 1385 m, N 37°09'485" – E 31°20'778", 26.iii.2011, Çinbilgel 7459, Medit., (G).
Asteraceae		
	<i>Achillea arabica</i> Kotschy	Manavgat, Semai Plateau, <i>Cedrus libani</i> forest – stony places, 1586 m, N 37°14'660" – E 31°19'747", 12.vi.2008, Çinbilgel 4029; İbradı, Melik Plateau,

		forest clearings – stony places, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6566, Ir. – Tur., (H).
	<i>*Achillea formosa</i> (Boiss.) Sch.Bip. subsp. <i>formosa</i>	İbradı, Melik Mountain, stony slopes, 1880 m, N 37°14'957" – E 31°23'372", 18.v.2008, Çinbilgel 3270; İbradı, Melik Mountain (Hasan Hill), slopes, 2000 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4348, End., E. Medit., (H).
	<i>*Achillea kotschyi</i> Boiss. subsp. <i>canescens</i> Bässler	İbradı, around of Çeşgar Plateau Breach, dwarf bush formation, 1588 m, N 37°17'590" – E 31°23'768", 01.vii.2010, Çinbilgel 6825, End., Medit. (mt.), (H).
	<i>*Achillea nobilis</i> L. subsp. <i>densissima</i> (O.Schwarz ex Bässler) Hub. – Mor.	İbradı, Melik Plateau, forest clearings – stony places, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6568; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çinbilgel 6707; İbradı, Melik Plateau, stony places, 1515 m, N 37°16'380" – E 31°24'492", 01.vii.2010, Çinbilgel 6928, End., Medit., (H).
	<i>Achillea nobilis</i> L. subsp. <i>neilreichii</i> (A.Kern) Velen	İbradı, Melik Plateau surroundings, streamsides, 1346 m, N 37°15'389" – E 31°24'545", 02.vii.2010, Çinbilgel 7016; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1006 m, N 37°11'054" – E 31°25'243", 03.vii.2011, Çinbilgel 8040; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 983 m, N 37°10'973" – E 31°25'293", 03.vii.2011, Çinbilgel 8045, 3566/a, Euro-Sib., (H).
	<i>Achillea santolinoides</i> Lag. subsp. <i>wilhelmsii</i> (K.Koch) Greuter	İbradı, Çeşgar Plateau, stony slopes, 1626 m, N 37°17'559" – E 31°23'696", 17.vi.2008, Çinbilgel 4257; Manavgat, Kovacık Mountain foothills (Gökçebel Location), thorny cushion formation, 1687 m, N 37°14'038" – E 31°19'336", 09.vi.2009, Çinbilgel 5189; İbradı, Melik Plateau, dwarf bush formation, 1728 m, N 37°16'359" – E 31°24'498", 01.vii.2010, Çinbilgel 6932, Ir. – Tur., (H).
	<i>Achillea setacea</i> Waldst. & Kit.	İbradı, Çeşgar Plateau, stony slopes, 1626 m, N 37°17'559" – E 31°23'696", 17.vi.2008, Çinbilgel 4255; İbradı, Melik Plateau, stony places, 1515 m, N 37°16'380" – E 31°24'492", 01.vii.2010, Çinbilgel 6927, Euro-Sib., (H).
	<i>*Anthemis aciphylla</i> Boiss. var. <i>aciphylla</i>	İbradı, Eynif Plain, arid meadows, 949 m, N 37°08'004" – E 31°27'963", 19.v.2011, Çinbilgel 7571, End., E. Medit., (H).
	<i>Anthemis cotula</i> L.	İbradı, around of Çukuryurt Plateau, stony places, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6174; İbradı, Melik Plateau surroundings, <i>Under the Pinus nigra forest</i> , 1304 m, N 37°15'106" – E 31°24'488", 02.vii.2010, Çinbilgel 6982; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7176, (Th).
	<i>Anthemis cretica</i> L. subsp. <i>albida</i> (Boiss.) Grierson	İbradı, Yukarıtoka Plateau (Between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3030; İbradı, Melik Mountain (Hasan Hill), rocky slopes, 2020 m, N 37°16'897" – E 31°23'181", 18.vi.2008, Çinbilgel 4303; İbradı, Lower Toka Plateau, dwarf bush formation, 1526 m, N 37°13'831" – E 31°22'356", 20.v.2011, Çinbilgel 7656, (Ch).
	<i>Anthemis cretica</i> L. subsp. <i>anatolica</i> (Boiss.) Grierson	İbradı, Toka Plateau (Göçebe Location), stony places, 1551 m, N 37°14'763" – E 31°22'308", 17.v.2008, Çinbilgel 3144; Manavgat, Kovacık Mountain (around of Dermebeli Location), dwarf bush formation, 1867 m, N 37°13'751" – E 31°20'136", 29.vi.2009, Çinbilgel 5463; İbradı, Melik Plateau, <i>Pinus nigra forest</i> , 1462 m, N 37°15'665" – E 31°24'474", 30.vi.2010, Çinbilgel 6606, (Ch).
	<i>Anthemis cretica</i> L. subsp. <i>leucanthemoides</i> (Boiss.) Grierson	İbradı, Dikenli Location, forest clearings, 1674 m, N 37°14'298" – E 31°23'004", 18.v.2008, Çinbilgel 3201, (Ch).
	<i>*Anthemis pauciloba</i> Boiss. var. <i>pauciloba</i>	İbradı, Melik Plateau Location, stony slopes, 1644 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4171; Manavgat, Kovacık Mountain, Gökçebel Location, thorny cushion formation, 1709 m, N 37°14'032" – E 31°19'421", 09.vi.2009, Çinbilgel 5221; İbradı, Melik Mountain (Orta Hill), stony hills, 2095 m, N 37°16'189" – E 31°22'824", 17.vi.2010, Çinbilgel 6352, End., E. Medit., (H).
	<i>Anthemis pseudocotula</i> Boiss.	İbradı, Between Akpınar Location and Suçikan Location, stony places, 1091 m, N 37°13'075" – E 31°24'456", 18.vi.2008, Çinbilgel 4429; İbradı, Melik Plateau, dwarf bush formation, 1665 m, N 37°17'156" – E 31°24'327", 01.vii.2010, Çinbilgel 6925; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia forest clearings</i> , 655 m, N 37°11'683" – E 31°16'103", 06.vi.2011, Çinbilgel 7752, (Th).
	<i>Artemisia absinthium</i> L.	İbradı, Toka Plateau (Göçebe Location), stony places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çinbilgel 3107; İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani forest</i> (mixed with <i>Pinus nigra</i>), 1660 m, N 37°13'683" – E 31°23'340", 09.v.2010, Çinbilgel 6065; İbradı, Yukarıtoka Plateau, dwarf bush formation, 1771 m, N 37°15'406" – E 31°22'669", 18.vi.2010, Çinbilgel 6547, (Ch).
	<i>Bellis perennis</i> L.	Manavgat, Akpınar Plateau way, <i>Pinus brutia forest</i> , 1000 m, N 37°08'705" – E 31°19'909", 26.iii.2009, Çinbilgel 4538; Manavgat, Kepez Plateau (İncayırı Location), wet places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5856; İbradı, Boğazyurt Location, <i>Juniperus excelsa forest</i> , 980 m, N 37°10'766" – E 31°25'403", 05.iii.2011, Çinbilgel 7333, Euro-Sib., (H).
	<i>Carduus argentatus</i> L.	İbradı, Suçikan Location way, stony rock places, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4415; İbradı, Cevzlidere Location, stony

		places, 1149 m, N 37°13'475" – E 31°24'190", 02.vii.2010, Çinbilgel 7142; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7735, E. Medit., (Th).
	<i>Carduus nutans</i> L. subsp. <i>nutans</i>	İbradı, around of Yirme Hill, stony slopes, 1783 m, N 37°14'667" – E 31°23'097", 18.v.2008, Çinbilgel 3195; Manavgat, Yalıncaadağ, dwarf bush formation, 2039 m, N 37°11'346" – E 31°18'682", 11.vi.2009, Çinbilgel 5407; İbradı, Akmuar Location, stony places, 1042 m, N 37°11'845" – E 31°24'839", 02.vii.2010, Çinbilgel 7157, (H).
	<i>Carduus pycnocephalus</i> L. subsp. <i>albidus</i> (M.Bieb.) Kazmi	İbradı, Akmuar Location, stony places, 1033 m, N 37°12'323" – E 31°24'784", 18.vi.2008, Çinbilgel 4436; Manavgat, Kepez Plateau (İnçayırı Location), wet places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5854; İbradı, Suçikan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7076, (Th).
	<i>Carduus pycnocephalus</i> L. subsp. <i>pycnocephalus</i>	İbradı, Eynif Plain, meadows, 941 m, N 37°08'989" – E 31°26'561", 19.v.2011, Çinbilgel 7575, Medit., (Th).
	<i>Carlina corymbosa</i> L.	Manavgat, around of Burmahan Neighbourhood, <i>Pinus brutia</i> forest, 695 m, N 37°11'319" – E 31°16'296", 06.vi.2011, Çinbilgel 7782, Medit., (H).
	<i>Carthamus dentatus</i> (Forssk.) Vahl.	İbradı, Boğazyurt-Akmuar Location, <i>Juniperus excelsa</i> forest, 964 m, N 37°11'201" – E 31°25'141", 18.vi.2008, Çinbilgel 4448; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7185, (Th).
	<i>Centaurea babylonica</i> (L.) L.	Manavgat, around of Burmahan Neighbourhood, roadsides, 630 m, 20.vii.2008, Çinbilgel 4533, E. Medit., (H).
	* <i>Centaurea calolepis</i> Boiss.	İbradı, Tahtalı Location, stony places, 1087 m, N 37°12'727" – E 31°24'595", 02.vii.2010, Çinbilgel 7122, End., E. Medit., (H).
	* <i>Centaurea drabifolia</i> Sibth. & Sm. subsp. <i>austro-occidentalis</i> Wagenitz	İbradı, Tahtalı Location, <i>Under the Pinus nigra forest</i> , 1078 m, N 37°12'750" – E 31°24'574", 03.vii.2011, Çinbilgel 8019, End., (H), (Conf. Eren).
	* <i>Centaurea drabifolia</i> Sibth. & Sm. subsp. <i>cappadocica</i> (DC.) Wagenitz	Manavgat, Çandır Hill, stony hills, 2037 m, N 37°16'626" – E 31°21'029", 11.vi.2008, Çinbilgel 3884; Manavgat, Kocatop Mountain (Koca Hill = Köygöründüğü Location), windy hills, 2150 m, N 37°12'346" – E 31°18'633", 10.vi.2009, Çinbilgel 5341; İbradı, Melik Mountain (Orta Hill), stony hills, 2095 m, N 37°16'189" – E 31°22'824", 17.vi.2010, Çinbilgel 6349, End., (H), (Conf. Eren).
	* <i>Centaurea drabifolia</i> Sibth. & Sm. subsp. <i>floccosa</i> (Boiss.) Wagenitz & Greuter	Manavgat, Dermebeli Location (Artaşı), on the rocks and stony places, 1874 m, N 37°13'519" – E 31°20'317", 10.vi.2008, Çinbilgel 3805; Manavgat, Dermebeli Location (Artaşı), rock, 1885 m, N 37°13'490" – E 31°20'297", 19.vii.2008, Çinbilgel 4467; Manavgat, Dermebeli Location, rock, 1929 m, N 37°13'694" – E 31°20'299", 29.vi.2009, Çinbilgel 5521, End., (H), (Conf. Eren).
	<i>Centaurea iberica</i> Trev. ex Sprengel	İbradı, Aşağıtoka Plateau, stony places, 1513 m, N 37°13'794" – E 31°22'474", 16.vi.2010, Çinbilgel 6249, (H).
	* <i>Centaurea solstitialis</i> L. subsp. <i>pyracantha</i> (Boiss.) Wagenitz	Manavgat, Çay Neighbourhood, Kızılıpınar Plateau way, streamsides, 628 m, 05.viii.2009, Çinbilgel 5739, End., E. Medit., (Th).
	<i>Centaurea solstitialis</i> L. subsp. <i>solstitialis</i>	İbradı, Melik Plateau, stony places, 1387 m, N 37°15'533" – E 31°24'528", 30.vi.2010, Çinbilgel 6725; İbradı, Melik Plateau, openings – stony places, 1387 m, N 37°15'533" – E 31°24'528", 30.vi.2010, Çinbilgel 6736, (Th).
	* <i>Centaurea urvillei</i> DC. subsp. <i>hayekiana</i> Wagenitz	İbradı, Yirme Hill, stony places, 1820 m, N 37°14'717" – E 31°23'178", 18.v.2008, Çinbilgel 3250; Manavgat, Düğnük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4086; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çinbilgel 6721, End., Ir. – Tur., (H).
	* <i>Centaurea urvillei</i> DC. subsp. <i>stepposa</i> Wagenitz	İbradı, between Akpınar Location and Suçikan Location, stony places, 1091 m, N 37°13'075" – E 31°24'456", 18.vi.2008, Çinbilgel 4425; İbradı, Akmuar Location, <i>Pinus nigra</i> forest clearings, 1047 m, N 37°11'563" – E 31°24'882", 02.vii.2010, Çinbilgel 7164, End., Ir. – Tur., (H).
	<i>Centaurea virgata</i> Lam.	İbradı, Aşağıtoka Plateau, stony places, 1534 m, N 37°12'825" – E 31°22'485", 16.vi.2010, Çinbilgel 6225, Ir. – Tur., (Ch).
	<i>Chondrilla juncea</i> L.	Manavgat, Kızılıpınar Plateau, dwarf bush formation, 1757 m, N 37°14'073" – E 31°19'973", 10.x.2009, Çinbilgel 5772; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest clearings – stony places, 1001 m, N 37°10'801" – E 31°25'318", 02.vii.2010, Çinbilgel 7214; İbradı, Melik Plateau, stony slopes, 1374 m, N 37°15'565" – E 31°24'628", 05.x.2010, Çinbilgel 7281, (H).
	* <i>Chrysophthalmum dichotomum</i> Boiss. & Heldr.	Manavgat, Derme Plateau, moist stony places, 1744 m, N 37°12'867" – E 31°20'405", 19.vii.2008, Çinbilgel 4472; Manavgat, Derme Plateau (Zahmedin Plateau Location), damp places, 1761 m, N 37°12'348" – E 31°20'456", 20.vii.2008, Çinbilgel 4520, End., E. Medit., (Ch).
	<i>Cichorium intybus</i> L.	Manavgat, Kepez Plateau (İnçayırı Location), wet places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5846; İbradı, Melik Plateau, openings – stony places, 1387 m, N 37°15'533" – E 31°24'528", 30.vi.2010, Çinbilgel 6737; İbradı, Gevenli Location, <i>Pinus brutia</i> forest clearings, 1071 m, N 37°12'300" – E 31°24'793", 03.vii.2011, Çinbilgel 8011, (H).

	<i>Cirsium leucocephalum</i> (Willd.) Spreng. subsp. <i>leucocephalum</i>	Manavgat, Kızılıpınar Plateau, dwarf bush formation, 1782 m, N 37°14'186" – E 31°20'257", 03.viii.2009, Çinbilgel 5603; İbradı, Yukarıtoka Plateau, stony places, 1525 m, N 37°14'271" – E 31°22'440", 02.vii.2010, Çinbilgel 7259; İbradı, Melik Plateau, open places, 1461 m, N 37°15'979" – E 31°24'568", 05.x.2010, Çinbilgel 7298, Ir. – Tur., (H).
	<i>Cnicus benedictus</i> L.	Manavgat, Çay Neighbourhood, Kızılıpınar Plateau way, Kocaarmut Location, <i>Pinus brutia</i> forest clearings – stony places, 709 m, N 37°12'174" – E 31°16'360", 27.iii.2011, Çinbilgel 7552, (Th).
	* <i>Cota pestalozzae</i> Boiss.	İbradı, Güzle Plateau – Yaylacık Mountain, damp places, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çinbilgel 2902; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4641; İbradı, Akmuar Location, Juniper Forest clearings, 1036 m, N 37°11'773" – E 31°24'864", 21.v.2011, Çinbilgel 7676, End., E. Medit., (Th).
	<i>Crepis commutata</i> (Spreng.) Greuter	İbradı, Melik Plateau, dwarf bush formation, 1523 m, N 37°16'395" – E 31°24'482", 01.vii.2010, Çinbilgel 6783; İbradı, Çeşgar Plateau Breach, stony places, 1618 m, N 37°16'826" – E 31°24'504", 01.vii.2010, Çinbilgel 6792, (Th).
	<i>Crepis foetida</i> L. subsp. <i>foetida</i>	Manavgat, Argıçca Breach Location, degraded <i>Pinus brutia</i> forest, 1090 m, N 37°13'042" – E 31°17'108", 13.vi.2008, Çinbilgel 4096; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7714, (Th).
	* <i>Crepis macropus</i> Boiss. & Heldr.	Manavgat, Argıçca Location, <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4132; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest clearing, 1486 m, N 37°16'052" – E 31°24'647", 05.x.2010, Çinbilgel 7314, End., Ir. – Tur., (H).
	<i>Crepis micrantha</i> Czerep.	İbradı, Toka Plateau (Göçebe Location), stony places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çinbilgel 3106; İbradı, Akpınar Plateau (Yumrutaş Location), mixed <i>Cedrus libani</i> forest, 1520 m, N 37°09'840" – E 31°21'028", 19.v.2009, Çinbilgel 4782; İbradı, Melik Mountain, dwarf bush-thorny cushion formation, 1882 m, N 37°15'510" – E 31°23'322", 18.vi.2010, Çinbilgel 6489, (Th).
	<i>Crepis reuterana</i> Boiss. & Heldr. subsp. <i>reuterana</i>	İbradı, Gevenli Location, under the <i>Pinus brutia</i> forest, 1108 m, N 37°12'316" – E 31°24'613", 17.v.2008, Çinbilgel 3025; İbradı, Tahtalı Location, under the forest, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4402; İbradı, Akmuar Location, open places, 1073 m, N 37°11'976" – E 31°24'786", 02.vii.2010, Çinbilgel 7150, E. Medit., (H).
	<i>Crepis sancta</i> (L.) Bornm.	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3459, (Th).
	<i>Cyanus thirkei</i> (Sch.Bip.) Holub	Manavgat, Eşekbeli Location, <i>Pinus nigra</i> forest, 1612 m, N 37°14'867" – E 31°19'864", 07.vi.2011, Çinbilgel 7903, E. Medit., (H).
	<i>Cyanus triumfettii</i> (All.) Dostál ex Á.Löve & D.Löve subsp. <i>triumfettii</i>	İbradı, Çeşgar Plateau, stony slopes, 1659 m, N 37°17'402" – E 31°24'250", 17.vi.2008, Çinbilgel 4243; Manavgat, Kaldırım Mountain (Kızlaroturağı Location), dwarf bush formation, 1710 m, N 37°10'890" – E 31°20'893", 03.vi.2009, Çinbilgel 5134; İbradı, Melik Mountain (Orta Hill), windy hills, 2175 m, N 37°16'298" – E 31°23'052", 17.vi.2010, Çinbilgel 6375, (H).
	<i>Doronicum orientale</i> Hoffm.	İbradı, around of Güzle Plateau, under the forest, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2813; Manavgat, Akpınar Plateau, under the <i>Cedrus libani</i> forest, 1480 m, N 37°09'593" – E 31°21'076", 19.v.2009, Çinbilgel 4811; İbradı, Toka Plateau way, <i>Pinus nigra</i> forest, 1662 m, N 37°13'925" – E 31°22'981", 08.v.2010, Çinbilgel 6000, (H).
	<i>Echinops ritro</i> L.	İbradı, around of Gevenli Location, stony places, 1092 m, N 37°12'523" – E 31°24'506", 02.vii.2010, Çinbilgel 7235, (Th).
	<i>Erigeron acer</i> L. subsp. <i>acer</i>	Manavgat, Kızılıpınar Plateau surroundings (Nurboğazı Location), dwarf bush formation, 1980 m, N 37°14'102" – E 31°20'550", 04.viii.2009, Çinbilgel 5651; Manavgat, Derme Plateau, dwarf bush formation, 1742 m, N 37°12'973" – E 31°20'498", 05.viii.2009, Çinbilgel 5723; Manavgat, Kızılıpınar Plateau, stony places, 1854 m, N 37°13'827" – E 31°20'182", 10.x.2009, Çinbilgel 5775, (H).
	* <i>Erigeron cilicicus</i> Boiss. ex Vierh.	Manavgat, Kızılıpınar Plateau, stony places, 1750 m, N 37°14'071" – E 31°19'981", 19.vii.2008, Çinbilgel 4461, End., E. Medit., (mt.), (H).
	<i>Filago arvensis</i> L.	Manavgat, Çağlan Breach, rocky stony slopes, 1678 m, N 37°16'114" – E 31°21'305", 12.vi.2008, Çinbilgel 3953; İbradı, Melik Plateau Location, stony slopes, 1644 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4164; İbradı, Çeşgar Plateau, dwarf bush formation, 1633 m, N 37°17'430" – E 31°23'654", 01.vii.2010, Çinbilgel 6875, (Th).
	<i>Filago eriocephala</i> Guss.	Manavgat, Çağlan Breach, stony places, 1681 m, N 37°16'230" – E 31°21'349", 12.vi.2008, Çinbilgel 3973; Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), arid meadows, 1766 m, N 37°13'788" – E 31°19'208", 09.vi.2009, Çinbilgel 5209; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7179, E. Medit., (Th).
	<i>Filago pyramidata</i> L.	Manavgat, Düğnük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4070, (Th).
	* <i>Helichrysum arenarium</i> (L.) Moench subsp. <i>aucherii</i>	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings, 1568 m, N 37°13'549" – E 31°23'433", 03.vii.2011, Çinbilgel 8000, End., Ir. – Tur., (H).

	(Boiss.) P.H.Davis & Kupicha	
	<i>*Helichrysum pamphylicum</i> P.H.Davis & Kupicha	Manavgat, Burmahan Neighbourhood, Akpınar Plateau way, <i>Pinus brutia</i> forest – roadsides, 798 m, N 37°09'483" – E 31°17'167", 11.x.2009, Çinbilgel 5814, End., E. Medit., (H).
	<i>*Helichrysum plicatum</i> DC. subsp. <i>isauricum</i> Parolly	Manavgat, Çağlan Breach, rock stony slopes, 1678 m, N 37°16'114" – E 31°21'305", 12.vi.2008, Çinbilgel 3950, End., Medit., (H).
	<i>Helichrysum plicatum</i> DC. subsp. <i>plicatum</i>	Manavgat, Between Çandır Hill and Çiğdem Hill, stony slopes, 2024 m, N 37°14'656" – E 31°20'981", 11.vi.2008, Çinbilgel 3894; Manavgat, Kocatop Mountain foothills, dwarf bush formation, 1765 m, N 37°12'731" – E 31°18'904", 11.vi.2009, Çinbilgel 5405; İbradı, Melik Mountain (Hasan Hill), stony slopes, 2079 m, N 37°16'594" – E 31°23'035", 17.vi.2010, Çinbilgel 6421, (H).
	<i>Helichrysum plicatum</i> DC. subsp. <i>polyphyllum</i> (Ledeb.) P.H.Davis & Kupicha	İbradı, Melik Mountain (the eastern slopes of Hasan Hill), rocky slopes, 2020 m, N 37°16'897" – E 31°23'181", 18.vi.2008, Çinbilgel 4311, (H).
	<i>*Hieracium lasiochaetum</i> (Bornm. & Zahn) P.D.Sell & C.West	Manavgat, Kovacık Mountain, rock, 1966 m, N 37°13'612" – E 31°20'057", 29.vi.2009, Çinbilgel 5494; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1508 m, N 37°15'558" – E 31°24'354", 30.vi.2010, Çinbilgel 6652; İbradı, Akmuar Location, rock, 1042 m, N 37°11'701" – E 31°24'851", 02.vii.2010, Çinbilgel 7227, End., (H), (Det: Eren).
	<i>Hieracium pannosum</i> Boiss.	Manavgat, Kızılıpınar Plateau surroundings (east of Nurboğazi Location), stony places, 1962 m, N 37°14'256" – E 31°20'649", 11.vi.2008, Çinbilgel 3853; Manavgat, Kızılıpınar Plateau, dwarf bush-thorny cushion formation, 1861 m, N 37°14'688" – E 31°20'385", 03.viii.2009, Çinbilgel 5631; İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1147 m, N 37°12'229" – E 31°24'506", 02.vii.2010, Çinbilgel 7236, E. Medit., (H).
	<i>Hirtellina lobelii</i> (DC.) Dittrich	İbradı, Akmuar Location, rock, 1042 m, N 37°11'701" – E 31°24'851", 02.vii.2010, Çinbilgel 7224, E. Medit., (Ch).
	<i>Inula acaulis</i> Schott & Kotschy ex Boiss. var. <i>acaulis</i>	Manavgat, Kepez Plateau (İnçayırı Location), wet places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5848, (H), (Det: Eren).
	<i>Inula conyzae</i> (Griess.) Meikle	İbradı, Cevizlidere Location, under the forest – moist rocky places, 1168 m, N 37°13'646" – E 31°24'123", 02.vii.2010, Çinbilgel 7244, Euro-Sib., (H), (Det: Eren).
	<i>Inula graveolens</i> (L.) Desf.	Manavgat, Çay Neighbourhood, Kızılıpınar Plateau way, Arğıçca Breach (cutting area), 1028 m, N 37°12'886" – E 31°17'080", 10.x.2009, Çinbilgel 5759, Medit., (Th).
	<i>Inula heterolepis</i> Boiss.	Manavgat, Kızılıpınar Plateau way, Kavaklı Location, rock, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4084; Manavgat, Kalbur Gözenekleri Location, rock, 1816 m, N 37°13'101" – E 31°20'680", 05.viii.2009, Çinbilgel 5726; İbradı, Akmuar Location, rock, 1042 m, N 37°11'701" – E 31°24'851", 02.vii.2010, Çinbilgel 7223, E. Medit., (Ch).
	<i>Inula montbretiana</i> DC.	Manavgat, Çağlan Breach, rocky stony slopes, 1678 m, N 37°16'114" – E 31°21'305", 12.vi.2008, Çinbilgel 3948; İbradı, Aşağıtoka Plateau, stony places, 1530 m, N 37°12'883" – E 31°22'500", 16.vi.2010, Çinbilgel 6214; İbradı, Melik Plateau, dwarf bush formation, 1665 m, N 37°17'156" – E 31°24'327", 01.vii.2010, Çinbilgel 6923, Ir. – Tur., (Ch).
	<i>Inula oculus – christi</i> L.	Manavgat, Çağlan Breach, rocky stony slopes, 1678 m, N 37°16'114" – E 31°21'305", 12.vi.2008, Çinbilgel 3949; Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), thorny cushion formation, 1758 m, N 37°13'773" – E 31°19'193", 09.vi.2009, Çinbilgel 5195; İbradı, Around of Çeşgar Plateau Breach, dwarf bush formation, 1588 m, N 37°17'590" – E 31°23'768", 01.vii.2010, Çinbilgel 6824, Euro-Sib., (H).
	<i>Inula viscosa</i> (L.) Aiton	Manavgat, between Burmahan Neighbourhood and Çay Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°11'284" – E 31°16'291", 11.x.2009, Çinbilgel 5793; Manavgat, Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest – watersides, 729 m, N 37°09'926" – E 31°17'026", 11.x.2009, Çinbilgel 5801, Medit., (Ch).
	<i>Klasea grandifolia</i> (P.H. Davis) Greuter & Wagenitz	İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7031; İbradı, Cevizlidere Location, <i>Pinus nigra</i> forest, 1126 m, N 37°14'042" – E 31°24'212", 03.vii.2011, Çinbilgel 8031, Ir. – Tur., (H), (Det: Duman).
	<i>Lactuca hispida</i> DC.	İbradı, around of Yirme Hill, damp places, 1805 m, N 37°14'690" – E 31°23'119", 18.v.2008, Çinbilgel 3258; İbradı, between Gevenli Location and Çukuryurt Plateau way, on the rock under the <i>Pinus nigra</i> forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3520; Manavgat, Burunçalı Location surroundings, mixed <i>Cedrus libani</i> forest, 1567 m, N 37°08'813" – E 31°19'104", 02.vi.2009, Çinbilgel 4995, (H).
	<i>Lactuca intricata</i> Boiss.	İbradı, around of Yirme Hill, stony places, 1805 m, N 37°14'690" – E 31°23'119", 18.v.2008, Çinbilgel 3264; Manavgat, Dermebeli Location (Arıtaşı), rock, 1885 m, N 37°13'490" – E 31°20'297", 19.vii.2008, Çinbilgel 4865; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1511 m, N 37°15'476" – E 31°24'352", 30.vi.2010, Çinbilgel 6663, E. Medit., (mt.), (H).
	<i>Lactuca muralis</i> (L.) Gaertn.	Manavgat, Dermebeli Location, rock, 1944 m, N 37°13'711" – E 31°20'308", 29.vi.2009, Çinbilgel 5527, Euro-Sib., (G).

	<i>Lactuca viminea</i> (L.) J.Presl & C. Presl	Manavgat, Çandır Location (Kocayokuş), dwarf bush formation, 1916 m, N 37°14'168" – E 31°20'833", 04.viii.2009, Çinbilgel 5663; İbradı, Melik Plateau, forest clearings – stony places, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6555; İbradı, Tahtalı Location, <i>Pinus brutia</i> forest, 1066 m, N 37°12'606" – E 31°24'675", 03.vii.2011, Çinbilgel 8014, (H).
	<i>Leontodon asperrimus</i> (Willd.) Endl.	İbradı, Melik Plateau, stony slopes, 1644 m, N37° 17' 210" – E031° 24' 327", 17.vi.2008, Çinbilgel 4172; Manavgat, Burunçalı Location (Aslanboğazi), stony places, 1817 m, N37° 09' 027" – E031° 19' 488", 02.vi.2009, Çinbilgel 5038; İbradı, Melik Plateau, dwarf bush-thorny cushion formation, 1516 m, N 37°16'359" – E 31°24'493", 01.vii.2010, Çinbilgel 6776, Ir.–Tur., (H).
	<i>Leontodon crispus</i> Vill. subsp. <i>asper</i> (Waldst. & Kit.) Röhl. var. <i>asper</i>	Manavgat, Derme Plateau (Zahmedin Plateau – Kovantaşı Location), damp places, 1804 m, N 37°11'937" – E 31°20'141", 20.vii.2008, Çinbilgel 4502; Manavgat, Kızılpınar Plateau surroundings (Nurboğazi Location), dwarf bush-thorny cushion formation, 1980 m, N 37°14'102" – E 31°20'550", 04.viii.2009, Çinbilgel 5653; İbradı, Melik Mountain (Orta Hill), windy hills, 1948 m, N 37°16'096" – E 31°22'564", 17.vi.2010, Çinbilgel 6304, (H).
	<i>Leontodon hispidus</i> L. var. <i>hispidus</i>	Manavgat, Sarıtaş Location, stony hillsides, 2080 m, N 37°14'845" – E 31°20'924", 11.vi.2008, Çinbilgel 3898; Manavgat, Kepez Plateau (İncayırı Location), stony places, 1624 m, N 37°11'525" – E 31°20'883", 11.x.2009, Çinbilgel 5834; İbradı, Melik Mountain (Orta Hill), windy hills, 1999 m, N 37°16'161" – E 31°22'671", 17.vi.2010, Çinbilgel 6328, Euro-Sib., (H).
	<i>Leontodon oxylepis</i> Boiss. & Heldr.	Manavgat, Kızılpınar Plateau, mixed <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1718 m, N 37°14'103" – E 31°19'803", 09.vi.2009, Çinbilgel 5252; Manavgat, West of Kocatop Mountain (Örencik Location), dwarf bush, 1861 m, N 37°11'829" – E 31°18'607", 11.vi.2009, Çinbilgel 5416; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1502 m, N 37°15'548" – E 31°24'365", 30.vi.2010, Çinbilgel 6657, E. Medit., (mt.), (H).
	<i>Onopordum acanthium</i> L.	Manavgat, Dermebeli Location (Arntaşı), rocky places, 1885 m, N 37°13'490" – E 31°20'297", 19.vii.2008, Çinbilgel 4463; Manavgat, Derme Plateau, stony places, 1759 m, N 37°12'512" – E 31°20'675", 19.vii.2008, Çinbilgel 4481; Yalıncaadağ, dwarf bush formation, 2039 m, N 37°11'346" – E 31°18'682", 11.vi.2009, Çinbilgel 5406, (H).
	<i>Picnomon acarna</i> (L.) Cass.	Manavgat, Sürüçikan Mountain, debris, 1979 m, N 37°12'744" – E 31°19'789", 10.vi.2008, Çinbilgel 3786; Manavgat, Çağlan Breach, stony places, 1680 m, N 37°15'751" – E 31°20'812", 12.vi.2008, Çinbilgel 4004; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest clearings – stony places, 1001 m, N 37°10'801" – E 31°25'318", 02.vii.2010, Çinbilgel 7208, Medit., (Th).
	<i>Picris cyprica</i> Lack	Manavgat, Akpınar Plateau (Doğruluk Location), stony places, 1424 m, N 37°08'886" – E 31°20'013", 02.vi.2009, Çinbilgel 4975; İbradı, Suçikan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7082; İbradı, Gevenli Location, stony places, 1075 m, N 37°12'345" – E 31°24'803", 02.vii.2010, Çinbilgel 7148, E. Medit., (Th).
	<i>Picris hieracioides</i> L. subsp. <i>hieracioides</i>	Manavgat, Derme Plateau, stony places, 1812 m, N 37°12'698" – E 31°20'052", 20.vii.2008, Çinbilgel 4497, Euro-Sib., (H), (Det: Eren).
	<i>Picris pauciflora</i> Willd.	İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1462 m, N 37°15'665" – E 31°24'474", 30.vi.2010, Çinbilgel 6608; İbradı, Melik Plateau, forest clearings – stony places, 1406 m, N 37°15'466" – E 31°24'469", 30.vi.2010, Çinbilgel 6685; İbradı, Melik Plateau, forest clearings – rocky places, 1501 m, N 37°15'537" – E 31°24'731", 02.vii.2010, Çinbilgel 6972, Medit., (Th).
	<i>Pilosella x auriculoides</i> (Lång) Arv. -Touv.	Manavgat, Eşekbeli Location, <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1750 m, N 37°15'074" – E 31°20'047", 12.vi.2008, Çinbilgel 4020; Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), thorny cushion formation, 1771 m, N 37°13'824" – E 31°19'246", 09.vi.2009, Çinbilgel 5262; İbradı, Melik Plateau, dwarf bush formation, 1503 m, N 37°16'190" – E 31°24'575", 01.vii.2010, Çinbilgel 6767, (H).
	<i>Pilosella hoppeana</i> (Schult.) F.W.Schultz & Sch.Bip. subsp. <i>cilicica</i> (Nägeli & Peter) P.D.Sell & C.West	Manavgat, Derme Plateau (towards Dermebeli Location), stony hillsides, 1827 m, N 37°13'386" – E 31°20'237", 10.vi.2008, Çinbilgel 3795; Manavgat, Kızılpınar Plateau (Kocaboğaz Location), arid meadows, 1790 m, N 37°13'949" – E 31°20'094", 29.vi.2009, Çinbilgel 5469; İbradı, Aşağıtoka Plateau, dwarf bush-thorny cushion formation, 1526 m, N 37°12'919" – E 31°22'639", 16.vi.2010, Çinbilgel 6188, (H).
	<i>Pilosella piloselloides</i> (Vill.) Soják subsp. <i>magyarica</i> (Peter) S.Bräut. & Greuter	Manavgat, Southern hillsides of Kaldırım Mountain, under the <i>Cedrus libani</i> forest, 1551 m, N 37°09'461" – E 31°20'374", 01.vi.2009, Çinbilgel 4885; Manavgat, Kızılpınar Plateau, dwarf bush-thorny cushion formation, 1781 m, N 37°14'111" – E 31°20'180", 03.viii.2009, Çinbilgel 5646; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest, 1344 m, N 37°15'376" – E 31°24'546", 02.vii.2010, Çinbilgel 7021, (H).
	<i>Pilosella piloselloides</i> (Vill.) Soják subsp. <i>piloselloides</i>	İbradı, Melik Plateau surroundings (Towards Suçikan Location), <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7027, (H).
	<i>Pilosella procera</i> (Fr.) F.W.Schultz & Sch.Bip.	İbradı, around of Yirme Hill, forest clearings, 1674 m, N 37°14'298" – E 31°23'004", 18.v.2008, Çinbilgel 3202; Manavgat, Derme Plateau, stony places, 1812 m, N 37°12'698" – E 31°20'052", 20.vii.2008, Çinbilgel 4500; İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings, 1568 m, N 37°13'549" – E 31°23'433", 03.vii.2011, Çinbilgel 7996, (H).

	<i>*Psephellus mucronifer</i> (DC.) Wagenitz	İbradı, around of Yirme Hill, stony hills, 1875 m, N 37°14'859" – E 31°23'148", 18.v.2008, Çinbilgel 3281; Manavgat, Dermebeli Location, rock, 1920 m, N 37°13'702" – E 31°20'304", 29.vi.2009, Çinbilgel 5528; İbradı, Melik Mountain (Orta Hill), windy hills, 2183 m, N 37°16'210" – E 31°22'965", 17.vi.2010, Çinbilgel 6366, End., Ir. – Tur., (H), (Conf. Eren).
	<i>*Ptilostemon afer</i> (Jacq.) Greuter subsp. <i>eburneus</i> Greuter	Manavgat, Derme Plateau – around of Kalbur Gözenekleri, rock, 1813 m, N 37°13'337" – E 31°20'464", 05.viii.2009, Çinbilgel 5733; İbradı, Melik Plateau, dwarf bush formation, 1405 m, N 37°15'738" – E 31°24'639", 01.vii.2010, Çinbilgel 6743, End., (H).
	<i>Rhagadiolus stellatus</i> (L.) Gaertn.	İbradı, Boğazyurt Location, <i>Pinus brutia</i> forest, 970 m, N 37°11'161" – E 31°25'266", 21.v.2011, Çinbilgel 7701, Medit., (Th).
	<i>*Rhaponticoides gokceoglui</i> Çinbilgel, Eren & H.Duman	İbradı, Melik Plateau surroundings (towards Suçkan Location), <i>Pinus nigra</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7029; İbradı, Cevizlidere Location (around of Suçkan Location), <i>Pinus nigra</i> forest, 1178 m, N 37°13'517" – E 31°24'177", 08.vi.2011, Çinbilgel 7962; İbradı, Suçkan Location, <i>Pinus nigra</i> forest clearings, 1186 m, N 37°14'441" – E 31°24'305", 03.vii.2011, Çinbilgel 8021 & Eren, End., E. Medit., (H).
	<i>Scolymus hispanicus</i> L. subsp. <i>hispanicus</i>	Manavgat, Kızılıpınar Plateau way, Göçek Location, <i>Pinus brutia</i> forest, 600 m, 13.vi.2008, Çinbilgel 4156; İbradı, Eynif Plain, stony places, 945 m, N 37°08'943" – E 31°26'971", 26.ix.2010, Çinbilgel 7195, Medit., (H).
	<i>Scorzonera cana</i> (C.A.Mey.) Griseb. var. <i>cana</i>	Manavgat, Kızılıpınar Plateau, dwarf bush formation, 1750 m, N 37°13'650" – E 31°20'051", 29.vi.2009, Çinbilgel 5513; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7182, (H), (Det: Eren).
	<i>Scorzonera cana</i> (C.A.Mey.) Griseb. var. <i>jaquiniana</i> (W. Koch) D.F.Chamb.	Manavgat, Kızılıpınar Plateau way, Kavaklı Location, stony places, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4033; İbradı, Melik Plateau, dwarf bush-thorny cushion formation, 1503 m, N 37°16'190" – E 31°24'575", 01.vii.2010, Çinbilgel 6771; İbradı, Akmuar Location, rock, 1024 m, N 37°11'720" – E 31°24'844", 19.v.2011, Çinbilgel 7603, (H), (Det: Eren).
	<i>Scorzonera cana</i> (C.A.Mey.) Griseb. var. <i>radicosa</i> (Boiss.) D.F.Chamb.	İbradı, between Güzle Plateau – Yaylacık Mountain, damp places, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çinbilgel 2897; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest, 1304 m, N 37°15'106" – E 31°24'488", 02.vii.2010, Çinbilgel 6983, (H), (Det: Eren).
	<i>Scorzonera cinerea</i> Boiss.	İbradı, Toka Plateau (Göçebe Location – towards Erenler Hill), stony places, 1551 m, N 37°14'763" – E 31°22'308", 17.v.2008, Çinbilgel 3143; Manavgat, Kaldırım Mountain, dwarf bush-thorny cushion formation, 1837 m, N 37°09'993" – E 31°20'451", 01.vi.2009, Çinbilgel 4925; Manavgat, Kaldırım Mountain, 1818 m, N 37°09'957" – E 31°20'583", 03.vi.2009, Çinbilgel 5099, Ir. – Tur., (H).
	<i>Scorzonera incisa</i> DC.	İbradı, Çeşgar Plateau, stony slopes, 1659 m, N 37°17'402" – E 31°24'250", 17.vi.2008, Çinbilgel 4238; Manavgat, southern of Kaldırım Mountain, slopes, 1943 m, N 37°09'635" – E 31°20'292", 01.vi.2009, Çinbilgel 4943; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1511 m, N 37°15'476" – E 31°24'352", 30.vi.2010, Çinbilgel 6665, Ir. – Tur., (H), (Det: Eren).
	<i>Scorzonera mollis</i> M.Bieb. subsp. <i>szowitzii</i> (DC.) D.F.Chamb.	Manavgat, around of Kepez Plateau– Kaldırım Mountain hillsides, slopes, 1657 m, N 37°10'485" – E 31°21'360", 03.v.2009, Çinbilgel 4713; İbradı, Toka Plateau way, <i>Pinus nigra</i> forest, 1662 m, N 37°13'925" – E 31°22'981", 08.v.2010, Çinbilgel 5995; İbradı, Aşağıtoka Plateau, wet meadows, 1520 m, N 37°14'220" – E 31°22'415", 20.v.2011, Çinbilgel 7640, Ir. – Tur., (H).
	<i>Scorzonera pseudolanata</i> Grossh.	Manavgat, Sarıtaş Location, windy hills, 2074 m, N 37°15'015" – E 31°20'984", 11.vi.2008, Çinbilgel 3912; İbradı, Toka Plateau, <i>Pinus nigra</i> forest clearings, 1577 m, N 37°14'030" – E 31°22'843", 09.v.2010, Çinbilgel 6102, Ir. – Tur., (H).
	<i>Senecio vernalis</i> Waldst. & Kit.	Manavgat, Akpınar Plateau, stony places, 1411 m, N 37°09'389" – E 31°20'748", 03.v.2009, Çinbilgel 4679; İbradı, Eynif Plain, meadows, 941 m, N 37°08'989" – E 31°26'561", 19.v.2011, Çinbilgel 7576, (Th).
	<i>Senecio vulgaris</i> L.	Manavgat, around of Kepezbeli Location, damp places, 1485 m, N 37°09'744" – E 31°21'083", 03.v.2009, Çinbilgel 4689; İbradı, Melik Mountain (Çeşgar Hill), stony hills, 2264 m, N 37°16'008" – E 31°23'667", 18.vi.2010, Çinbilgel 6526; Manavgat, Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> clearings, 611 m, N 37°09'918" – E 31°16'962", 26.iii.2011, Çinbilgel 7468, (Th).
	<i>Solidago virgaurea</i> L. subsp. <i>virgaurea</i>	İbradı, Suçkan Location, streamsides, 1252 m, N 37°14'487" – E 31°24'182", 02.vii.2010, Çinbilgel 7100, Euro-Sib., (H), (Det: Eren).
	<i>Sonchus asper</i> (L.) Hill. subsp. <i>glaucescens</i> (Jord.) Ball	Manavgat, Arğıçca Breach Location (burning area), <i>Pinus brutia</i> forest clearings, 1090 m, N 37°13'042" – E 31°17'108", 13.vi.2008, Çinbilgel 4094; Manavgat, Kızılıpınar Plateau way, Putraklı Location, stony places, 734 m, N 37°12'249" – E 31°16'454", 27.iii.2011, Çinbilgel 7561; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7728, (H).
	<i>Sonchus oleraceus</i> L.	Manavgat, Between Çay and Burmahan Neighbourhoods, streamsides, 706 m, N 37°11'423" – E 31°16'306", 06.vi.2011, Çinbilgel 7758, (H).
	<i>*Tanacetum argenteum</i> (Lam.) Willd. subsp. <i>canum</i>	İbradı, Aşağıtoka Plateau, moist stony places, 1513 m, N 37°13'806" – E 31°22'419", 16.vi.2010, Çinbilgel 6260, End., Ir. – Tur., (Ch).

	(K.Koch) Grierson var. <i>pumilum</i> Grierson	
	<i>Tanacetum armenum</i> (DC.) Sch.Bip.	Manavgat, Kocatop Mountain, on the rock, 1811 m, N 37°13'994" – E 31°20'286", 09.vi.2008, Çinbilgel 3663; Manavgat, Kaldırım Mountain, windy hills, 2138 m, N 37°10'445" – E 31°20'597", 03.vi.2009, Çinbilgel 5121; İbradı, Melik Mountain (Orta Hill), windy hills, 2187 m, N 37°16'210" – E 31°22'985", 17.vi.2010, Çinbilgel 6382, (Ch).
	<i>Tanacetum aucheri</i> DC.	Manavgat, Susuzkır Hill, <i>Pinus nigra</i> forest, 1718 m, N 37°16'910" – E 31°21'829", 12.vi.2008, Çinbilgel 3980; Manavgat, Çağlan Breach, stony places, 1641 m, N 37°15'823" – E 31°20'867", 12.vi.2008, Çinbilgel 4002; Manavgat, Derme Plateau, wet meadows, 1798 m, N 37°13'070" – E 31°20'203", 30.vi.2009, Çinbilgel 5588, E. Medit., (H).
	<i>Taraxacum bessarabicum</i> (Hornem.) Hand. –Mazz.	İbradı, Güzle Plateau, Juniper Forest, 1306 m, N 37°09'817" – E 31°23'980", 20.iv.2008 Çinbilgel 2829; Manavgat, eastern hillsides of Kaldırım Mountain, dwarf bush formation, 1985 m, N 37°10'405" – E 31°20'777", 03.vi.2009, Çinbilgel 5116; Manavgat, Eşekbeli Location, <i>Pinus nigra</i> forest, 1612 m, N 37°14'867" – E 31°19'864", 07.vi.2011, Çinbilgel 7910, (H).
	<i>Taraxacum bithynicum</i> DC.	Manavgat, Kızılpınar Plateau, stony places, 1750 m, N 37°14'071" – E 31°19'981", 19.vii.2008, Çinbilgel 4459; Manavgat, Derme Plateau, stony places, 1825 m, N 37°13'356" – E 31°20'215", 10.x.2009, Çinbilgel 5784; İbradı, Melik Plateau, slopes, 1536 m, N 37°16'507" – E 31°24'493", 05.x.2010, Çinbilgel 7305, (H).
	<i>Taraxacum syriacum</i> Boiss.	Manavgat, Burunçalı Location, slopes, 1883 m, N 37°09'116" – E 31°18'988", 02.vi.2009, Çinbilgel 5054; İbradı, Melik Plateau surroundings (towards Suçkan Location), <i>Pinus nigra</i> forest, 1347 m, N 37°15'332" – E 31°24'499", 02.vii.2010, Çinbilgel 6998; İbradı, Aşağıtoka Plateau, wet meadows, 1520 m, N 37°14'220" – E 31°22'415", 20.v.2011, Çinbilgel 7641, Ir. – Tur., (H).
	<i>Tolpis virgata</i> (Desf.) Bertol.	İbradı, Suçkan Location, streamsides, 1259 m, N 37°14'503" – E 31°24'173", 02.vii.2010, Çinbilgel 7070, Medit., (H).
	<i>Tragopogon latifolius</i> Boiss. subsp. <i>angustifolius</i> Boiss.	İbradı, Toka Plateau way, Belova Location, stony places, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3465; Manavgat, Hillsides of Kovacık Mountain, dwarf bush formation, 1924 m, N 37°13'235" – E 31°19'993", 30.vi.2009, Çinbilgel 5570; İbradı, Yukarıtoka Plateau, dwarf bush formation, 1635 m, N 37°15'828" – E 31°22'221", 17.vi.2010, Çinbilgel 6275, Ir. – Tur., (H).
	<i>Tragopogon porrifolius</i> L. subsp. <i>abbreviatus</i> (Boiss.) Coşkunçelebi & M.Gultepe	İbradı, Tahtalı Location, stony places, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4398; İbradı, Akmuar Location, stony places, 1024 m, N 37°11'720" – E 31°24'844", 19.v.2011, Çinbilgel 7605; İbradı, Gevenli Location, <i>Pinus brutia</i> forest clearings, 1099 m, N 37°12'084" – E 31°24'524", 03.vii.2011, Çinbilgel 8006, (H).
	<i>Tragopogon porrifolius</i> L. subsp. <i>longirostris</i> (Sch.Bip.) Greuter	Manavgat, Çağlan Breach, stony places, 1681 m, N 37°16'230" – E 31°21'349", 12.vi.2008, Çinbilgel 3970; İbradı, Melik Plateau, forest clearings – rocky places, 1501 m, N 37°15'537" – E 31°24'731", 02.vii.2010, Çinbilgel 6977; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7734, (H).
	<i>Tussilago farfara</i> L.	İbradı, Suçkan Location (Taşkesiği Location), <i>Pinus nigra</i> forest – roadsides, 1308 m, N 37°15'191" – E 31°24'495", 18.vi.2008, Çinbilgel 4364; İbradı, Melik Plateau, forest clearings, 1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6968; İbradı, around of Suçkan Location, moist stony places, 1160 m, N 37°14'079" – E 31°24'218", 05.iii.2011, Çinbilgel 7362, Euro-Sib., (H).
	<i>Urospermum picroides</i> (L.) Scop. ex F.W.Schmidt	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 622 m, N 37°11'707" – E 31°16'064", 06.vi.2011, Çinbilgel 7747; Manavgat, Çay Neighbourhood, roadsides, 690 m, N 37°12'122" – E 31°16'461", 07.vi.2011, Çinbilgel 7940, Medit., (Th).
Berberidaceae		
	<i>Berberis crataegina</i> DC.	İbradı, hillsides of Melik Mountain, rocky slopes, 1880 m, N 37°15'258" – E 31°23'363", 18.v.2008, Çinbilgel 3297; İbradı, hillsides of Melik Mountain, stony-rocky slopes, 1955 m, N 37°15'593" – E 31°23'351", 18.v.2008, Çinbilgel 3302; İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest, 1695 m, N 37°13'638" – E 31°23'243", 19.v.2008, Çinbilgel 3428, (Ph).
Betulaceae		
	<i>Ostrya carpinifolia</i> Scop.	İbradı, Taşkesiği Location, <i>Pinus nigra</i> forest – roadsides, 1308 m, N 37°15'191" – E 31°24'495", 18.vi.2008, Çinbilgel 4363; İbradı, Cevizlidere Location, rocky places, 1132 m, N 37°13'382" – E 31°24'226", 02.vii.2010, Çinbilgel 7243; Manavgat, Çay Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°12'133" – E 31°16'452", 27.iii.2011, Çinbilgel 7477, Medit., (Ph).
Boraginaceae		
	<i>Alkanna areolata</i> Boiss.	Manavgat, Kızılpınar Plateau, stony – rocky places, 1720 m, N 37°14'073" – E 31°19'695", 09.vi.2009, Çinbilgel 5274; İbradı, Melik Mountain, dwarf bush formation, 2106 m, N 37°16'066" – E 31°23'466", 18.vi.2010, Çinbilgel 6497; İbradı, Gevenli Location, <i>Pinus brutia</i> forest – stony places, 1133 m, N 37°11'988" – E 31°24'420", 19.v.2011, Çinbilgel 7623, E. Medit., (H).
	* <i>Alkanna pamphylica</i> Hub. – Mor. & Reese	Manavgat, Kızılpınar Plateau surroundings (Nurboğazı Location), rocky places, 1907 m, N 37°14'167" – E 31°20'459", 11.vi.2008, Çinbilgel 3839;

		Manavgat, Akpınar Plateau– hillsides of Kaldırım Mountain, stony places, 1541 m, N 37°09'456" – E 31°20'380", 01.vi.2009, Çinbilgel 4820; Manavgat, Çay Neighbourhood surroundings (Bıçıklık stream Location), stony places, 1164 m, N 37°13'134" – E 31°17'608", 07.vi.2011, Çinbilgel 7927, End., E. Medit., (H).
	<i>Alkanna tinctoria</i> (L.) Tausch subsp. <i>anatolica</i> Hub. – Mor.	İbradı, Melik Mountain (Southwest slope of Orta Hill, stony places, 1842 m, N 37°15'907" – E 31°22'558", 17.vi.2010, Çinbilgel 6298; İbradı, Çeşgar Plateau, rock, 1590 m, N 37°17'623" – E 31°23'753", 01.vii.2010, Çinbilgel 6829; İbradı, around of Toka Plateau Breach (Kartalyuvası Location), rock, 1719 m, N 37°13'584" – E 31°23'231", 21.v.2011, Çinbilgel 7675, E. Medit., (H).
	<i>Anchusa azurea</i> Mill. var. <i>azurea</i>	Manavgat, Burmahan Neighbourhood surroundings, stony places, 704 m, N 37°11'401" – E 31°16'295", 06.vi.2011, Çinbilgel 7765, (H).
	<i>Anchusa hybrida</i> Ten.	Manavgat, around of Kepez Plateau–hillsides of Kaldırım Mountain, slopes, 1657 m, N 37°10'485" – E 31°21'360", 03.v.2009, Çinbilgel 4721; İbradı, Gevenli Location, under the forest, 1145 m, N 37°11'875" – E 31°24'457", 28.iii.2010, Çinbilgel 5936; İbradı, Suçikan Location, streamsides, 1252 m, N 37°14'487" – E 31°24'182", 02.vii.2010, Çinbilgel 7103, Çinbilgel, (H).
	<i>Anchusa leptophylla</i> Roem. & Schult. subsp. <i>leptophylla</i>	İbradı, Çeşgar Plateau, rock, 1618 m, N 37°17'456" – E 31°23'710", 01.vii.2010, Çinbilgel 6847, (H).
	<i>Asperugo procumbens</i> L.	Manavgat, Çağlan Breach, stony places, 1641 m, N 37°15'823" – E 31°20'867", 12.vi.2008, Çinbilgel 3995, Euro-Sib., (Th).
	<i>Buglossoides arvensis</i> (L.) I.M.Johnst. subsp. <i>sibthorpiana</i> (Griseb.) R.Fern.	İbradı, Gavur Breach – Etülgenboğazi, <i>Pinus brutia</i> forest, 977 m, N 37°09'100" – E 31°25'485", 20.iv.2008, Çinbilgel 3001; Manavgat, Akpınar Plateau, mixed <i>Cedrus libani</i> forest, 1485 m, N 37°09'577" – E 31°21'003", 03.v.2009, Çinbilgel 4676; İbradı, Akmuar Location, stony places, 1044 m, N 37°11'687" – E 31°24'880", 28.iii.2010, Çinbilgel 5950, (Th).
	<i>Buglossoides incrassata</i> (Guss.) I.M.Johnst. subsp. <i>incrassata</i>	İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 2083 m, N 37°16'553" – E 31°22'889", 17.vi.2010, Çinbilgel 6444, Medit., (Th).
	<i>Cerinth minor</i> L. subsp. <i>minor</i>	İbradı, Çeşgar Plateau Location, on the rock, 1642 m, N 37°17'662" – E 31°23'989", 17.vi.2008, Çinbilgel 4189; Manavgat, Burunçalı Location (Aslanboğazi), 1817 m, N 37°09'027" – E 31°19'488", 02.vi.2009, Çinbilgel 5040; İbradı, Çeşgar Plateau Breach, stony places, 1682 m, N 37°17'288" – E 31°24'324", 01.vii.2010, Çinbilgel 6915, Euro-Sib., (H).
	<i>Cynoglossum creticum</i> Mill.	Manavgat, Çay Neighbourhood surroundings (Değirmenderesi Bridge), streamsides, 705 m, N 37°12'138" – E 31°16'452", 07.vi.2011, Çinbilgel 7879, (H).
	<i>Cynoglossum montanum</i> L.	İbradı, Melik Plateau, stony slopes, 1644 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4160; İbradı, Toka Plateau Breach, stony slopes, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5974; İbradı, Melik Plateau surroundings (towards Suçikan Location), <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7036, Euro-Sib., (H).
	* <i>Cynoglottis chetikiana</i> Vural & Kit Tan subsp. <i>chetikiana</i>	Manavgat, Kepezbeli Location, stony slopes, 1616 m, N 37°10'421" – E 31°21'504", 19.v.2009, Çinbilgel 4802; Manavgat, Kaldırım Mountain, <i>Cedrus libani</i> forest clearings, 1753 m, N 37°09'809" – E 31°20'339", 01.vi.2009, Çinbilgel 4854; İbradı, Melik Plateau, <i>Pinus nigra</i> forest – stony places, 1486 m, N 37°15'604" – E 31°24'398", 30.vi.2010, Çinbilgel 6591, End., E. Medit., (H).
	* <i>Cynoglottis chetikiana</i> Vural & Kit Tan subsp. <i>paphlagonica</i> (Hausskn. ex Bornm.) Vural & Kit Tan	Manavgat, Kovacık Mountain surroundings, Kocayokuş Location, rocky places, 1887 m, N 37°14'043" – E 31°20'377", 18.v.2008, Çinbilgel 3688; İbradı, Yaylacık Mountain, <i>Cedrus libani</i> forest clearings, 1302 m, N 37°09'869" – E 31°25'078", 19.iv.2009, Çinbilgel 4611; İbradı, Gevenli Location, stony places, 1133 m, N 37°11'988" – E 31°24'420", 19.v.2011, Çinbilgel 7624, End., (H).
	<i>Echium italicum</i> L.	İbradı, Suçikan Location surroundings (Taşkesiği Location), roadsides, 1308 m, N 37°15'191" – E 31°24'495", 18.vi.2008, Çinbilgel 4370; Manavgat, Çay Neighbourhood surroundings, roadsides, 665 m, N 37°11'982" – E 31°16'333", 07.vi.2011, Çinbilgel 7876, Medit., (H).
	<i>Heliotropium hirsutissimum</i> Grauer	İbradı, Eynif Plain, arid meadows, 945 m, N 37°08'943" – E 31°26'971", 26.ix.2010, Çinbilgel 7192; İbradı, Eynif Plain, arid meadows, 943 m, N 37°08'721" – E 31°26'542", 05.x.2010, Çinbilgel 7269; İbradı, Eynif Plain, stony places, 953 m, N 37°09'488" – E 31°26'114", 05.x.2010, Çinbilgel 7270, E. Medit., (Th).
	<i>Lithodora hispidula</i> (Sm.) Griseb.	Manavgat, Çay Neighbourhood– Burmahan Neighbourhood way, under the <i>Pinus brutia</i> forest, 642 m, N 37°11'726" – E 31°16'096", 26.iii.2011, Çinbilgel 7397; Manavgat, Çay Neighbourhood, Kızılpinar Plateau way, Kocaarmut Location, <i>Pinus brutia</i> forest, forest clearings, 747 m, N 37°12'260" – E 31°16'287", 27.iii.2011, Çinbilgel 7542, E. Medit., (Ch).
	<i>Lycopsis orientalis</i> L.	İbradı, Between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çinbilgel 2929; Manavgat, Çay Neighbourhood surroundings (Değirmenderesi Bridge), streamsides, 705 m, N 37°12'138" – E 31°16'452", 07.vi.2011, Çinbilgel 7882, (Th).
	<i>Myosotis alpestris</i> F.W.Schmidt subsp. <i>alpestris</i>	İbradı, Toka Plateau (Göbebe Location), stony places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çinbilgel 3102; İbradı, Toka Plateau way, Belova

		Location, <i>Cedrus libani</i> forest, 1723 m, N 37°13'458" – E 31°23'314", 19.v.2008, Çinbilgel 3456; Manavgat, Akpınar Plateau (Yumrutaş Location), <i>Cedrus libani</i> forest clearings, 1514 m, N 37°09'991" – E 31°21'166", 19.v.2009, Çinbilgel 4730, (H).
	<i>Myosotis lithospermifolia</i> Hornem.	İbradı, Yirme Hill, forest clearings, 1674 m, N 37°14'298" – E 31°23'004", 18.v.2008, Çinbilgel 3208; İbradı, Melik Mountain (Orta Hill), dwarf bush formation, 2153 m, N 37°16'186" – E 31°22'903", 17.vi.2010, Çinbilgel 6337; İbradı, Toka Plateau Breach, meadows, 1647 m, N 37°13'777" – E 31°23'191", 21.v.2011, Çinbilgel 7660, (H).
	<i>Myosotis minutiflora</i> Boiss. & Reut.	Manavgat, Kızılıpınar Plateau surroundings (Nurboğazi Location), rocky places, 1907 m, N 37°14'167" – E 31°20'459", 11.vi.2008, Çinbilgel 3830; Manavgat, Kızılıpınar Plateau (Nurboğazi Location surroundings), stony places, 1962 m, N 37°14'256" – E 31°20'649", 11.vi.2008, Çinbilgel 3859, Medit., (Th).
	<i>Myosotis ramosissima</i> Rochel.	İbradı, between Gavur Breach and Etülgenboğazi, <i>Pinus brutia</i> forest, 977 m, N 37°09'100" – E 31°25'485", 20.iv.2008, Çinbilgel 3000; Manavgat, Kaldırım Mountain, <i>Cedrus libani</i> forest clearings, 1753 m, N 37°09'809" – E 31°20'339", 01.vi.2009, Çinbilgel 4857, (Th).
	<i>Myosotis refracta</i> Boiss. subsp. <i>refracta</i>	Manavgat, Kızılıpınar Plateau, under the <i>Cedrus libani</i> forest, 1717 m, N 37°14'110" – E 31°19'899", 09.vi.2009, Çinbilgel 5238; İbradı, Çukuryurt Plateau surroundings, stony places, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6170, Medit., (Th).
	<i>Myosotis stricta</i> Roem. & Schult.	İbradı, Yukarıtoka Plateau (Between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3083; Manavgat, Akpınar Plateau (Kelsu Location), stony places, 1591 m, N 37°09'534" – E 31°20'305", 01.vi.2009, Çinbilgel 4840; Manavgat, eastern hillsides of Kaldırım Mountain, dwarf bush-thorny cushion formation, 1985 m, N 37°10'405" – E 31°20'777", 03.vi.2009, Çinbilgel 5117, Euro-Sib., (Th).
	* <i>Omphalodes luciliae</i> Boiss. subsp. <i>luciliae</i>	İbradı, Melik Mountain (Orta Hill), windy hills, 2126 m, N 37°16'348" – E 31°23'067", 17.vi.2010, Çinbilgel 6394, End., E. Medit., (mt.), (H).
	* <i>Omphalodes ripleyana</i> P.H.Davis	İbradı, around of Yirme Hill, rock crevices, 1814 m, N 37°14'700" – E 31°23'139", 18.v.2008, Çinbilgel 3225; İbradı, Melik Mountain (Hasan Hill), stony hillsides, 2049 m, N 37°16'896" – E 31°23'156", 18.vi.2008, Çinbilgel 4328; Manavgat, Burunçalı Location (Aslanboğazi), rock, 1907 m, N 37°08'951" – E 31°19'501", 02.vi.2009, Çinbilgel 5050, End., E. Medit. (mt.), (H).
	* <i>Onosma armena</i> DC.	İbradı, Toka Plateau Breach surroundings, <i>Cedrus libani</i> forest clearings, 1570 m, N 37°13'602" – E 31°23'417", 09.v.2010, Çinbilgel 6123, End., (H).
	<i>Onosma bracteosa</i> Hausskn. & Bornm.	Manavgat, Kocatop Mountain, windy hills, 2106 m, N 37°12'578" – E 31°18'867", 10.vi.2009, Çinbilgel 5362; İbradı, Melik Mountain (Orta Hill), rocky places, 1967 m, N 37°16'123" – E 31°22'618", 17.vi.2010, Çinbilgel 6318; İbradı, Suçkan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7085, End., Ir. – Tur., (H).
	* <i>Onosma inexpectata</i> Teppner	İbradı, Çukuryurt Plateau, forest, 1260 m, N 37°12'026" – E 31°24'001", 19.v.2011, Çinbilgel 7625, End., E. Medit., (H).
	* <i>Onosma isaurica</i> Boiss. & Heldr.	İbradı, Toka Plateau way, under the <i>Cedrus libani</i> forest – stony places, 1460 m, N 37°13'078" – E 31°23'776", 19.v.2008, Çinbilgel 3493; İbradı, Melik Mountain (Hasan Hill), stony places, 2049 m, N 37°16'896" – E 31°23'156", 18.vi.2008, Çinbilgel 4322; İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1048 m, N 37°12'283" – E 31°24'732", 19.v.2011, Çinbilgel 7614, End., Ir. – Tur., (H).
	<i>Onosma oreodoxa</i> Boiss.	Manavgat, Burmahan Neighbourhood – Kovacık Mountain way, <i>Pinus brutia</i> forest, 886 m, N 37°13'095" – E 31°16'406", 25.v.2008, Çinbilgel 3602/a; Manavgat, Arıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4104; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7731, E. Medit., (H).
	<i>Paracaryum lithospermifolium</i> (Lam.) Grande subsp. <i>cariense</i> (Boiss.) R.R.Mill var. <i>cariense</i>	Manavgat, around of Çağlan Breach, moving rock and stony places, 1665 m, N 37°15'851" – E 31°21'000", 12.vi.2008, Çinbilgel 3928; Manavgat, Kaldırım Mountain (Aktünek Hill), windy hills, 2198 m, N 37°10'556" – E 31°20'329", 03.vi.2009, Çinbilgel 5133; İbradı, Melik Mountain, stony places, 2266 m, N 37°16'071" – E 31°23'703", 18.vi.2010, Çinbilgel 6521, Medit., (mt.), (H).
Brassicaceae		
	<i>Aethionema arabicum</i> (L.) Andr. ex DC.	İbradı, Yukarıtoka Plateau (Between Beydiğın Plateau and Göçebe Location), stony places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3069; İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3477, (Th).
	<i>Aethionema cordatum</i> (Desf.) Boiss.	İbradı, Gevenli Location, <i>Pinus nigra</i> forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3515; Manavgat, Kızılıpınar Plateau, rocky places, 1735 m, N 37°14'127" – E 31°19'929", 25.v.2008, Çinbilgel 3515/a, Ir. – Tur., (H).
	<i>Aethionema oppositifolium</i> (Pers.) Hedge	Manavgat, Sarıtaş Location, stony places, 2084 m, N 37°15'212" – E 31°20'984", 11.vi.2008, Çinbilgel 3906; Manavgat, Kaldırım Mountain (Aktünek Hill), windy hills, 2190 m, N 37°10'583" – E 31°20'323", 03.vi.2009,

		Çinbilgel 5156; İbradı, Melik Mountain (Orta Hill), windy hills, 2183 m, N 37°16'210" – E 31°22'965", 17.vi.2010, Çinbilgel 6364, (H).
	<i>*Aethionema subulatum</i> (Boiss. & Heldr.) Boiss.	Manavgat, hillsides of Çandır Hill, stony hills, 2037 m, N 37°16'626" – E 31°21'029", 11.vi.2008, Çinbilgel 3879; Manavgat, Kaldırım Mountain (Aktünek Hill), windy hills, 2198 m, N 37°10'556" – E 31°20'329", 03.vi.2009, Çinbilgel 5129; İbradı, Melik Mountain (Orta Hill), windy hills, 2053 m, N 37°16'182" – E 31°22'761", 17.vi.2010, Çinbilgel 6323, End., (Ch).
	<i>*Alyssum argyrophyllum</i> Schott & Kotschy	İbradı, hillsides of Melik Mountain (Çataldüzlügen Location), slopes, 1843 m, N 37°15'258" – E 31°23'363", 18.v.2008, Çinbilgel 3289; Manavgat, Kaldırım Mountain, 1818 m, N 37°09'957" – E 31°20'583", 03.vi.2009, Çinbilgel 5100; İbradı, Melik Mountain (southern hillsides of Hasan Hill), dwarf bush, 2073 m, N 37°16'539" – E 31°23'096", 17.vi.2010, Çinbilgel 6436, End., (Ch).
	<i>Alyssum desertorum</i> Stapf.	İbradı, Yirme Location, rocky hills, stony places, 1820 m, N 37°14'717" – E 31°23'178", 18.v.2008, Çinbilgel 3247; İbradı, Aşağıtoka Plateau, stony places, 1530 m, N 37°13'109" – E 31°22'633", 20.v.2011, Çinbilgel 7646, (Th).
	<i>Alyssum hirsutum</i> M.Bieb. subsp. <i>hirsutum</i>	İbradı, Yukarıtoka Plateau (Between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3054, (Th).
	<i>*Alyssum lepidotum</i> Boiss.	İbradı, Aşağıtoka Plateau, dwarf bush-thorny cushion formation, 1523 m, N 37°12'927" – E 31°22'506", 16.vi.2010, Çinbilgel 6210; İbradı, Melik Plateau, dwarf bush-thorny cushion formation, 1503 m, N 37°16'190" – E 31°24'575", 01.vii.2010, Çinbilgel 6766; İbradı, Aşağıtoka Plateau, dwarf bush, 1529 m, N 37°12'869" – E 31°22'570", 02.vii.2011, Çinbilgel 7981, End., (Ch).
	<i>Alyssum minutum</i> Schlecht. ex DC.	Manavgat, around of Kepezbeli Location, dolinense, 1485 m, N 37°09'744" – E 31°21'083", 03.v.2009, Çinbilgel 4690, (Th).
	<i>Alyssum mouradicum</i> Boiss. & Balansa	İbradı, hillsides of Melik Mountain, rocky slopes, 1880 m, N 37°15'258" – E 31°23'363", 18.v.2008, Çinbilgel 3291; Manavgat, Around of Kepez Plateau (hillsides of Kaldırım Mountain), slopes, 1657 m, N37° 10' 485" – E031° 21' 360", 03.v.2009, Çinbilgel 4718; İbradı, around of Toka Plateau Breach, 1570 m, N 37°13'602" – E 31°23'417", 09.v.2010, Çinbilgel 6125, (H).
	<i>Alyssum murale</i> Waldst. & Kit. subsp. <i>murale</i> var. <i>murale</i>	İbradı, Çeşgar Plateau, on the rock, 1642 m, N 37°17'662" – E 31°23'989", 17.vi.2008, Çinbilgel 4187, (Ch).
	<i>Alyssum simplex</i> Rudolph	İbradı, Güzle Plateau surroundings, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2815; Manavgat, around of Kepez Plateau (hillsides of Kaldırım Mountain), slopes, 1657 m, N 37°10'485" – E 31°21'360", 03.v.2009, Çinbilgel 4716; İbradı, around of Toka Plateau Breach, <i>Cedrus libani</i> – <i>Pinus nigra</i> forest, 1679 m, N 37°13'808" – E 31°23'202", 21.v.2011, Çinbilgel 7669, (Th).
	<i>Alyssum strictum</i> Willd.	İbradı, Yukarıtoka Plateau (Between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3089; Manavgat, Kızılpınar Plateau, under the <i>Cedrus libani</i> forest, 1717 m, N 37°14'110" – E 31°19'899", 09.vi.2009, Çinbilgel 5231; İbradı, Melik Mountain (Hasan Hill), slopes, 2079 m, N 37°16'594" – E 31°23'035", 17.vi.2010, Çinbilgel 6420, Ir. – Tur., (Th).
	<i>Alyssum strigosum</i> Banks & Sol. subsp. <i>strigosum</i>	İbradı, hillsides of Yaylacık Mountain, <i>Pinus brutia</i> forest, 1155 m, N 37°09'781" – E 31°25'476", 19.iv.2009, Çinbilgel 4582; İbradı, Akmuar Location, stony slopes, 1044 m, N 37°11'687" – E 31°24'880", 28.iii.2010, Çinbilgel 5951; İbradı, Melik Plateau surroundings, rocky places, forest clearings, 1501 m, N 37°15'537" – E 31°24'731", 02.vii.2010, Çinbilgel 6976, (Th).
	<i>Anchonium elichrysofolium</i> (DC.) Boiss. subsp. <i>elichrysofolium</i> var. <i>elichrysofolium</i>	Manavgat, Kocatop Mountain, snow patch-stony places, 2080 m, N 37°12'599" – E 31°18'884", 10.vi.2009, Çinbilgel 5281; Manavgat, Kocatop Mountain, windy hills, 2080 m, N 37°12'398" – E 31°18'732", 10.vi.2009, Çinbilgel 5352; İbradı, Toka Plateau Breach, stony places, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5969, (H), (Det: Eren).
	<i>Arabis alpina</i> L. subsp. <i>brevifolia</i> (DC.) Cullen	Manavgat, between Dermebeli Location and Sürüçkan Mountain, stony slopes, 1914 m, N 37°13'471" – E 31°20'139", 10.vi.2008, Çinbilgel 3735; İbradı, Toka Plateau Breach, stony slopes, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5987; İbradı, Melik Mountain (Orta Hill), windy hills, 2126 m, N 37°16'348" – E 31°23'067", 17.vi.2010, Çinbilgel 6390, E. Medit., (mt.), (H).
	<i>*Arabis deflexa</i> Boiss.	İbradı, around of Ekinlik Location, <i>Pinus nigra</i> forest (mixed with <i>Cedrus libani</i>), 1458 m, N 37°13'114" – E 31°23'740", 09.v.2010, Çinbilgel 6154; İbradı, Suçikan Location, stony places, 1130 m, N 37°14'407" – E 31°24'267", 05.iii.2011, Çinbilgel 7365; Manavgat, Çay Neighbourhood, Kavaklı Location, <i>Juniperus excelsa</i> forest, 1323 m, N 37°13'974" – E 31°18'405", 07.vi.2011, Çinbilgel 7925, End., E. Medit., (H).
	<i>Arabis ionocalyx</i> Boiss.	İbradı, southern slopes of Yaylacık Mountain, stony places, 1541 m, N 37°10'262" – E 31°24'049", 18.iv.2009, Çinbilgel 4577; İbradı, southeast hillsides of Yaylacık Mountain, <i>Cedrus libani</i> forest clearings, 1263 m, N 37°09'958" – E 31°25'227", 19.iv.2009, Çinbilgel 4610; İbradı, Suçikan Location, streamsides, 1259 m, N 37°14'503" – E 31°24'173", 02.vii.2010, Çinbilgel 7067, E. Medit., (H).

	<i>Arabis nova</i> Vill.	İbradı, Suçkan Location way, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2768; Manavgat, Burunçalı Location, slopes, 1883 m, N 37°09'116" – E 31°18'988", 02.vi.2009, Çinbilgel 5058; Manavgat, Çay Neighbourhood, Eşekbeli Location, <i>Pinus nigra</i> forest, 1612 m, N 37°14'867" – E 31°19'864", 07.vi.2011, Çinbilgel 7914, (Th).
	<i>Arabis verna</i> (L.) R.Br.	İbradı, Gavur Breach way – Etülgenboğazi, <i>Pinus brutia</i> forest, 977 m, N 37°09'100" – E 31°25'485", 20.iv.2008, Çinbilgel 2995; İbradı, southeast hillsides of Yaylacık Mountain, <i>Pinus brutia</i> forest, 1155 m, N 37°09'781" – E 31°25'476", 19.iv.2009, Çinbilgel 4579; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 990 m, N 37°10'778" – E 31°25'386", 21.v.2011, Çinbilgel 7691, Medit., (Th).
	* <i>Aubrieta canescens</i> (Boiss.) Bornm. subsp. <i>canescens</i>	Manavgat, Dermebeli–Sürüçkan Mountain, rocky slopes, 1917 m, N 37°12'891" – E 31°19'890", 10.vi.2008, Çinbilgel 3771; İbradı, Melik Mountain (Hasan Hill), rocky slopes, 2000 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4346; Manavgat, Dermebeli Location (Artaşı), rocky, 1823 m, N 37°13'506" – E 31°20'339", 30.vi.2009, Çinbilgel 5545, End., (H).
	<i>Barbarea</i> Boiss. subsp. <i>minor</i> (K. Koch) Parolly & Eren var. <i>minor</i>	İbradı, Suçkan Location way, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2775; Manavgat, Akpınar Plateau, mixed <i>Cedrus libani</i> forest, 1485 m, N 37°09'577" – E 31°21'003", 03.v.2009, Çinbilgel 4673; İbradı, Boğazyurt Location, stony places, 1048 m, N 37°11'404" – E 31°25'002", 28.iii.2010, Çinbilgel 5961, Ir. – Tur., (H).
	* <i>Bornmuellara cappadocica</i> (Willd.) Cullen & T.R.Dudley	İbradı, Toka Plateau, moist stony places, 1500 m, N 37°14'332" – E 31°22'569", 08.v.2010, Çinbilgel 6050 & Eren, End., Ir. – Tur., (H).
	<i>Calepina irregularis</i> (Asso) Thell.	İbradı, Güzle Plateau – Mezarlı Location, 1287 m, N 37°10'193" – E 31°23'650", 18.iv.2009, Çinbilgel 4566; İbradı, Akmuar Location, rocky places, 1024 m, N 37°11'720" – E 31°24'844", 19.v.2011, Çinbilgel 7602, (Th).
	<i>Capsella bursa – pastoris</i> (L.) Medik.	İbradı, Toka Plateau, stony places, 1528 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3325; Manavgat, Akpınar Plateau, mixed <i>Cedrus libani</i> forest, 1485 m, N 37°09'577" – E 31°21'003", 03.v.2009, Çinbilgel 4668; İbradı, Toka Plateau, wet meadows, 1528 m, N 37°14'074" – E 31°22'444", 09.v.2010, Çinbilgel 6098, (Th).
	<i>Capsella rubella</i> Reut.	Manavgat, Akpınar Plateau way, <i>Pinus brutia</i> forest, 900 m, N 37°09'068" – E 31°17'826", 26.iii.2009, Çinbilgel 4551; İbradı, hillsides of Yaylacık Mountain, under the <i>Cedrus libani</i> forest, 1356 m, N 37°09'929" – E 31°25'029", 19.iv.2009, Çinbilgel 4586; Manavgat, Kızılıpınar Plateau, under the <i>Cedrus libani</i> forest, 1717 m, N 37°14'110" – E 31°19'899", 09.vi.2009, Çinbilgel 5237, Medit., (Th).
	<i>Cardamine graeca</i> L.	İbradı, Gevenli Location, <i>Pinus nigra</i> forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3525; İbradı, Çukuryurt Plateau, meadows, 1268 m, N 37°12'117" – E 31°24'023", 28.iii.2010, Çinbilgel 5932; İbradı, Boğazyurt Location, stony places, 1048 m, N 37°11'404" – E 31°25'002", 28.iii.2010, Çinbilgel 5957, (Th).
	<i>Cardamine hirsuta</i> L.	Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4651; İbradı, around of Yirme Hill, s forest, 1550 m, N 37°14'328" – E 31°22'786", 08.v.2010, Çinbilgel 6024; Burmağan Neighbourhood, around of Erenli Location, streamsides, 753 m, N 37°09'844" – E 31°16'991", 26.iii.2011, Çinbilgel 7438, (Th).
	* <i>Clypeola ciliata</i> Boiss.	İbradı, around of Yirme Hill (Dikenli Karlığı), rocky stony places, 1814 m, N 37°14'700" – E 31°23'139", 18.v.2008, Çinbilgel 3216; Manavgat, Akpınar Plateau (Hincirak Location), rubble rocks, 1468 m, N 37°09'337" – E 31°20'412", 01.vi.2009, Çinbilgel 4871; İbradı, Çeşgar Plateau, rock, 1623 m, N 37°17'463" – E 31°23'699", 01.vii.2010, Çinbilgel 6852, End., E. Medit., (Th).
	<i>Clypeola jonthlaspi</i> L.	Manavgat, Burunçalı Location, rock, 1815 m, N 37°09'107" – E 31°18'850", 02.vi.2009, Çinbilgel 5083, (Th).
	<i>Draba bruniifolia</i> Steven subsp. <i>bruniifolia</i>	Manavgat, Kaldırım Mountain (Aktünek Hill), windy hills, 2208 m, N 37°10'562" – E 31°20'327", 03.vi.2009, Çinbilgel 5108; Manavgat, Kaldırım Mountain, windy hills, 2192 m, N 37°10'494" – E 31°20'513", 03.vi.2009, Çinbilgel 5123; Manavgat, Kocatop Mountain, windy hills, 2174 m, N 37°12'467" – E 31°18'805", 10.vi.2009, Çinbilgel 5324, (Ch).
	<i>Draba heterocoma</i> Fenzl	İbradı, Yirme Location, rocky hills, stony places, 1820 m, N 37°14'717" – E 31°23'178", 18.v.2008, Çinbilgel 3230; Manavgat, Sarıtaş Location, windy stony hillsides, 2074 m, N 37°15'015" – E 31°20'984", 11.vi.2008, Çinbilgel 3919; İbradı, Melik Mountain (Orta Hill), windy hills, 2175 m, N 37°16'298" – E 31°23'052", 17.vi.2010, Çinbilgel 6376, (Ch).
	<i>Draba verna</i> L.	İbradı, Toka Plateau, under the rock, 1512 m, N 37°13'755" – E 31°22'602", 28.iii.2010, Çinbilgel 5898; İbradı, Toka Plateau, wet meadows, 1528 m, N 37°14'074" – E 31°22'444", 09.v.2010, Çinbilgel 6097; Manavgat, Çay Neighbourhood, Kızılıpınar Plateau way (Düğnük Plateau), degraded <i>Pinus brutia</i> forest, 1163 m, N 37°13'385" – E 31°17'716", 27.iii.2011, Çinbilgel 7502; İbradı, Toka Plateau, moist stony places, 1522 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3342; İbradı, Eynif Plain, Kışla Location, stony places, 954 m, N 37°09'724" – E 31°26'060", 19.iv.2009, Çinbilgel 4631; İbradı, Akpınar (Akmuar) Location, moist stony places, 1062 m, N 37°11'985" – E 31°24'815", 05.iii.2011, Çinbilgel 7373 (Th).

	<i>*Erysimum kotschyannum</i> J.Gay	İbradı, Melik Mountain (Çeşgar Hill), stony slopes, 2100 m, N 37°15'610" – E 31°23'438", 18.v.2008, Çinbilgel 3314; İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings, rock, 1695 m, N 37°13'638" – E 31°23'243", 19.v.2008, Çinbilgel 3432; İbradı, Toka Plateau way, Belova Location, rock, 1695 m, N 37°13'638" – E 31°23'243", 19.v.2008, Çinbilgel 3440, End., (Ch).
	<i>Erysimum scabrum</i> DC.	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest, roadsides, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3485; İbradı, Toka Plateau Breach, stony places, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5975; İbradı, Melik Plateau, streamsides, 1537 m, N 37°15'549" – E 31°24'633", 02.vii.2010, Çinbilgel 6951, (H).
	<i>Erysimum smyrnaeum</i> Boiss. & Balansa	İbradı, around of Yirme Hill, forest clearings, 1783 m, N 37°14'667" – E 31°23'097", 18.v.2008, Çinbilgel 3180; Manavgat, Sarıtaş Location, stony hillsides, 2074 m, N 37°15'015" – E 31°20'984", 11.vi.2008, Çinbilgel 3911; Manavgat, Çay Neighbourhood, Kavaklı Location, <i>Juniperus excelsa</i> forest, 1374 m, N 37°14'135" – E 31°18'658", 07.vi.2011, Çinbilgel 7918, (H).
	<i>Fibigia clypeata</i> (L.) Medik. subsp. <i>clypeata</i> var. <i>clypeata</i>	İbradı, Melik Mountain, stony slopes, 1880 m, N 37°14'957" – E 31°23'372", 18.v.2008, Çinbilgel 3269; İbradı, Hillsides of Melik Mountain, rocky slopes, 1880 m, N 37°15'258" – E 31°23'363", 18.v.2008, Çinbilgel 3293; İbradı, Çukuryurt Plateau surroundings, rocky places, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6175, (H).
	<i>Heldreichia bupleurifolia</i> Boiss. subsp. <i>bupleurifolia</i>	İbradı, Güzle Plateau – Yaylacık Mountain, screes, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çinbilgel 2894; Manavgat, Kaldırım Mountain, 1839 m, N 37°09'906" – E 31°20'407", 01.vi.2009, Çinbilgel 4950; İbradı, Melik Mountain (Çeşgar Hill), stony hills, 2270 m, N 37°16'036" – E 31°23'689", 18.vi.2010, Çinbilgel 6522, (H).
	<i>*Hesperis bicuspidata</i> (Willd.) Poir.	Manavgat, Yalıncaadağ, rocky hills, 2068 m, N 37°11'428" – E 31°18'822", 11.vi.2009, Çinbilgel 5368, End., Ir. – Tur., (H), (Conf. Duran).
	<i>Iberis carnososa</i> Willd.	İbradı, Gevenli Location, under the <i>Pinus brutia</i> forest, 1108 m, N 37°12'316" – E 31°24'613", 17.v.2008, Çinbilgel 3019; İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1048 m, N 37°12'283" – E 31°24'732", 19.v.2011, Çinbilgel 7613/a, E. Medit., (Th).
	<i>Iberis sempervirens</i> L.	Manavgat, Kızılpınar Plateau surroundings, Nurboğazi Location, rocky places, 1907 m, N 37°14'167" – E 31°20'459", 11.vi.2008, Çinbilgel 3842; Manavgat, Kızılpınar Plateau, dwarf bush-thorny cushion formation, 1861 m, N 37°14'688" – E 31°20'385", 03.viii.2009, Çinbilgel 5630; İbradı, Çeşgar Plateau, rock, 1625 m, N 37°17'470" – E 31°23'695", 01.vii.2010, Çinbilgel 6841, E. Medit., (Ch).
	<i>Iberis simplex</i> DC.	İbradı, Gevenli Location, under the <i>Pinus brutia</i> forest, 1108 m, N 37°12'316" – E 31°24'613", 17.v.2008, Çinbilgel 3020; İbradı, Çukuryurt Plateau surroundings, stony places, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6165; İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1048 m, N 37°12'283" – E 31°24'732", 19.v.2011, Çinbilgel 7613, (H).
	<i>Isatis tinctoria</i> L. subsp. <i>corymbosa</i> (Boiss.) P.H.Davis	İbradı, Aşağıtoka Plateau, debris, 1547 m, N 37°13'903" – E 31°22'326", 19.v.2008, Çinbilgel 3361; İbradı, Melik Mountain (Hasan Hill), slopes, 2000 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4339; Manavgat, Dermebeli Location (Artaşı), on the rock, 1910 m, N37°13'547" – E 31°20'315", 30.vi.2009, Çinbilgel 5597, (H).
	<i>Lepidium chalepense</i> L.	İbradı, Toka Plateau (Beydiğin Plateau), stony places, 1544 m, N 37°14'435" – E 31°22'650", 18.v.2008, Çinbilgel 3155; Manavgat, Akpınar Plateau, stony places, 1403 m, N 37°09'414" – E 31°20'759", 19.v.2009, Çinbilgel 4750; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çinbilgel 6716, (H).
	<i>Malcolmia chia</i> (L.) DC.	Manavgat, Burunçalı Location, mixed <i>Cedrus libani</i> forest, 1567 m, N 37°08'813" – E 31°19'104", 02.vi.2009, Çinbilgel 4993; Manavgat, Burunçalı Location, 1581 m, N37°08'893" – E 31°18'959", 02.vi.2009, Çinbilgel 5086, E. Medit., (Th).
	<i>Nasturtium officinale</i> R.Br.	İbradı, Melik Plateau, streams, 1368 m, N 37°15'549" – E 31°24'633", 02.vii.2010, Çinbilgel 6953, (H).
	<i>Neslia paniculata</i> (L.) Desv. subsp. <i>thracica</i> (Velen.) Borm.	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way (Kocaarmut Location), <i>Pinus brutia</i> forest clearings – stony places, 709 m, N 37°12'174" – E 31°16'360", 27.iii.2011, Çinbilgel 7545, (Th).
	<i>Raphanus raphanistrum</i> L. subsp. <i>raphanistrum</i>	İbradı, Melik Plateau, openings, stony places, 1387 m, N 37°15'533" – E 31°24'528", 30.vi.2010, Çinbilgel 6731, (Th).
	<i>*Ricotia varians</i> B.L. Burttt	Manavgat, Burunçalı Location (Aslanboğazi), screes, 1732 m, N 37°08'976" – E 31°19'499", 02.vi.2009, Çinbilgel 5016; Manavgat, Kovacık Mountain, 1871 m, N 37°13'099" – E 31°19'006", 10.vi.2009, Çinbilgel 5307; İbradı, Melik Mountain, (Çeşgar Hill), screes, 2270 m, N 37°16'036" – E 31°23'689", 18.vi.2010, Çinbilgel 6524, End., Ir. – Tur., (H).
	<i>Sinapis arvensis</i> L.	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, Kocaarmut Location, <i>Pinus brutia</i> forest clearings, 709 m, N 37°12'174" – E 31°16'360", 27.iii.2011, Çinbilgel 7555, (Th).
	<i>Sisymbrium officinale</i> (L.) Scop.	Manavgat, Between Burmahan Neighbourhood and Kovacık Mountain, rocky places, 1214 m, N 37°13'606" – E 31°17'996", 25.v.2008, Çinbilgel 3591; İbradı, Akmuar Location, rock, 1024 m, N 37°11'720" – E 31°24'844",

		19.v.2011, Çinbilgel 7601; Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7741, (Th).
	<i>Sisymbrium orientale</i> L.	Manavgat, Arğuca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4119; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7724, (Th).
	<i>Sobolewsia clavata</i> (Boiss.) Fenzl	Manavgat, Around of Burunçalı Location, mixed <i>Cedrus libani</i> forest, 1567 m, N 37°08'813" – E 31°19'104", 02.vi.2009, Çinbilgel 4994; Manavgat, Kovacık Mountain, 1871 m, N 37°13'099" – E 31°19'006", 10.vi.2009, Çinbilgel 5308; İbradı, Toka Plateau Breach, stony places, 1691 m, N 37°13'748" – E 31°23'142", 09.v.2010, Çinbilgel 6112, (H).
	<i>Thlaspi perfoliatum</i> L.	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest, 1723 m, N 37°13'458" – E 31°23'314", 19.v.2008, Çinbilgel 3458; İbradı, Çukuryurt Plateau, meadows, 1268 m, N 37°12'117" – E 31°24'023", 28.iii.2010, Çinbilgel 5922; Çay Neighbourhood, Kızılpınar Plateau way (Düğük Plateau), degraded <i>Pinus brutia</i> forest, 1163 m, N 37°13'385" – E 31°17'716", 27.iii.2011, Çinbilgel 7511, (Th).
	<i>Thlaspi ochroleucum</i> Boiss. & Heldr.	İbradı, Dikenli Location, forest clearings, 1674 m, N 37°14'298" – E 31°23'004", 18.v.2008, Çinbilgel 3209; Manavgat, Kepez Plateau (İnçayırı Location), meadows, 1632 m, N 37°10'647" – E 31°21'180", 19.v.2009, Çinbilgel 4777; İbradı, Toka Plateau way, <i>Pinus nigra</i> forest clearings, 1554 m, N 37°14'132" – E 31°22'822", 09.v.2010, Çinbilgel 6107, (Ch).
	<i>Turritis laxa</i> (Sibth. & Sm.) Hayek	İbradı, Suçikan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7081; Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 749 m, N 37°09'726" – E 31°17'001", 06.vi.2011, Çinbilgel 7815, (Th).
Campanulaceae		
	<i>Asyneuma compactum</i> (Boiss. & Heldr.) Damboldt var. <i>compactum</i>	Manavgat, Çandır Hill, stony hills, 2037 m, N 37°16'626" – E 31°21'029", 11.vi.2008, Çinbilgel 3882; Manavgat, Kocatop Mountain, windy hills, 2076 m, N 37°12'643" – E 31°18'882", 10.vi.2009, Çinbilgel 5282; İbradı, Melik Mountain (Orta Hill), windy hills, 2187 m, N 37°16'210" – E 31°22'985", 17.vi.2010, Çinbilgel 6388; Manavgat, Kocatop Mountain (Koca Hill = Köygöründüğü Location), windy hills, 2150 m, N 37°12'346" – E 31°18'633", 10.vi.2009, Çinbilgel 5336, End., E. Medit., (mt.), (H).
	<i>Asyneuma limonifolium</i> (L.) Janehen subsp. <i>limonifolium</i>	Manavgat, Kepezbeli Location, slopes, 1674 m, N 37°10'329" – E 31°21'582", 03.v.2009, Çinbilgel 4706; Manavgat, Kovacık Mountain foothills, thorny cushion formation, 1832 m, N 37°13'397" – E 31°19'049", 10.vi.2009, Çinbilgel 5316; Manavgat, Kızılpınar Plateau (Kocaboğaz Location), arid meadows, 1790 m, N 37°13'949" – E 31°20'094", 29.vi.2009, Çinbilgel 5477, (H).
	* <i>Asyneuma limonifolium</i> (L.) Janehen subsp. <i>pestalozzae</i> (Boiss.) Damboldt	Manavgat, between Derme Plateau and Sürüçikan Mountain, dwarf bush formation 2223 m, N 37°12'479" – E 31°19'531", 30.vi.2009, Çinbilgel 5594, End., (H).
	* <i>Asyneuma linifolium</i> (Boiss. & Heldr.) Bornm. subsp. <i>linifolium</i>	İbradı, Çeşgar Plateau, rock, 1642 m, N 37°17'662" – E 31°23'989", 17.vi.2008, Çinbilgel 4196; Manavgat, Burunçalı Location, rock, 1815 m, N 37°09'107" – E 31°18'850", 02.vi.2009, Çinbilgel 5075; İbradı, around of Çeşgar Plateau Breach, stony places, 1681 m, N 37°17'294" – E 31°24'331", 01.vii.2010, Çinbilgel 6801, End., E. Medit., (mt.), (H).
	<i>Asyneuma lobelioides</i> (Willd.) Hand.–Mazz.	Manavgat, Çandır Hill, stony hills, 2037 m, N 37°16'626" – E 31°21'029", 11.vi.2008, Çinbilgel 3882; İbradı, Melik Mountain (Hasan Hill), stony hillsides, 2000 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4351; İbradı, Melik Plateau, stony paths, 1553 m, N 37°16'606" – E 31°24'539", 01.vii.2010, Çinbilgel 6789, Ir. – Tur., (H).
	<i>Asyneuma rigidum</i> (Willd.) Grossh. subsp. <i>rigidum</i>	İbradı, Çeşgar Plateau Location, stony slopes, 1659 m, N 37°17'402" – E 31°24'250", 17.vi.2008, Çinbilgel 4239; İbradı, Tahtalı Location, moist stony places, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4400; Manavgat, Dermebeli Location, stony places, 1880 m, N 37°12'698" – E 31°20'052", 20.vii.2008, Çinbilgel 4532, Ir. – Tur., (H).
	<i>Asyneuma rigidum</i> (Willd.) Grossh. subsp. <i>sibthorpiantum</i> (Roemer & Schultes) Damboldt	Manavgat, Kızılpınar Plateau, dwarf bush-thorny cushion formation, 1798 m, N 37°14'544" – E 31°20'315", 03.viii.2009, Çinbilgel 5624; Manavgat, Kızılpınar Plateau, dwarf bush-thorny cushion formation, 1754 m, N 37°14'090" – E 31°19'969", 04.viii.2009, Çinbilgel 5689; İbradı, Melik Plateau, dwarf bush-thorny cushion formation, 1503 m, N 37°16'190" – E 31°24'575", 01.vii.2010, Çinbilgel 6765, Ir. – Tur., (H).
	<i>Asyneuma virgatum</i> (Labill.) Bornm. subsp. <i>cichoriiforme</i> (Boiss.) Damboldt	İbradı, Çeşgar Plateau, rock, 1620 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4254, E. Medit., (H).
	<i>Asyneuma virgatum</i> (Labill.) Bornm. subsp. <i>virgatum</i>	Manavgat, Kocatop Mountain, windy hills, 2106 m, N 37°12'578" – E 31°18'867", 10.vi.2009, Çinbilgel 5360; Manavgat, Kızılpınar Plateau, stony slopes, 1908 m, N 37°14'355" – E 31°20'499", 03.viii.2009, Çinbilgel 5625, (H).
	<i>Campanula cymbalaria</i> Sibth. & Sm.	Manavgat, Around of Dermebeli Location, under the rock, 1944 m, N 37°13'667" – E 31°20'270", 10.vi.2008, Çinbilgel 3707; İbradı, Melik Mountain (Hasan Hill), rock, 2049 m, N 37°16'896" – E 31°23'156", 18.vi.2008, Çinbilgel 4316; Manavgat, Kocatop Mountain foothills (Ocakyeri

		Location), rock, 1782 m, N 37°12'226" – E 31°18'317", 11.vi.2009, Çinbilgel 5432, E. Medit., (mt.), (H).
	<i>Campanula involucrata</i> Aucher ex A.DC.	İbradı, Melik Mountain (Hasan Hill), slopes, 2000 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4343; Manavgat, Kızılıpınar Plateau, mixed <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1718 m, N 37°14'103" – E 31°19'803", 09.vi.2009, Çinbilgel 5246; İbradı, Çeşgar Plateau surroundings (Çeşgar Plateau Breach), stony places, 1682 m, N 37°17'288" – E 31°24'324", 01.vii.2010, Çinbilgel 6912, Ir. – Tur., (H).
	* <i>Campanula isaurica</i> Constandr., Quezel & A. Pamukçuoğlu	Manavgat, Kızılıpınar Plateau surroundings (Nurboğazı Location), rocky places, 1907 m, N 37°14'167" – E 31°20'459", 11.vi.2008, Çinbilgel 3836; Manavgat, Burunçalı Location, rock, 1815 m, N 37°09'107" – E 31°18'850", 02.vi.2009, Çinbilgel 5076; Manavgat, Kızılıpınar Plateau, rock, 1850 m, N 37°14'263" – E 31°20'414", 03.viii.2009, Çinbilgel 5626, End. E. Medit., (H).
	<i>Campanula lyrata</i> Lam. subsp. <i>lyrata</i>	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3483; İbradı, Suçikan Location way, under the forest – stony places, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4414; İbradı, Suçikan Location surroundings, <i>Cedrus libani</i> – <i>Pinus nigra</i> forest, 1271 m, N 37°14'863" – E 31°24'369", 02.vii.2010, Çinbilgel 7042, (H).
	<i>Campanula peregrina</i> L.	Manavgat, Çay Neighbourhood, Kızılıpınar Plateau way, streamsides, 628 m, 05.viii.2009, Çinbilgel 5734, E. Medit., (H).
	<i>Campanula phrygia</i> Jaub. & Spach	Manavgat, around of Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest, 702 m, N 37°09'905" – E 31°16'998", 06.vi.2011, Çinbilgel 7835, E. Medit., (Th).
	<i>Campanula propinqua</i> Fisch. & C.A.Mey.	Manavgat, Burunçalı Location (Aslanboğazı), stony places, 1687 m, N 37°08'939" – E 31°19'500", 02.vi.2009, Çinbilgel 5019; İbradı, around of Çeşgar Plateau Breach, stony places, 1672 m, N 37°17'262" – E 31°24'324", 01.vii.2010, Çinbilgel 6808; Manavgat, north of Çay Neighbourhood (Dikmen Hill), <i>Pinus brutia</i> forest, 816 m, N 37°12'492" – E 31°16'225", 07.vi.2011, Çinbilgel 7893, Ir. – Tur., (Th).
	<i>Campanula stricta</i> L. var. <i>libanotica</i> (A.DC.) Boiss.	Manavgat, Saritaş Location, rock, 1998 m, N 37°15'078" – E 31°20'867", 04.viii.2009, Çinbilgel 5668, Ir. – Tur., (Th).
	<i>Legousia falcata</i> (Ten.) Fritsch ex Janch.	Manavgat, Dutluca Plateau, under the rock – stony places, 1282 m, N 37°13'785" – E 31°18'213", 13.vi.2008, Çinbilgel 4063; Manavgat, Akpınar Plateau (Kelsu Location), watersides, 1573 m, N 37°09'507" – E 31°20'312", 01.vi.2009, Çinbilgel 4889; Manavgat, north of Çay Neighbourhood (Dikmen Hill), <i>Pinus brutia</i> forest, 816 m, N 37°12'492" – E 31°16'225", 07.vi.2011, Çinbilgel 7894, Medit., (Th).
	<i>Legousia speculum – veneris</i> (L.) Durende ex Vill.	İbradı, Akmuar Location, Juniper Forest clearings, 1036 m, N 37°11'773" – E 31°24'864", 21.v.2011, Çinbilgel 7681; İbradı, Akmuar Location, <i>Juniperus excelsa</i> forest, 1044 m, N 37°11'404" – E 31°24'996", 03.vii.2011, Çinbilgel 8033, Medit., (Th).
Capparaceae		
	<i>Capparis spinosa</i> L.	Manavgat, between Çay and Burmahan Neighbourhoods, <i>Pinus brutia</i> forest, 700 m, N 37°11'519" – E 31°16'218", 06.vi.2011, Çinbilgel 7774, (Ph).
Caprifoliaceae		
	* <i>Cephalaria lycica</i> V.A.Matthews	İbradı, Melik Mountain (eastern slopes of Hasan Hill), stony slopes, 1890 m, N 37°16'914" – E 31°23'287", 18.vi.2008, Çinbilgel 4285; Manavgat, Çandır Hill, dwarf bush-thorny cushion formation, 1987 m, N 37°14'492" – E 31°21'151", 04.viii.2009, Çinbilgel 5685; İbradı, Yukartoka Plateau, stony places, 1525 m, N 37°14'271" – E 31°22'440", 02.vii.2010, Çinbilgel 7258, End., E. Medit.(mt.), (H).
	<i>Knautia integrifolia</i> (L.) Bert. var. <i>bidens</i> (Sm.) Borbás	Manavgat, around of Burmahan Neighbourhood, roadsides, 670 m, N 37°10'760" – E 31°16'790", 06.vi.2011, Çinbilgel 7781; Manavgat, around of Burmahan Neighbourhood (Degirmenderesi Location), damp places, 685 m, N 37°10'723" – E 31°16'850", 06.vi.2011, Çinbilgel 7789, E. Medit., (Th).
	<i>Lonicera etrusca</i> Santi var. <i>etrusca</i>	İbradı, Akmuar Location, <i>Juniperus excelsa</i> forest, 1019 m, N 37°11'322" – E 31°25'008", 03.vii.2011, Çinbilgel 8032, Medit., (Ph).
	<i>Lonicera nummularifolia</i> Jaub. & Spach subsp. <i>nummularifolia</i>	Manavgat, Derme Plateau, rock, 1759 m, N 37°12'512" – E 31°20'675", 19.vii.2008, Çinbilgel 4482; İbradı, Boğazyurt Location, stony places, 1048 m, N 37°11'404" – E 31°25'002", 28.iii.2010, Çinbilgel 5958; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 980 m, N 37°10'766" – E 31°25'403", 05.iii.2011, Çinbilgel 7336, (Ph).
	<i>Pterocephalus pinardii</i> Boiss.	Manavgat, Kovacık Mountain surroundings, rocky places, 1887 m, N 37°14'043" – E 31°20'377", 09.vi.2008, Çinbilgel 3676; Manavgat, Kızılıpınar Plateau, rock, 1875 m, N 37°14'249" – E 31°20'465", 03.viii.2009, Çinbilgel 5620; İbradı, Çeşgar Plateau, rocky places, 1665 m, N 37°17'211" – E 31°24'325", 01.vii.2010, Çinbilgel 6871, End., E. Medit., (Ch).
	<i>Pterocephalus plumosus</i> (L.) Coulter	İbradı, Suçikan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7077; Manavgat, Genger Location, <i>Pinus brutia</i> forest, 1150 m, N 37°09'242" – E 31°17'835", 06.vi.2011, Çinbilgel 7871; İbradı, around of Çukuryurt Plateau, <i>Pinus nigra</i> forest, 1227 m, N 37°12'049" – E 31°24'169", 03.vii.2011, Çinbilgel 8003, (Th).

	<i>Scabiosa micrantha</i> Desf.	İbradı, Tahtalı Location, stony places, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4388; İbradı, Aşağıtoka Plateau, stony slopes, 1544 m, N 37°13'888" – E 31°22'352", 16.vi.2010, Çinbilgel 6255; İbradı, Akpınar Location, <i>Pinus nigra</i> forest clearings, 1047 m, N 37°11'563" – E 31°24'882", 02.vii.2010, Çinbilgel 7162, (Th).
	* <i>Scabiosa reuteriana</i> Boiss.	Manavgat, Kızılıpınar Plateau way, Kavaklı Location, stony places, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4045; İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1223 m, N 37°14'468" – E 31°24'259", 02.vii.2010, Çinbilgel 7054; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1006 m, N 37°11'054" – E 31°25'243", 03.vii.2011, Çinbilgel 8039, End., E. Medit., (Th).
	<i>Valeriana dioscoridis</i> Sm.	İbradı, Gevenli Location, under the <i>Pinus nigra</i> forest– under the rock, 1145 m, N 37°11'875" – E 31°24'457", 28.iii.2010, Çinbilgel 5937, E. Medit., (H).
	<i>Valerianella carinata</i> Loisel	İbradı, Melik Mountain (Orta Hill), windy hills, 2053 m, N 37°16'182" – E 31°22'761", 17.vi.2010, Çinbilgel 6319, (Th).
	<i>Valerianella costata</i> (Steven) Betcke	İbradı, Akmuar Location, Juniper Forest clearings, 1036 m, N 37°11'773" – E 31°24'864", 21.v.2011, Çinbilgel 7679, Medit., (Th).
	<i>Valerianella muricata</i> (Steven ex Roem. & Schult.) W.H.Baxter	İbradı, Toka Plateau Breach, meadows, 1676 m, N 37°13'774" – E 31°23'168", 21.v.2011, Çinbilgel 7662, (Th).
	<i>Valerianella turgida</i> (Steven) Betcke	İbradı, between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çinbilgel 2922; Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4836, (Th).
	<i>Valerianella vesicaria</i> (L.) Moench	İbradı, Söbüova Location (around of Akpınar Location), under the forest, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3551; İbradı, Akmuar Location, stony places, 1042 m, N 37°12'255" – E 31°24'780", 21.v.2011, Çinbilgel 7685, (Th).
Caryophyllaceae		
	* <i>Arenaria deflexa</i> Decne subsp. <i>microsepala</i> McNeill	İbradı, Suçikan Location way, stony rocky places, under the forest, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4424; İbradı, Yukarıtoka Plateau, rock, 1733 m, N 37°15'405" – E 31°22'545", 18.vi.2010, Çinbilgel 6474; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1508 m, N 37°15'558" – E 31°24'354", 30.vi.2010, Çinbilgel 6651, End., E. Medit., (H).
	* <i>Arenaria macrosepala</i> Boiss.	Manavgat, Kocatop Mountain, windy hills, 2174 m, N 37°12'467" – E 31°18'805", 10.vi.2009, Çinbilgel 5325, End., (Th).
	<i>Arenaria serpyllifolia</i> L. subsp. <i>leptoclados</i> (Rchb.) Nyman	İbradı, around of Yirme Hill, forest clearings, 1783 m, N 37°14'667" – E 31°23'097", 18.v.2008, Çinbilgel 3189; Manavgat, Kızılıpınar Plateau, mixed <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1718 m, N 37°14'103" – E 31°19'803", 09.vi.2009, Çinbilgel 5255; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'491" – E 31°24'478", 30.vi.2010, Çinbilgel 6694, (Th).
	<i>Arenaria serpyllifolia</i> L. subsp. <i>serpyllifolia</i>	İbradı, Güzle Plateau, Juniper Forest, 1306 m, N 37°09'817" – E 31°23'980", 20.iv.2008, Çinbilgel 2832; Manavgat, Akpınar Plateau (Yumrutaş Location), mixed <i>Cedrus libani</i> forest, 1520 m, N 37°09'840" – E 31°21'028", 19.v.2009, Çinbilgel 4791; İbradı, Çukuryurt Plateau surroundings, stony places, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6168, (Th).
	* <i>Arenaria tmolea</i> Boiss.	İbradı, Melik Mountain (Hasan Hill), stony hillsides, 1800 – 2000 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4341; Manavgat, Kovacık Mountain (Uzunçaşır Location), on the rock, 1966 m, N 37°13'620" – E 31°20'082", 29.vi.2009, Çinbilgel 5452; İbradı, Melik Mountain (Orta Hill), windy hills, 1999 m, N 37°16'161" – E 31°22'671", 17.vi.2010, Çinbilgel 6326, End., E. Medit. (mt.), (H).
	<i>Cerastium brachypetalum</i> Pers. subsp. <i>roeseri</i> (Boiss. & Heldr.) Nyman	İbradı, Güzle Plateau, Juniper forest, 1306 m, N 37°09'817" – E 31°23'980", 20.iv.2008, Çinbilgel 2839; Manavgat, Akpınar Plateau (Yumrutaş Location), under the mixed <i>Cedrus libani</i> forest, 1514 m, N 37°09'738" – E 31°21'024", 19.v.2009, Çinbilgel 4755; Manavgat, Eşekbeli Location, <i>Pinus nigra</i> forest, 1612 m, N 37°14'867" – E 31°19'864", 07.vi.2011, Çinbilgel 7913, (Th).
	<i>Cerastium dichotomum</i> L. subsp. <i>dichotomum</i>	Manavgat, Çiğdem Hill, rocky – stony slopes, 2015 m, N 37°15'429" – E 31°21'200", 11.vi.2008, Çinbilgel 3923; Manavgat, Akpınar Plateau, under the <i>Cedrus libani</i> forest, 1480 m, N 37°09'593" – E 31°21'076", 19.v.2009, Çinbilgel 4809; İbradı, Melik Mountain, dwarf bush-thorny cushion formation, 2244 m, N 37°16'042" – E 31°23'633", 18.vi.2010, Çinbilgel 6532, (Th).
	<i>Cerastium dubium</i> (Bastard) O.Schwarz	İbradı, southeast slopes of Yaylacık Mountain, <i>Pinus brutia</i> forest, 1089 m, N 37°09'775" – E 31°25'648", 19.iv.2009, Çinbilgel 4619; İbradı, Toka Plateau, meadows, 1528 m, N 37°14'074" – E 31°22'444", 09.v.2010, Çinbilgel 6096; İbradı, Aşağıtoka Plateau, meadows, 1520 m, N 37°14'220" – E 31°22'415", 20.v.2011, Çinbilgel 7642, (Th).
	<i>Cerastium fragillimum</i> Boiss.	İbradı, Toka Plateau, moist – stony places, 1522 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3340, (Th).
	<i>Cerastium glomeratum</i> Thuill.	Manavgat, Çiğdem Hill, rocky – stony slopes, 2015 m, N 37°15'429" – E 31°21'200", 11.vi.2008, Çinbilgel 3923; Manavgat, Burunçalı Location, slopes, 1883 m, N 37°09'116" – E 31°18'988", 02.vi.2009, Çinbilgel 5059; İbradı, Aşağıtoka Plateau, stony places, 1555 m, N 37°13'162" – E 31°22'666", 20.v.2011, Çinbilgel 7647, (Th).

	<i>Dianthus anatolicus</i> Boiss.	Manavgat, Dermebeli Location, dwarf bush-thorny cushion formation, 1880 m, N 37°12'698" – E 31°20'052", 20.vii.2008, Çinbilgel 4528; Kızılpınar Plateau (around of Dermebeli Location), stony places, 1904 m, N 37°13'703" – E 31°20'275", 10.x.2009, Çinbilgel 5776; İbradı, Yukarıtoka Plateau, stony places, 1525 m, N 37°14'271" – E 31°22'440", 02.vii.2010, Çinbilgel 7262, (H).
	<i>Dianthus crinitus</i> Sm. var. <i>crinitus</i>	Manavgat, Dermebeli Location (Artaşı), stony places, 1874 m, N 37°13'519" – E 31°20'317", 10.vi.2008, Çinbilgel 3811, (H).
	<i>Dianthus micranthus</i> Boiss. & Heldr.	Manavgat, Kızılpınar Plateau, stony places, 1750 m, N 37°14'071" – E 31°19'981", 19.vii.2008, Çinbilgel 4460; Manavgat, Kızılpınar Plateau, dwarf bush-thorny cushion formation, 1747 m, N 37°14'087" – E 31°19'983", 29.vi.2009, Çinbilgel 5504; İbradı, Yukarıtoka Plateau - Melik Mountain (Orta Hill) foothills, 1879 m, N 37°15'975" – E 31°22'558", 17.vi.2010, Çinbilgel 6288, (H).
	<i>Dianthus zonatus</i> Fenzl var. <i>zonatus</i>	Manavgat, Dermebeli Location (Artaşı), rock, 1885 m, N 37°13'490" – E 31°20'297", 19.vii.2008, Çinbilgel 4465; Manavgat, Dermebeli Location, rock, 1920 m, N 37°13'702" – E 31°20'304", 29.vi.2009, Çinbilgel 5531; İbradı, Çeşgar Plateau, rock, 1618 m, N 37°17'456" – E 31°23'710", 01.vii.2010, Çinbilgel 6846, (Ch).
	* <i>Eremogone isaurica</i> (Boiss.)Ikonn.	Manavgat, Sarıtaş Location, stony hillsides, 2074 m, N 37°15'015" – E 31°20'984", 11.vi.2008, Çinbilgel 3910; Kocatop Mountain, windy hills, 2080 m, N 37°12'398" – E 31°18'732", 10.vi.2009, Çinbilgel 5349; İbradı, Melik Mountain (Orta Hill), slopes, 2195 m, N 37°16'355" – E 31°23'060", 17.vi.2010, Çinbilgel 6401, End., (H).
	* <i>Eremogone ledebouriana</i> (Fenzl) Ikonn.	İbradı, Melik Mountain (Orta Hill), windy hills, 2053 m, N 37°16'182" – E 31°22'761", 17.vi.2010, Çinbilgel 6320; İbradı, Melik Mountain (Orta Hill), windy hills, 2183 m, N 37°16'210" – E 31°22'965", 17.vi.2010, Çinbilgel 6362; İbradı, Melik Mountain (Orta Hill), windy hills, 2187 m, N 37°16'210" – E 31°22'985", 17.vi.2010, Çinbilgel 6385, End., (H).
	* <i>Gypsophila curvifolia</i> Fenzl	Manavgat, Kovacık Mountain foothills (Gökçebel Location), dwarf bush-thorny cushion formation, 1695 m, N 37°13'984" – E 31°19'318", 11.vi.2009, Çinbilgel 5441; İbradı, Aşağıtoka Plateau, dwarf bush-thorny cushion formation, 1526 m, N 37°12'919" – E 31°22'639", 16.vi.2010, Çinbilgel 6187; İbradı, Aşağıtoka Plateau, stony places, 1508 m, N 37°13'377" – E 31°22'478", 02.vii.2011, Çinbilgel 7973, End., E. Medit. (mt.), (Ch).
	<i>Herniaria glabra</i> L.	Manavgat, Susuzkır Hill surroundings, open places, 1666 m, N 37°16'797" – E 31°21'755", 12.vi.2008, Çinbilgel 3983; Manavgat, Kızılpınar Plateau (Kocaboğaz Location), arid meadows, 1790 m, N 37°13'949" – E 31°20'094", 29.vi.2009, Çinbilgel 5472; İbradı, Aşağıtoka Plateau, dwarf bush-thorny cushion formation, 1526 m, N 37°12'919" – E 31°22'639", 16.vi.2010, Çinbilgel 6185, (H).
	<i>Herniaria incana</i> Lam.	Manavgat, Kepezbeli Location, stony slopes, 1641 m, N 37°10'300" – E 31°21'722", 19.v.2009, Çinbilgel 4772; Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), dwarf bush-thorny cushion formation, 1771 m, N 37°13'824" – E 31°19'246", 09.vi.2009, Çinbilgel 5264; İbradı, Aşağıtoka Plateau, stony places, 1530 m, N 37°12'883" – E 31°22'500", 16.vi.2010, Çinbilgel 6217, (H).
	<i>Holosteum umbellatum</i> L. var. <i>umbellatum</i>	İbradı, southern slopes of Yaylacık Mountain, stony places, 1541 m, N 37°10'262" – E 31°24'049", 18.iv.2009, Çinbilgel 4571; İbradı, southeast slopes of Yaylacık Mountain, under the <i>Cedrus libani</i> forest, 1356 m, N 37°09'929" – E 31°25'029", 19.iv.2009, Çinbilgel 4589; İbradı, Çukuryurt Plateau, meadows, 1268 m, N 37°12'117" – E 31°24'023", 28.iii.2010, Çinbilgel 5927, (Th).
	<i>Minuartia anatolica</i> (Boiss.) Woronow var. <i>polymorpha</i> McNeill	İbradı, Melik Mountain (Hasan Hill), stony slopes, 2136 m, N 37°16'753" – E 31°23'122", 18.vi.2008, Çinbilgel 4332; Manavgat, Sürüçkan Mountain, stony slopes, 2107 m, N 37°12'335" – E 31°19'690", 30.vi.2009, Çinbilgel 5554; İbradı, Toka Plateau, stony slopes, 1546 m, N 37°14'038" – E 31°22'783", 09.v.2010, Çinbilgel 6071, (Ch).
	<i>Minuartia erythrosepala</i> (Boiss.) Hand.–Mazz. var. <i>erythrosepala</i>	İbradı, Melik Mountain (Hasan Hill), stony hillsides, 2000 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4349; İbradı, Melik Mountain (Hasan Hill), slopes, 2076 m, N 37°16'581" – E 31°23'005", 17.vi.2010, Çinbilgel 6423; İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings, 1568 m, N 37°13'549" – E 31°23'433", 03.vii.2011, Çinbilgel 7997, (Ch).
	<i>Minuartia globulosa</i> (Labill.) Schinz & Thell.	Manavgat, Çiğdem Hill surroundings, dolines, 2001 m, N 37°14'664" – E 31°21'051", 04.viii.2009, Çinbilgel 5654; İbradı, Melik Mountain (Hasan Hill), dwarf bush-thorny cushion formation, 1840 m, N 37°16'441" – E 31°22'564", 17.vi.2010, Çinbilgel 6456; Manavgat, Çay Neighbourhood, Eşekbeli Location, stony places, 1589 m, N 37°14'740" – E 31°19'747", 07.vi.2011, Çinbilgel 7915, E. Medit., (Th).
	<i>Minuartia hamata</i> (Hausskn.) Mattf.	İbradı, Yukarıtoka Plateau (between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3096; Manavgat, west of Kocatop Mountain (Örencik Location), dwarf bush-thorny cushion formation, 1861 m, N 37°11'829" – E 31°18'607", 11.vi.2009, Çinbilgel 5421; İbradı, Yukarıtoka Plateau, dwarf bush-thorny cushion

		formation, 1760 m, N 37°15'903" – E 31°22'402", 17.vi.2010, Çinbilgel 6281, (Th).
	<i>Minuartia juniperina</i> (L.) Maire & Petitm.	Manavgat, Burunçalı Location, rocky places, 1660 m, N 37°08'970" – E 31°18'852", 02.vi.2009, Çinbilgel 4969; Manavgat, Dermebeli Location, rock, 1920 m, N 37°13'702" – E 31°20'304", 29.vi.2009, Çinbilgel 5533; İbradı, Melik Mountain, windy hills, 2256 m, N 37°16'044" – E 31°23'646", 18.vi.2010, Çinbilgel 6536, (Ch).
	* <i>Minuartia leucocephala</i> (Boiss.) Mattf.	İbradı, Yukarıtoka Plateau (Between Beydiğin Plateau and Göcebe Location), stony places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çinbilgel 3108; Manavgat, Kaldırım Mountain (Aktünek Hill), windy hills, 2198 m, N 37°10'556" – E 31°20'329", 03.vi.2009, Çinbilgel 5130; İbradı, Melik Mountain (Hasan Hill), stony slopes, 2079 m, N 37°16'594" – E 31°23'035", 17.vi.2010, Çinbilgel 6418, End., Medit., (Ch).
	<i>Minuartia mediterranea</i> (Link) K.Maly	İbradı, Güzle Plateau, Juniper Forest, 1306 m, N 37°09'817" – E 31°23'980", 20.iv.2008 Çinbilgel 2832, Medit., (Th).
	<i>Minuartia mesogitana</i> (Boiss.) Hand. – Mazz. subsp. <i>kotschyana</i> (Boiss.) McNeill	İbradı, Söbüova Location (around of Akpınar Location), under the forest, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3540; Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4834; İbradı, Boğazyurt Location, stony places, 974 m, N 37°10'338" – E 31°25'772", 19.v.2011, Çinbilgel 7585, (Th).
	<i>Minuartia mesogitana</i> (Boiss.) Hand.–Mazz. subsp. <i>mesogitana</i>	İbradı, Çukuryurt Plateau surroundings, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6169; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7178; İbradı, Boğazyurt Location, stony places, 974 m, N 37°10'338" – E 31°25'772", 19.v.2011, Çinbilgel 7586, E. Medit., (Th).
	<i>Minuartia meyeri</i> (Boiss.) Bornm.	Manavgat, Kaldırım Mountain (Hıncırak Location), mixed <i>Cedrus libani</i> forest, 1475 m, N 37°09'368" – E 31°20'410", 01.vi.2009, Çinbilgel 4898; Manavgat, Kocatop Mountain (Koca Hill = Köygöründüğü Location), windy hills, 2150 m, N 37°12'346" – E 31°18'633", 10.vi.2009, Çinbilgel 5342; İbradı, Yukarıtoka Plateau surroundings, <i>Pinus nigra</i> forest clearing, 1783 m, N 37°16'349" – E 31°22'313", 17.vi.2010, Çinbilgel 6464, Ir. – Tur., (Th).
	<i>Minuartia multinervis</i> (Boiss.) Bornm.	Manavgat, around of Çağlan Breach, debris and stony, 1665 m, N 37°15'851" – E 31°21'000", 12.vi.2008, Çinbilgel 3930; Manavgat, Akpınar Plateau(Hıncırak Location), rock debris, 1512 m, N 37°09'442" – E 31°20'412", 01.vi.2009, Çinbilgel 4915; İbradı, Melik Mountain (Orta Hill), rocky places, 1967 m, N 37°16'123" – E 31°22'618", 17.vi.2010, Çinbilgel 6316, (Th).
	* <i>Minuartia umbellulifera</i> (Boiss.) McNeill subsp. <i>umbellulifera</i> var. <i>umbellulifera</i>	Manavgat, around of Burunçalı Location, rocky places, 1581 m, N 37°08'893" – E 31°18'959", 02.vi.2009, Çinbilgel 5087, End., (Ch).
	<i>Moenchia mantica</i> (L.) Bartl. subsp. <i>caerulea</i> (Boiss.) Clapham	İbradı, Eynif Plain, arid meadows, 949 m, N 37°08'004" – E 31°27'963", 19.v.2011, Çinbilgel 7573; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 990 m, N 37°10'778" – E 31°25'386", 21.v.2011, Çinbilgel 7695, (Th).
	* <i>Paronychia argyroloba</i> Satpf	Manavgat, Kocatop Mountain (Koca Hill = Köygöründüğü Location), windy hills, 2150 m, N 37°12'346" – E 31°18'633", 10.vi.2009, Çinbilgel 5335; İbradı, Melik Mountain (Orta Hill), windy hills, 2187 m, N 37°16'210" – E 31°22'985", 17.vi.2010, Çinbilgel 6383; İbradı, Aşağıtoka Plateau, rocky hills, 1538 m, N 37°12'892" – E 31°22'648", 02.vii.2011, Çinbilgel 7979, End., (H).
	<i>Petrorhagia dubia</i> (Raf.) G.López & Romo	İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7169; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest clearings, 965 m, N 37°10'640" – E 31°25'709", 19.v.2011, Çinbilgel 7594; Manavgat, around of Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest, 702 m, N 37°09'905" – E 31°16'998", 06.vi.2011, Çinbilgel 7839, (Th).
	<i>Saponaria pumilio</i> Boiss.	Manavgat, Çandır Hill, stony slopes, 2034 m, N 37°14'500" – E 31°21'087", 11.vi.2008, Çinbilgel 3872; İbradı, Melik Mountain (Hasan Hill), stony hillsides, 2049 m, N 37°16'896" – E 31°23'156", 18.vi.2008, Çinbilgel 4318; Manavgat, eastern hillsides of Kaldırım Mountain, windy hills, 2026 m, N 37°10'434" – E 31°20'702", 03.vi.2009, Çinbilgel 5161, E. Medit., (Ch).
	<i>Silene aegyptiaca</i> (L.) L.f. subsp. <i>aegyptiaca</i>	Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4663; İbradı, around of Gevenli Location, under the rock, 1145 m, N 37°11'947" – E 31°24'445", 28.iii.2010, Çinbilgel 5938; İbradı, Melik Mountain (Orta Hill), slopes, 2195 m, N 37°16'355" – E 31°23'060", 17.vi.2010, Çinbilgel 6400, (Th).
	<i>Silene behen</i> L.	Manavgat, Burunçalı Location (Aslanboğazı), 1703 m, N 37°08'958" – E 31°19'504", 02.vi.2009, Çinbilgel 5045, (Th).
	* <i>Silene capitellata</i> Boiss.	İbradı, Toka Plateau, moist stony places, 1522 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3343; İbradı, Toka Plateau, stony places, 1520 m, N 37°14'038" – E 31°22'783", 09.v.2010, Çinbilgel 6087; İbradı, around of Toka Plateau Breach, <i>Cedrus libani</i> – <i>Pinus nigra</i> forest, 1679 m, N 37°13'808" – E 31°23'202", 21.v.2011, Çinbilgel 7673, End., Ir. – Tur., (Ch).
	* <i>Silene caramanica</i> Boiss. & Heldr. var. <i>ilarstanii</i> Aytaç & Dural	Manavgat, Kızılıpınar Plateau, dwarf bush-thorny cushion formation, 1786 m, N 37°14'176" – E 31°20'229", 03.viii.2009, Çinbilgel 5628, End., Ir. – Tur., (H).

	<i>Silene compacta</i> Fisch. ex Hornem	Manavgat, Arğıça Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4130, (H).
	<i>Silene dichotoma</i> Ehrh. subsp. <i>dichotoma</i>	İbradı, around of Ekinlik Location, <i>Cedrus libani</i> – <i>Pinus nigra</i> forest, 1439 m, N 37°12'987" – E 31°23'789", 20.v.2011, Çinbilgel 7629, (H).
	<i>Silene gigantea</i> L. var. <i>incana</i>	İbradı, Gevenli Location, under the <i>Pinus brutia</i> forest, 1108 m, N 37°12'316" – E 31°24'613", 17.v.2008, Çinbilgel 3014; Manavgat, between Akin – Karahan Locations, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çinbilgel 4148, E. Medit., (H).
	<i>Silene italica</i> (L.) Pers. subsp. <i>italica</i>	İbradı, Ekinlik Location surroundings, <i>Cedrus libani</i> forest (mixed with <i>Pinus nigra</i>), 1532 m, N 37°13'668" – E 31°23'510", 09.v.2010, Çinbilgel 6140; İbradı, Çukuryurt Plateau surroundings, <i>Pinus nigra</i> forest, 1357 m, N 37°12'461" – E 31°23'914", 09.v.2010, Çinbilgel 6159, Medit., (H).
	* <i>Silene lycaonica</i> Chowdhuri	Manavgat, Kızılpınar Plateau, dwarf bush-thorny cushion formation, 1754 m, N 37°14'090" – E 31°19'969", 04.viii.2009, Çinbilgel 5687; İbradı, Melik Mountain (Hasan Hill), dwarf bush-thorny cushion formation, 2085 m, N 37°16'595" – E 31°22'998", 17.vi.2010, Çinbilgel 6428; İbradı, Melik Plateau, rocky places, 1425 m, N 37°15'482" – E 31°24'716", 02.vii.2010, Çinbilgel 6980, End., Ir. – Tur., (H).
	<i>Silene macrodonta</i> Boiss.	İbradı, between Akpınar Location and Suçikan Location, 1091 m, N 37°13'075" – E 31°24'456", 18.vi.2008, Çinbilgel 4431, (Th).
	<i>Silene odontopetala</i> Fenzl	İbradı, Çeşgar Plateau, on the rock, 1642 m, N 37°17'662" – E 31°23'989", 17.vi.2008, Çinbilgel 4193; Manavgat, Kızılpınar Plateau, Nurboğazi Location, rock, 1913 m, N 37°14'166" – E 31°20'469", 04.viii.2009, Çinbilgel 5674; İbradı, Melik Mountain (Orta Hill), stony places, 2084 m, N 37°16'414" – E 31°23'077", 17.vi.2010, Çinbilgel 6406, (Ch).
	* <i>Silene oreades</i> Boiss. & Heldr.	İbradı, Melik Mountain (Çeşgar Hill), screes, 2148 m, N 37°16'119" – E 31°23'536", 18.vi.2010, Çinbilgel 6501, End., E. Medit. (mt.), (H).
	<i>Silene rhynchocarpa</i> Boiss.	İbradı, hillsides of Melik Mountain (Çatal düzlüğü), stony slopes, 1843 m, N 37°15'258" – E 31°23'363", 18.v.2008, Çinbilgel 3288; İbradı, Çeşgar Plateau, stony slopes, 1659 m, N 37°17'402" – E 31°24'250", 17.vi.2008, Çinbilgel 4245; İbradı, Melik Mountain (Orta Hill), stony places, 2084 m, N 37°16'414" – E 31°23'077", 17.vi.2010, Çinbilgel 6407, E. Medit., (H).
	<i>Silene spergulifolia</i> (Desf.) M.Bieb.	İbradı, around of Yirme Hill, forest clearings, 1674 m, N 37°14'298" – E 31°23'004", 18.v.2008, Çinbilgel 3200; Manavgat, Kocatop Mountain, windy hills, 2106 m, N 37°12'578" – E 31°18'867", 10.vi.2009, Çinbilgel 5358; İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1048 m, N 37°12'283" – E 31°24'732", 19.v.2011, Çinbilgel 7615, Ir. – Tur., (H).
	<i>Silene subconica</i> Friv.	İbradı, Melik Mountain (southeastern hillsides of Hasan Hill), stony slopes, 1937 m, N 37°16'885" – E 31°23'249", 18.vi.2008, Çinbilgel 4295; İbradı, Toka Plateau, <i>Pinus nigra</i> forest clearings, 1577 m, N 37°14'030" – E 31°22'843", 09.v.2010, Çinbilgel 6103; İbradı, Gevenli Location, <i>Pinus brutia</i> forest clearings, 1102 m, N 37°12'102" – E 31°24'531", 19.v.2011, Çinbilgel 7621, (Th).
	<i>Silene vulgaris</i> (Moench) Garcke var. <i>vulgaris</i>	İbradı, Toka Plateau (=Beydiğın Plateau), <i>Pinus nigra</i> forest clearings-stony rocky places, 1544 m, N 37°14'435" – E 31°22'650", 18.v.2008, Çinbilgel 3164; Manavgat, Düğnük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4077; İbradı, Melik Plateau, dwarf bush-thorny cushion formation, 1523 m, N 37°16'395" – E 31°24'482", 01.vii.2010, Çinbilgel 6784, (H).
	<i>Spergularia rubra</i> (L.) J. Presl & C. Presl	Manavgat, Kızılpınar Plateau, stony places, 1750 m, N 37°14'071" – E 31°19'981", 19.vii.2008, Çinbilgel 4458; Manavgat, Kızılpınar Plateau (Kocaboğaz Location), arid meadows, 1790 m, N 37°13'949" – E 31°20'094", 29.vi.2009, Çinbilgel 5476; Manavgat, Kepez Plateau, meadows, 1631 m, N 37°11'208" – E 31°21'007", 11.x.2009, Çinbilgel 5844, (H).
	<i>Stellaria cilicica</i> Boiss. & Balansa	İbradı, southeast slopes of Yaylacık Mountain, under the <i>Cedrus libani</i> forest, 1356 m, N 37°09'929" – E 31°25'029", 19.iv.2009, Çinbilgel 4593; Manavgat, Burmahan Neighbourhood, around of Erenli Location, streamsides, 753 m, N 37°09'844" – E 31°16'991", 26.iii.2011, Çinbilgel 7440; Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, Putraklı Location, 734 m, N 37°12'249" – E 31°16'454", 27.iii.2011, Çinbilgel 7569, (Th).
	<i>Stellaria media</i> (L.) Vill.	İbradı, around of Yirme Hill, <i>Cedrus libani</i> forest, 1783 m, N 37°14'667" – E 31°23'097", 18.v.2008, Çinbilgel 3177; Manavgat, Akpınar Plateau (Yumrutaş Location), mixed <i>Cedrus libani</i> forest, 1520 m, N 37°09'840" – E 31°21'028", 19.v.2009, Çinbilgel 4788; İbradı, around of Yirme Hill, <i>Pinus nigra</i> forest, 1550 m, N 37°14'328" – E 31°22'786", 08.v.2010, Çinbilgel 6026, (Th).
	<i>Telephium imperati</i> L. subsp. <i>orientale</i> (Boiss.) Nyman	Manavgat, Dermebeli Location (Arıtışı), on the rock and stony places, 1874 m, N 37°13'519" – E 31°20'317", 10.vi.2008, Çinbilgel 3821; Manavgat, around of Burunçalı Location, debris, 1857 m, N 37°09'063" – E 31°19'473", 02.vi.2009, Çinbilgel 5011; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çinbilgel 6715, (H).
	<i>Vaccaria hispanica</i> (Mill.) Rauschert	Manavgat, Arğıça Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4135; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest clearings (roadsides), 655 m, N 37°11'683" – E 31°16'103", 06.vi.2011, Çinbilgel 7753, (Th).

	<i>Velezia rigida</i> L.	Manavgat, Susuzkır Hill surroundings, stony places, 1706 m, N 37°16'425" – E 31°21'369", 12.vi.2008, Çinbilgel 3963; İbradı, Melik Plateau, forest clearings – stony places, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6559; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 983 m, N 37°10'973" – E 31°25'293", 03.vii.2011, Çinbilgel 8044, (Th).
Cistaceae		
	<i>Cistus creticus</i> L.	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, <i>Pinus brutia</i> forest, 628 m, N 37°11'815" – E 31°16'212", 10.x.2009, Çinbilgel 5743; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7730, Omni – Medit., (Ch).
	<i>Cistus salviifolius</i> L.	Manavgat, Çay Neighbourhood, Arğıçca Breach Location, burning area, 1052 m, N 37°12'995" – E 31°17'113", 07.vi.2011, Çinbilgel 7929, (Ch).
	<i>Fumana thymifolia</i> (L.) Spach	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 622 m, N 37°11'707" – E 31°16'064", 06.vi.2011 Çinbilgel 7744; Manavgat, Çay Neighbourhood, <i>Pinus brutia</i> forest, 630 m, N 37°11'879" – E 31°16'167", 07.vi.2011, Çinbilgel 7942, Medit., (H).
	<i>Helianthemum nummularium</i> (L.) Mill. subsp. <i>nummularium</i>	İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1048 m, N 37°12'283" – E 31°24'732", 19.v.2011, Çinbilgel 7618; İbradı, Toka Plateau way, <i>under the Pinus nigra</i> forest, 1600 m, N 37°13'955" – E 31°22'921", 08.vi.2011, Çinbilgel 7959, (H).
	<i>Helianthemum oelandicum</i> (L.) DC. subsp. <i>incanum</i> (Willk.) G.López	İbradı, Toka Plateau, moist stony places, 1522 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3346; İbradı, Toka Plateau, stony places, 1500 m, N 37°14'332" – E 31°22'569", 08.v.2010, Çinbilgel 6052; İbradı, Toka Plateau, stony places, 1538 m, N 37°14'038" – E 31°22'783", 09.v.2010, Çinbilgel 6079, (H).
	<i>Helianthemum salicifolium</i> (L.) Mill.	İbradı, Söbüova Location (around of Akpınar Location), under the forest, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3539; İbradı, Akmuar Location, Juniper Forest, 1047 m, N 37°11'401" – E 31°25'006", 21.v.2011, Çinbilgel 7689, (Th).
	<i>Helianthemum tomentosum</i> Gray	İbradı, between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çinbilgel 2930; İbradı, Çeşgar Plateau, stony slopes, 1659 m, N 37°17'402" – E 31°24'250", 17.vi.2008, Çinbilgel 4240; İbradı, Çeşgar Plateau, rocky places, 1665 m, N 37°17'211" – E 31°24'325", 01.vii.2010, Çinbilgel 6869, (H).
Cleomaceae		
	<i>Cleome ornithopodioides</i> L.	İbradı, Akmuar Location, streamsides, 1044 m, N 37°11'719" – E 31°24'820", 02.vii.2010, Çinbilgel 7221, E. Medit., (Th).
Colchicaceae		
	<i>Colchicum dolichantherum</i> K.Perss.	İbradı, between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çinbilgel 2921; İbradı, Toka Plateau, stony rocky places, 1544 m, N 37°14'435" – E 31°22'650", 18.v.2008, Çinbilgel 3167; İbradı, Çukuryurt Plateau, under the forest, 1286 m, N 37°12'113" – E 31°24'003", 02.vii.2010, Çinbilgel 7250, E. Medit., (G).
	<i>Colchicum triphyllum</i> Kunze	İbradı, Çukuryurt Plateau, meadows, 1267 m, N 37°12'129" – E 31°24'049", 05.iii.2011, Çinbilgel 7346, Medit., (G).
	<i>Colchicum variegatum</i> L.	Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4653; Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, <i>Pinus brutia</i> forest clearings – stony places, 628 m, N 37°11'815" – E 31°16'212", 10.x.2009, Çinbilgel 5741; İbradı, Melik Plateau, <i>Pinus nigra</i> forest clearings, 1431 m, N 37°15'838" – E 31°24'585", 05.x.2010, Çinbilgel 7290, E. Medit., (G).
Convolvulaceae		
	<i>Convolvulus arvensis</i> L.	Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4118; Manavgat, Kovacık Mountain (Gökçebel Location), dwarf bush-thorny cushion formation, 1709 m, N 37°14'032" – E 31°19'421", 09.vi.2009, Çinbilgel 5223; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çinbilgel 6718, (H).
	<i>Convolvulus cantabrica</i> L.	İbradı, Aşağıtoka Plateau, stony places, 1534 m, N 37°12'825" – E 31°22'485", 16.vi.2010, Çinbilgel 6222, Medit., (H).
	<i>Convolvulus libanoticus</i> Boiss.	Manavgat, around of Susuzkır Hill, dolines, stony places, 1666 m, N 37°16'797" – E 31°21'755", 12.vi.2008, Çinbilgel 3982; İbradı, Aşağıtoka Plateau, stony places, 1531 m, N 37°12'882" – E 31°22'682", 16.vi.2010, Çinbilgel 6243; İbradı, Aşağıtoka Plateau, dwarf bush formation, 1517 m, N 37°12'759" – E 31°22'661", 02.vii.2011, Çinbilgel 7985, E. Medit., (H).
	<i>Cuscuta palaestina</i> Boiss.	İbradı, Melik Mountain (Hasan Hill), slopes, 2076 m, N 37°16'581" – E 31°23'005", 17.vi.2010, Çinbilgel 6425; İbradı, Toka Plateau way, Belova Location, stony places, 1660 m, N 37°13'649" – E 31°23'311", 02.vii.2010, Çinbilgel 7254 (on <i>Sideritis erythrantha</i> var. <i>erythrantha</i>), (Vp).
Cornaceae		
	<i>Cornus mas</i> L.	Manavgat, Burmahan Neighbourhood, Erenli Location, streamsides, 713 m, N 37°10'048" – E 31°16'974", 11.x.2009, Çinbilgel 5811, Euro-Sib., (Ph).
Crassulaceae		

	<i>*Prometheum chrysanthum</i> (Boiss.) 't Hart	Manavgat, Derme Plateau, rocky places, 1740 m, N 37°12'694" – E 31°20'565", 19.vii.2008, Çinbilgel 4490; Dermebeli Location (Artaşı), on the rock, 1910 m, N 37°13'547" – E 31°20'315", 30.vi.2009, Çinbilgel 5598; İbradı, Çeşgar Plateau Breach, stony places, 1648 m, N 37°17'056" – E 31°24'397", 01.vii.2010, Çinbilgel 6796, End., E. Medit. (mt.), (Ch).
	<i>Prometheum sempervivoides</i> (Fischer ex M.Bieb.) H.Ohba	İbradı, Melik Mountain (Hasan Hill), stony slopes, 2136 m, N 37°16'753" – E 31°23'122", 18.vi.2008, Çinbilgel 4329; Manavgat, Çandır Hill, windy hills, 1987 m, N 37°14'492" – E 31°21'151", 04.viii.2009, Çinbilgel 5678; İbradı, Melik Mountain (Orta Hill), windy hills, 2059 m, N 37°16'461" – E 31°23'118", 17.vi.2010, Çinbilgel 6402, (Ch).
	<i>*Rosularia sempervivum</i> (M.Bieb.) A.Berger subsp. <i>glaucophylla</i> Egglı	İbradı, Melik Mountain (eastern slopes of Hasan Hill), rocky slopes, 2020 m, N 37°16'897" – E 31°23'181", 18.vi.2008, Çinbilgel 4301; Manavgat, Sulucan Hill, rock, 2081 m, N 37°11'492" – E 31°18'857", 11.vi.2009, Çinbilgel 5394; İbradı, Çeşgar Plateau Breach, stony places, 1648 m, N 37°17'056" – E 31°24'397", 01.vii.2010, Çinbilgel 6797, End., E. Medit., (Ch).
	<i>Sedum acre</i> L. subsp. <i>acre</i>	Manavgat, around of Çağlan Breach, stony places, 1681 m, N 37°16'230" – E 31°21'349", 12.vi.2008, Çinbilgel 3972; İbradı, Çeşgar Plateau, stony places, 1617 m, N 37°17'485" – E 31°23'565", 17.vi.2008, Çinbilgel 4220; İbradı, Çeşgar Plateau, stony places, 1619 m, N 37°17'489" – E 31°23'946", 01.vii.2010, Çinbilgel 6885, (H).
	<i>Sedum album</i> L.	Manavgat, Kaldırım Mountain, stony places, 1712 m, N 37°09'693" – E 31°20'284", 01.vi.2009, Çinbilgel 4936; Manavgat, Dermebeli Location, rock, 1928 m, N 37°13'717" – E 31°20'306", 29.vi.2009, Çinbilgel 5517; İbradı, Çeşgar Plateau, rock, 1623 m, N 37°17'463" – E 31°23'699", 01.vii.2010, Çinbilgel 6850, (Ch).
	<i>Sedum amplexicaule</i> DC. subsp. <i>tenuifolium</i> (Sm.) Greuter & Burdet	İbradı, Suçukan Location way, damp places, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2748; Manavgat, Akpınar Plateau (Sumaklı Location), mixed <i>Cedrus libani</i> forest, 1517 m, N 37°08'847" – E 31°19'742", 02.vi.2009, Çinbilgel 5025; İbradı, around of Çeşgar Plateau Breach, dwarf bush formation, 1681 m, N 37°17'317" – E 31°24'321", 01.vii.2010, Çinbilgel 6820, Medit., (Ch).
	<i>Sedum hispanicum</i> L.	Manavgat, Kaldırım Mountain, stony places, 1839 m, N 37°09'906" – E 31°20'407", 01.vi.2009, Çinbilgel 4951; Manavgat, Sürüçkan Mountain, windy hills, 2107 m, N 37°12'335" – E 31°19'690", 30.vi.2009, Çinbilgel 5558; İbradı, Yukarıtoka Plateau, rock, 1733 m, N 37°15'405" – E 31°22'545", 18.vi.2010, Çinbilgel 6473, Ir. – Tur., (Th).
	<i>Sedum laconicum</i> Boiss. & Heldr.	İbradı, around of Çeşgar Plateau Breach, stony places, 1672 m, N 37°17'262" – E 31°24'324", 01.vii.2010, Çinbilgel 6803, Medit., (Ch).
	<i>Sedum litoreum</i> Guss. var. <i>litoreum</i>	Manavgat, Akpınar Plateau way, <i>Pinus brutia</i> forest, 1310 m, N 37°08'495" – E 31°18'855", 06.vi.2011, Çinbilgel 7872, Medit., (Th).
	<i>Sedum magellense</i> Ten.	Manavgat, Dermebeli Location (Artaşı), on the rock and stony places, 1874 m, N 37°13'519" – E 31°20'317", 10.vi.2008, Çinbilgel 3813; Manavgat, Kovacık Mountain, rock, 1952 m, N 37°13'631" – E 31°20'057", 29.vi.2009, Çinbilgel 5487; İbradı, Melik Mountain (Orta Hill), windy hills, 2126 m, N 37°16'348" – E 31°23'067", 17.vi.2010, Çinbilgel 6392, Medit., (mt.), (H).
	<i>Sedum pallidum</i> M.Bieb.	İbradı, Toka Plateau (Göçebe Location), stony places, 1551 m, N 37°14'763" – E 31°22'308", 17.v.2008, Çinbilgel 3142; Manavgat, Kaldırım Mountain, 1818 m, N 37°09'957" – E 31°20'583", 03.vi.2009, Çinbilgel 5097; Manavgat, Kızılpınar Plateau, dwarf bush formation, 1747 m, N 37°14'087" – E 31°19'983", 29.vi.2009, Çinbilgel 5501, (H).
	<i>Sedum rubens</i> L.	Manavgat, Kızılpınar Plateau, stony paths, 1742 m, N 37°14'095" – E 31°19'986", 09.vi.2008, Çinbilgel 3642/a, Medit., (H).
	<i>Umbilicus horizontalis</i> DC.	Manavgat, around of Burmahan Neighbourhood, roadsides, 700 m, N 37°10'623" – E 31°16'847", 06.vi.2011, Çinbilgel 7794, E. Medit., (Ch).
	<i>Umbilicus luteus</i> (Huds.) Webb & Berthel.	Manavgat, Dermebeli Location (Artaşı), on the rock and stony places, 1874 m, N 37°13'519" – E 31°20'317", 10.vi.2008, Çinbilgel 3812; Manavgat, Dermebeli Location (Artaşı), rock, 1823 m, N 37°13'506" – E 31°20'339", 30.vi.2009, Çinbilgel 5544; İbradı, around of Çeşgar Plateau, rocky hills, 1631 m, N 37°17'458" – E 31°24'005", 01.vii.2010, Çinbilgel 6887, (Ch).
Cucurbitaceae		
	<i>Bryonia multiflora</i> Boiss. & Heldr.	İbradı, around of Yirme Hill, damp places, 1814 m, N 37°14'700" – E 31°23'139", 18.v.2008, Çinbilgel 3222, Ir. – Tur., (H).
Cyperaceae		
	<i>*Carex coriogyne</i> Nelmess	İbradı, between Gavur Breach and Etülgenboğazi, stony places, 1186 m, N 37°09'074" – E 31°25'116", 20.iv.2008, Çinbilgel 2977; Manavgat, Derme Plateau, stony hillsides, 1827 m, N 37°13'386" – E 31°20'237", 10.vi.2008, Çinbilgel 3798; İbradı, Aşağıtoka Plateau, moist stony places, 1555 m, N 37°13'162" – E 31°22'666", 20.v.2011, Çinbilgel 7653, End., E. Medit., (H).
	<i>Carex divisa</i> Huds.	Manavgat, Kızılpınar Plateau, watersides, 1805 m, N 37°14'086" – E 31°20'164", 09.vi.2008, Çinbilgel 3702; Manavgat, Kızılpınar Plateau, watersides, 1780 m, N 37°14'087" – E 31°20'158", 03.viii.2009, Çinbilgel 5639; İbradı, Toka Plateau Breach, meadows, 1676 m, N 37°13'774" – E 31°23'168", 21.v.2011, Çinbilgel 7665, Euro-Sib., (H).

	<i>Carex flacca</i> Schreb. subsp. <i>erythrostrachys</i> (Hoppe) Holub	İbradı, Melik Plateau surroundings (towards Suçikan Location), wet places, 1340 m, N 37°15'336" – E 31°24'546", 02.vii.2010, Çinbilgel 7025; İbradı, Melik Plateau surroundings (towards Suçikan Location), <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7035, Medit., (H).
	<i>Carex muricata</i> L. subsp. <i>muricata</i>	Manavgat, Kızılpınar Plateau way, between Akin – Karahan Locations, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çinbilgel 4146; İbradı, Melik Plateau, streamsides, 1537 m, N 37°15'549" – E 31°24'633", 02.vii.2010, Çinbilgel 6946, Euro-Sib., (H).
	<i>Carex stenophylla</i> Wahlenb. subsp. <i>stenophylloides</i> (V.I.Krecz.) Egorova	Manavgat, Kepez Plateau, meadows, 1628 m, N 37°11'110" – E 31°21'021", 03.vi.2009, Çinbilgel 5177; İbradı, Melik Plateau, meadows, 1446 m, N 37°15'954" – E 31°24'591", 01.vii.2010, Çinbilgel 6752, Ir. – Tur., (H).
	<i>Scirpioides holoschoneus</i> (L.) Soják subsp. <i>holoschoneus</i>	Manavgat, Akpınar Plateau (Kelsu Location), watersides, 1573 m, N 37°09'507" – E 31°20'312", 01.vi.2009, Çinbilgel 4893, (H).
Dioscoreaceae		
	<i>Dioscorea communis</i> (L.) Caddick & Wilkin	Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 680 m, N 37°10'071" – E 31°16'971", 06.vi.2011, Çinbilgel 7827, (G).
Ericaceae		
	<i>Arbutus andrachne</i> L.	Manavgat, Akpınar Plateau way, damaged forest, 1207 m, N 37°08'797" – E 31°18'139", 11.x.2009, Çinbilgel 5824; Manavgat, Burmahan Neighbourhood, Akpınar Plateau way, Yellibel Location, <i>Pinus brutia</i> forest, 896 m, N 37°09'186" – E 31°17'076", 26.iii.2011, Çinbilgel 7441; Manavgat, Çay Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°12'133" – E 31°16'452", 27.iii.2011, Çinbilgel 7473, (Ph).
	<i>Erica manipuliflora</i> Salisb.	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, under the <i>Pinus brutia</i> forest, 635 m, N 37°11'846" – E 31°16'238", 10.x.2009, Çinbilgel 5752, E. Medit., (Ph).
Euphorbiaceae		
	<i>Euphorbia condylocarpa</i> M.Bieb.	İbradı, Between Gavur Breach and Etülgenboğazi, <i>Pinus brutia</i> forest, 1186 m, N 37°09'074" – E 31°25'116", 20.iv.2008, Çinbilgel 2975; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1502 m, N 37°15'548" – E 31°24'365", 30.vi.2010, Çinbilgel 6659, (H).
	<i>Euphorbia falcata</i> L. subsp. <i>falcata</i> var. <i>galilaea</i> (Boiss.) Boiss.	İbradı, Gevenli Location, under the <i>Pinus nigra</i> forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3504; Manavgat, Kızılpınar Plateau way, Kavaklı Location, stony places, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4032, (Th).
	<i>Euphorbia helioscopia</i> L. subsp. <i>helioscopia</i>	Manavgat, Kızılpınar Plateau way, Putraklı Location, stony places, 734 m, N 37°12'249" – E 31°16'454", 27.iii.2011, Çinbilgel 7564, (Th).
	<i>Euphorbia herniariifolia</i> Willd. var. <i>herniariifolia</i>	Manavgat, Kovacık Mountain surroundings (Kocayokuş Location), rocky slopes, 1807 m, N 37°13'996" – E 31°20'274", 09.vi.2008, Çinbilgel 3656; Manavgat, Kızılpınar Plateau, rock, 1894 m, N 37°14'315" – E 31°20'502", 03.viii.2009, Çinbilgel 5621; İbradı, Melik Mountain (Orta Hill), rock, 2077 m, N 37°16'553" – E 31°23'091", 17.vi.2010, Çinbilgel 6415, (H).
	<i>Euphorbia kotschyana</i> Fenzl	İbradı, Gevenli Location, under the <i>Pinus brutia</i> forest, 1108 m, N 37°12'316" – E 31°24'613", 17.v.2008, Çinbilgel 3007; İbradı, Söbüova Location (around of Akpınar Location), under the forest, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3535; İbradı, Between Akpınar Location and Suçikan Location, stony places, 1091 m, N 37°13'075" – E 31°24'456", 18.vi.2008, Çinbilgel 4426, E. Medit. (mt.), (Ch).
	<i>Euphorbia peplus</i> L. var. <i>peplus</i>	İbradı, between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çinbilgel 2932; Manavgat, Burunçalı Location surroundings, 1581 m, N 37°08'893" – E 31°18'959", 02.vi.2009, Çinbilgel 5092; Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, Putraklı Location, stony places, 734 m, N 37°12'249" – E 31°16'454", 27.iii.2011, Çinbilgel 7568, (Th).
	<i>Euphorbia rigida</i> M.Bieb.	Manavgat, Akpınar Plateau way, <i>Pinus brutia</i> forest, ~1000 m, N 37°08'705" – E 31°19'909", 26.iii.2009, Çinbilgel 4543; Manavgat, Çay Neighbourhood, Burmahan Neighbourhood way, under the <i>Pinus brutia</i> forest, 642 m, N 37°11'726" – E 31°16'096", 26.iii.2011, Çinbilgel 7409; Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, Kabakulak Location, <i>Pinus brutia</i> forest clearings, 1014 m, N 37°12'843" – E 31°16'064", 27.iii.2011, Çinbilgel 7500, Medit., (Ch).
	<i>Euphorbia taurinensis</i> All.	İbradı, southeast slopes of Yaylacık Mountain, under the <i>Cedrus libani</i> forest, 1356 m, N 37°09'929" – E 31°25'029", 19.iv.2009, Çinbilgel 4592; İbradı, Akmuar Location, Juniper forest, 995 m, N 37°10'992" – E 31°25'278", 19.v.2011, Çinbilgel 7600; Manavgat, Çay Neighbourhood surroundings (Dikmen Hill), <i>Pinus brutia</i> forest, 816 m, N 37°12'492" – E 31°16'225", 07.vi.2011, Çinbilgel 7887, (Th).
	<i>Mercurialis annua</i> L.	Manavgat, Arıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4110, (Th).
Fabaceae		
	<i>Adenocarpus complicatus</i> (L.) Gay	Manavgat, between Burmahan Neighbourhood and Akpınar Plateau, under the <i>Pinus brutia</i> forest, 784 m, N 37°09'537" – E 31°17'054", 11.x.2009, Çinbilgel

		5815; Manavgat, Burmahan Neighbourhood, Erenli Location, streamsides, 749 m, N 37°09'726" – E 31°17'001", 06.vi.2011, Çinbilgel 7816, Medit., (Ph).
	<i>Anthyllis vulneraria</i> L. subsp. <i>praepropera</i> (Kerner) Bornm.	İbradı, Melik Mountain (Hasan Hill), stony slopes, 2136 m, N 37°16'753" – E 31°23'122", 18.vi.2008, Çinbilgel 4336; Manavgat, Kocatop Mountain, windy hills, 2174 m, N 37°12'467" – E 31°18'805", 10.vi.2009, Çinbilgel 5332; İbradı, Toka Plateau, stony places, 1538 m, N 37°14'038" – E 31°22'783", 09.v.2010, Çinbilgel 6082, Medit., (Th).
	* <i>Astragalus acmonotrichus</i> Fenzl	İbradı, around of Yirme Hill, forest clearings, 1674 m, N 37°14'298" – E 31°23'004", 18.v.2008, Çinbilgel 3210; İbradı, Toka Plateau Breach, stony slopes, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5990; İbradı, around of Toka Plateau Breach, <i>Cedrus libani</i> – <i>Pinus nigra</i> forest, 1679 m, N 37°13'808" – E 31°23'202", 21.v.2011, Çinbilgel 7672; İbradı, Toka Plateau way, <i>Pinus nigra</i> forest, 1662 m, N 37°13'925" – E 31°22'981", 08.v.2010, Çinbilgel 5998, End, E. Medit., (H), (Conf. Aytaç).
	* <i>Astragalus amoenus</i> Fenzl	Manavgat, Kepez Plateau, arid meadows, 1631 m, N 37°10'707" – E 31°21'189", 11.x.2009, Çinbilgel 5830; İbradı, Toka Plateau Breach, meadows, 1687 m, N 37°13'756" – E 31°23'188", 09.v.2010, Çinbilgel 6119; İbradı, Aşağıtoka Plateau, stony places, 1530 m, N 37°13'109" – E 31°22'633", 20.v.2011, Çinbilgel 7644, End., Ir. – Tur., (H).
	<i>Astragalus angustifolius</i> Lam. subsp. <i>angustifolius</i>	Manavgat, Kocatop Mountain (Koca Hill = Köygöründüğü Location), windy hills, 2150 m, N 37°12'346" – E 31°18'633", 10.vi.2009, Çinbilgel 5340; Manavgat, Kızılpınar Plateau surroundings, Nurboğazi Location, rock, 1913 m, N 37°14'166" – E 31°20'469", 04.viii.2009, Çinbilgel 5672; İbradı, Melik Mountain (Orta Hill), windy hills, 1999 m, N 37°16'161" – E 31°22'671", 17.vi.2010, Çinbilgel 6331, (Ch).
	<i>Astragalus angustifolius</i> Lam. subsp. <i>pungens</i> (Willd.) Hayek	İbradı, Toka Plateau (Göçebe Location), rocky places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çinbilgel 3126; İbradı, Melik Plateau, stony slopes, 1644 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4161; İbradı, Melik Mountain (southeastern hillsides of Hasan Hill), stony slopes, 1937 m, N 37°16'885" – E 31°23'249", 18.vi.2008, Çinbilgel 4295, (Ch).
	* <i>Astragalus cadmicus</i> Boiss.	Manavgat, Kocatop Mountain, windy hills, 2174 m, N 37°12'467" – E 31°18'805", 10.vi.2009, Çinbilgel 5322; Manavgat, Kocatop Mountain (Koca Hill = Köygöründüğü Location), windy hills, 2150 m, N 37°12'346" – E 31°18'633", 10.vi.2009, Çinbilgel 5338; Dermebeli Location (Arıtışı), rock, 1823 m, N 37°13'506" – E 31°20'339", 30.vi.2009, Çinbilgel 5540, End., (H).
	<i>Astragalus creticus</i> Lam.	İbradı, Aşağıtoka Plateau, thorny cushion formation, 1526 m, N 37°13'831" – E 31°22'356", 20.v.2011, Çinbilgel 7657, Ir. – Tur., (Ch), (Conf. Duran).
	<i>Astragalus depressus</i> L. var. <i>depressus</i>	Manavgat, Akpınar Plateau (Yumrutaş Location), <i>Cedrus libani</i> forest clearings, 1514 m, N 37°09'991" – E 31°21'166", 19.v.2009, Çinbilgel 4731; Manavgat, Akpınar Plateau, stony places, 1422 m, N 37°09'248" – E 31°20'517", 03.vi.2009, Çinbilgel 5151; İbradı, Toka Plateau Breach, stony places, 1691 m, N 37°13'748" – E 31°23'142", 09.v.2010, Çinbilgel 6115, (H).
	* <i>Astragalus heldreichii</i> Boiss.	Manavgat, Dermebeli Location, stony places, 1881 m, N 37°13'500" – E 31°20'284", 10.x.2009, Çinbilgel 5780, End., (H).
	* <i>Astragalus lycius</i> Boiss.	İbradı, Aşağıtoka Plateau, stony places, 1531 m, N 37°12'882" – E 31°22'682", 16.vi.2010, Çinbilgel 6241; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1487 m, N 37°15'663" – E 31°24'435", 30.vi.2010, Çinbilgel 6621; İbradı, around of Çeşgar Plateau Breach, stony places, 1681 m, N 37°17'294" – E 31°24'331", 01.vii.2010, Çinbilgel 6799, End., (H).
	* <i>Astragalus mesogitanus</i> Boiss.	Manavgat, Çağlan Breach – around of Susuzkır Hill, stony places, 1681 m, N 37°16'230" – E 31°21'349", 12.vi.2008, Çinbilgel 3971; Manavgat, Kovacık Mountain foothills (Gökçebel Location), dwarf bush-thorny cushion formation, 1687 m, N 37°14'038" – E 31°19'336", 09.vi.2009, Çinbilgel 5187; İbradı, Aşağıtoka Plateau, stony places, 1516 m, N 37°13'511" – E 31°22'567", 08.vi.2011, Çinbilgel 7949, End., Ir. – Tur., (H).
	<i>Astragalus odoratus</i> Lam.	İbradı, Aşağıtoka Plateau, stony places, 1519 m, N 37°13'935" – E 31°22'417", 16.vi.2010, Çinbilgel 6253; İbradı, Melik Plateau, openings – stony places, 1387 m, N 37°15'533" – E 31°24'528", 30.vi.2010, Çinbilgel 6724, (H).
	<i>Astragalus onobrychis</i> L.	Manavgat, Akpınar Plateau (Kelsu Location), dwarf bush formation, 1591 m, N 37°09'534" – E 31°20'305", 01.vi.2009, Çinbilgel 4842; Manavgat, Kocatop Mountain (Örencik Location), dwarf bush formation, 1861 m, N 37°11'829" – E 31°18'607", 11.vi.2009, Çinbilgel 5422, (H).
	<i>Astragalus plumosus</i> Willd.	Manavgat, Dermebeli Location, thorny cushion formation, 1775 m, N 37°12'927" – E 31°20'580", 05.viii.2009, Çinbilgel 5728; İbradı, Yukarıtoka Plateau, thorny cushion formation, 1615 m, N 37°15'329" – E 31°22'423", 18.vi.2010, Çinbilgel 6542; İbradı, Melik Plateau, under the <i>Pinus nigra</i> forest, 1470 m, N 37°15'607" – E 31°24'429", 30.vi.2010, Çinbilgel 6636, Ir. – Tur., (Ch).
	<i>Astragalus prusianus</i> Boiss.	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, Putraklı Location, 734 m, N 37°12'249" – E 31°16'454", 27.iii.2011, Çinbilgel 7562; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7722, E. Medit., (Ch).
	<i>Astragalus schizopterus</i> Boiss.	İbradı, between Gavur Breach and Etülgenboğazi, <i>Pinus brutia</i> forest, 1186 m, N 37°09'074" – E 31°25'116", 20.iv.2008, Çinbilgel 2980; Manavgat, Yelibelen Location, under the <i>Pinus brutia</i> forest, 896 m, N 37°09'186" – E

		31°17'076", 26.iii.2011, Çinbilgel 7443; İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1048 m, N 37°12'283" – E 31°24'732", 19.v.2011, Çinbilgel 7617, E. Medit., (Ch).
	<i>Astragalus sibthorpianus</i> Boiss.	Manavgat, Kovacık Mountain (around of Dermebeli Location), dwarf bush formation, 1867 m, N 37°13'751" – E 31°20'136", 29.vi.2009, Çinbilgel 5464; İbradı, Hillsides of Melik Mountain, dwarf bush formation, 1882 m, N 37°15'510" – E 31°23'322", 18.vi.2010, Çinbilgel 6488; İbradı, Aşağıtoka Plateau, wet meadows, 1514 m, N 37°13'772" – E 31°22'587", 08.vi.2011, Çinbilgel 7953, (H).
	<i>Bituminaria butiminosa</i> (L.) C.H.Stirt.	Manavgat, around of Burmahan Neighbourhood, streamsides, 693 m, N 37°11'151" – E 31°16'304", 06.vi.2011, Çinbilgel 7769, Medit., (H).
	<i>Calicotome villosa</i> (Poir.) Link	Manavgat, Çay Neighbourhood–Burmahan Neighbourhood way, <i>Pinus brutia</i> forest, 642 m, N 37°11'726" – E 31°16'096", 26.iii.2011, Çinbilgel 7404, Medit., (Ph).
	<i>Ceratonia siliqua</i> L.	Manavgat, Kızılpınar Plateau way, Göçek Location, <i>Pinus brutia</i> forest, 600 m, 13.vi.2008, Çinbilgel 4153, Medit., (Ph).
	<i>Cercis siliquastrum</i> L. subsp. <i>hebecarpa</i> (Bornm.) Yalt.	(Ph), (Observation).
	<i>Cicer incisum</i> (Willd.) K.Maly	Manavgat, Burunçalı Location (Aslanboğazi), screes, 1716 m, N 37°08'966" – E 31°19'501", 02.vi.2009, Çinbilgel 5018; Manavgat, Kocatop Mountain foothills, screes, 1814 m, N 37°12'479" – E 31°18'421", 11.vi.2009, Çinbilgel 5373, (H).
	* <i>Cicer isauricum</i> P.H.Davis	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3472; İbradı, Ekinlik Location surroundings, <i>Cedrus libani</i> forest (mixed with <i>Pinus nigra</i>), 1532 m, N 37°13'668" – E 31°23'510", 09.v.2010, Çinbilgel 6147; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7030, End., E. Medit., (Ch).
	* <i>Colutea melanocalyx</i> Boiss. & Heldr. subsp. <i>melanocalyx</i>	Manavgat, around of Burmahan Neighbourhood, <i>Pinus brutia</i> forest, 695 m, N 37°11'319" – E 31°16'296", 06.vi.2011, Çinbilgel 7783, End., E. Medit., (Ph).
	* <i>Dorycnium pentaphyllum</i> Scop. subsp. <i>haussknechtii</i> (Boiss.) Gams	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7736, End., Ir. – Tur., (H).
	<i>Genista acanthoclada</i> DC.	Manavgat, Çay Neighbourhood–Burmahan Neighbourhood way, <i>Pinus brutia</i> forest, 642 m, N 37°11'726" – E 31°16'096", 26.iii.2011, Çinbilgel 7401; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7723, E. Medit., (Ph).
	<i>Genista albida</i> Willd.	İbradı, Toka Plateau, moist stony places, 1522 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3347; İbradı, Toka Plateau, stony places, 1500 m, N 37°14'332" – E 31°22'569", 08.v.2010, Çinbilgel 6053; İbradı, Toka Plateau, meadows, 1514 m, N 37°13'924" – E 31°22'564", 09.v.2010, Çinbilgel 6093, (Ch).
	<i>Hippocrepis emerus</i> (L.) Lassen subsp. <i>emeroides</i> (Boiss. & Spruner) Lassen	İbradı, Suçikan Location way, stony – rocky places under the forest, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4413; İbradı, Cevizlidere Location, on the rock, 1132 m, N 37°13'382" – E 31°24'226", 02.vii.2010, Çinbilgel 7242, (Ph).
	<i>Hymenocarpus circinnatus</i> (L.) Savi	Manavgat, Çay Neighbourhood, Eşekbeli Location, <i>Pinus nigra</i> forest, 1612 m, N 37°14'867" – E 31°19'864", 07.vi.2011, Çinbilgel 7892, Medit., (Th).
	<i>Lathyrus aureus</i> (Steven) D.Brandza	İbradı, between Gavur Breach and Etülgenboğazi, <i>Pinus brutia</i> forest, 1186 m, N 37°09'074" – E 31°25'116", 20.iv.2008, Çinbilgel 2972; İbradı, Boğazyurt Location, <i>Pinus brutia</i> forest, 970 m, N 37°11'161" – E 31°25'266", 21.v.2011, Çinbilgel 7700; Manavgat, around of Burmahan Neighbourhood, <i>Pinus brutia</i> forest, 695 m, N 37°11'319" – E 31°16'296", 06.vi.2011, Çinbilgel 7785, Medit., (Th).
	<i>Lathyrus digitatus</i> (M.Bieb.) Fiori	İbradı, Toka Plateau way, under the <i>Cedrus libani</i> forest, 1460 m, N 37°13'078" – E 31°23'776", 19.v.2008, Çinbilgel 3492; İbradı, Ekinlik Location surroundings, under the <i>Cedrus libani</i> forest, 1691 m, N 37°13'304" – E 31°23'423", 09.v.2010, Çinbilgel 6138; İbradı, Akmuar Location, Juniper Forest clearings, 1036 m, N 37°11'773" – E 31°24'864", 21.v.2011, Çinbilgel 7680, E. Medit., (H).
	<i>Lathyrus hirsutus</i> L.	Manavgat, Çay Neighbourhood surroundings, open places, 691 m, N 37°12'049" – E 31°16'444", 07.vi.2011, Çinbilgel 7878; Manavgat, north of Çay Neighbourhood (Dikmen Hill), <i>Pinus brutia</i> forest, 816 m, N 37°12'492" – E 31°16'225", 07.vi.2011, Çinbilgel 7890, Medit., (Th).
	<i>Lathyrus inconspicuus</i> L. var. <i>inconspicuus</i>	İbradı, Akmuar Location, Juniper Forest, 995 m, N 37°10'992" – E 31°25'278", 19.v.2011, Çinbilgel 7597, (Th).
	<i>Lathyrus laxiflorus</i> (Desf.) O.Kuntze subsp. <i>laxiflorus</i>	İbradı, between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çinbilgel 2938; Manavgat, Kovacık Mountain foothills (Elmalı Location), <i>Pinus nigra</i> forest, 1615 m, N 37°13'748" – E 31°18'967", 11.vi.2009, Çinbilgel 5412; İbradı, Suçikan Location, streamsides, 1220 m, N 37°14'429" – E 31°24'260", 02.vii.2010, Çinbilgel 7110, (H).
	<i>Lathyrus nissolia</i> L.	İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1487 m, N 37°15'663" – E 31°24'435", 30.vi.2010, Çinbilgel 6633, (Th).

	<i>Lathyrus setifolius</i> L.	İbradı, Gavur Breach – Etülgen boğazi, <i>Pinus brutia</i> forest, 977 m, N 37°09'100" – E 31°25'485", 20.iv.2008, Çinbilgel 2987; Manavgat, Akpınar Plateau– hillsides of Kaldırım Mountain, stony places, 1541 m, N 37°09'456" – E 31°20'380", 01.vi.2009, Çinbilgel 4822; İbradı, Toka Plateau way, under the <i>Pinus nigra</i> forest, 1600 m, N 37°13'955" – E 31°22'921", 08.vi.2011, Çinbilgel 7960, E. Medit., (Th).
	<i>Lathyrus sphaericus</i> Retz.	Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4826, Medit., (Th).
	<i>Lathyrus stenophyllus</i> Boiss. & Heldr.	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7710; Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 749 m, N 37°09'726" – E 31°17'001", 06.vi.2011, Çinbilgel 7814; Manavgat, around of Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest, 702 m, N 37°09'905" – E 31°16'998", 06.vi.2011, Çinbilgel 7833, E. Medit., (Th).
	<i>Lens culinaris</i> Medik. subsp. <i>orientalis</i> (Boiss.) Ponert	Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest clearings, 1430 m, N 37°09'494" – E 31°20'838", 19.v.2009, Çinbilgel 4813; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1487 m, N 37°15'592" – E 31°24'391", 30.vi.2010, Çinbilgel 6646; around of Burmahan Neighbourhood (Öküzzaşı Location), <i>Pinus brutia</i> forest, 683 m, N 37°10'293" – E 31°16'897", 06.vi.2011, Çinbilgel 7810, (Th).
	<i>Lens ervoides</i> (Brign.) Grande	İbradı, between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çinbilgel 2934; İbradı, between Gavur Breach and Etülgenboğazi, <i>Pinus brutia</i> forest, 1186 m, N 37°09'074" – E 31°25'116", 20.iv.2008, Çinbilgel 2981, Medit., (Th).
	<i>Lotus corniculatus</i> L. var. <i>alpinus</i> Ser.	Manavgat, Kızılpinar Plateau, dwarf bush formation, 1747 m, N 37°14'087" – E 31°19'983", 29.vi.2009, Çinbilgel 5509; İbradı, Aşağıtoka Plateau, thorny cushion formation, 1526 m, N 37°12'919" – E 31°22'639", 16.vi.2010, Çinbilgel 6183; İbradı, Melik Plateau surroundings, streamsides, 1346 m, N 37°15'389" – E 31°24'545", 02.vii.2010, Çinbilgel 7017, (H).
	<i>Lotus corniculatus</i> L. var. <i>corniculatus</i>	Manavgat, Kovacık Mountain foothills (Gökçebel Location), thorny cushion formation, 1695 m, N 37°13'984" – E 31°19'318", 11.vi.2009, Çinbilgel 5444; İbradı, Aşağıtoka Plateau, thorny cushion formation, 1526 m, N 37°12'919" – E 31°22'639", 16.vi.2010, Çinbilgel 6183; İbradı, Melik Plateau surroundings, streamsides, 1346 m, N 37°15'389" – E 31°24'545", 02.vii.2010, Çinbilgel 7017, (H).
	<i>Lotus corniculatus</i> L. var. <i>tenuifolius</i> L.	Manavgat, Kepez Plateau (İncayırı Location), damp places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5851; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çinbilgel 6712; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest, 1347 m, N 37°15'332" – E 31°24'499", 02.vii.2010, Çinbilgel 7009, (H).
	<i>Medicago brachycarpa</i> M.Bieb.	Manavgat, southern of Kaldırım Mountain, slopes, 1943 m, N 37°09'635" – E 31°20'292", 01.vi.2009, Çinbilgel 4946; Manavgat, Kızılpinar Plateau, mixed <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1718 m, N 37°14'103" – E 31°19'803", 09.vi.2009, Çinbilgel 5257; İbradı, Yukarıtoka Plateau, dwarf bush formation, 1635 m, N 37°15'828" – E 31°22'221", 17.vi.2010, Çinbilgel 6276, Ir. – Tur., (Th).
	<i>Medicago coronata</i> (L.) Bartal.	Manavgat, Kaldırım Mountain (Hıncırak Location), mixed <i>Cedrus libani</i> forest, 1475 m, N 37°09'368" – E 31°20'410", 01.vi.2009, Çinbilgel 4902; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7715, Medit., (Th).
	<i>Medicago disciformis</i> DC.	İbradı, Söbüova Location (around of Akpınar Location), stony places, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3549; İbradı, Melik Plateau, forest clearings, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6558; İbradı, Eynif Plain, meadows, 941 m, N 37°08'989" – E 31°26'561", 19.v.2011, Çinbilgel 7580, Medit., (Th).
	<i>Medicago doliata</i> Carmign. var. <i>muricata</i> (Benth.) Heyn	Manavgat, north of Çay Neighbourhood (Dikmen Hill), <i>Pinus brutia</i> forest, 816 m, N 37°12'492" – E 31°16'225", 07.vi.2011, Çinbilgel 7898, (Th).
	<i>Medicago lupulina</i> L.	Manavgat, Akpınar Plateau (Kelsu Location), 1591 m, N 37°09'534" – E 31°20'305", 01.vi.2009, Çinbilgel 4845; Manavgat, Kepez Plateau (İncayırı Location), stony places, 1624 m, N 37°11'525" – E 31°20'883", 11.x.2009, Çinbilgel 5832; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1487 m, N 37°15'663" – E 31°24'435", 30.vi.2010, Çinbilgel 6627, (H).
	<i>Medicago minima</i> (L.) Bartal. var. <i>minima</i>	Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4105; İbradı, Melik Plateau, forest clearings, 1445 m, N 37°15'662" – E 31°24'501", 30.vi.2010, Çinbilgel 6575; İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1223 m, N 37°14'468" – E 31°24'259", 02.vii.2010, Çinbilgel 7052, (Th).
	<i>Medicago orbicularis</i> (L.) Bartal.	Manavgat, Düğnük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4082; Manavgat, Çay Neighbourhood, opennings – roadsides, 690 m, N 37°12'122" – E 31°16'461", 07.vi.2011, Çinbilgel 7932; İbradı, Gevenli Location, <i>Pinus brutia</i> forest clearings, 1071 m, N 37°12'300" – E 31°24'793", 03.vii.2011, Çinbilgel 8007, (Th).
	<i>Medicago radiata</i> L.	İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1223 m, N 37°14'468" – E 31°24'259", 02.vii.2010, Çinbilgel 7051; Manavgat, around of Burmahan Neighbourhood (Öküzzaşı Location), <i>Pinus brutia</i> forest, 683 m, N 37°10'293" – E 31°16'897", 06.vi.2011, Çinbilgel 7802, Ir. – Tur., (Th).

	<i>Medicago rigidula</i> (L.) All. var. <i>rigidula</i>	İbradı, Gavur Breach way – Etülgen boğazı, <i>Pinus brutia</i> forest, 977 m, N 37°09'100" – E 31°25'485", 20.iv.2008, Çinbilgel 2990, (Th).
	<i>Medicago sativa</i> L. subsp. <i>sativa</i>	İbradı, Çeşgar Plateau, stony slopes, 1659 m, N 37°17'402" – E 31°24'250", 17.vi.2008, Çinbilgel 4242; Manavgat, Derme Plateau, dwarf bush formation, 1805 m, N 37°13'393" – E 31°20'371", 05.viii.2009, Çinbilgel 5708; İbradı, Aşağıtoka Plateau, stony places, 1523 m, N 37°12'924" – E 31°22'540", 16.vi.2010, Çinbilgel 6197, (H).
	<i>Melilotus indicus</i> (L.) All.	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7721, (Th).
	<i>Onobrychis caput – galli</i> (L.) Lam.	Manavgat, Between Çay and Burmahan Neighbourhoods, <i>Pinus brutia</i> forest, 700 m, N 37°11'519" – E 31°16'218", 06.vi.2011, Çinbilgel 7773, Medit., (Th).
	<i>Onobrychis montana</i> DC. subsp. <i>cadmea</i> (Boiss.) P.W.Ball	Manavgat, Derme Plateau (towards Dermebeli Location), stony places, 1827 m, N 37°13'386" – E 31°20'237", 10.vi.2008, Çinbilgel 3796; Manavgat, Kızılpınar Plateau, stony places, 1750 m, N 37°14'071" – E 31°19'981", 19.vii.2008, Çinbilgel 4456; Manavgat, Dermebeli Location, dwarf bush formation, 1935 m, N 37°13'656" – E 31°20'290", 29.vi.2009, Çinbilgel 5457, (H).
	<i>Onobrychis oxyodonta</i> Boiss. var. <i>armena</i>	Manavgat, eastern slopes of Sürüçikan Mountain, dwarf bush formation, 2051 m, N 37°12'401" – E 31°19'742", 30.vi.2009, Çinbilgel 5538; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 2085 m, N 37°16'595" – E 31°22'998", 17.vi.2010, Çinbilgel 6429; İbradı, Akmuar Location, <i>Juniperus excelsa</i> forest, 1044 m, N 37°11'404" – E 31°24'996", 03.vii.2011, Çinbilgel 8037, (H).
	<i>Onobrychis oxyodonta</i> Boiss. var. <i>oxyodonta</i>	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 622 m, N 37°11'707" – E 31°16'064", 06.vi.2011, Çinbilgel 7743; Manavgat, Çay Neighbourhood, Kavaklı Location, <i>Juniperus excelsa</i> forest, 1323 m, N 37°13'974" – E 31°18'405", 07.vi.2011, Çinbilgel 7924, (H).
	<i>Ononis adenotricha</i> Boiss.	Manavgat, Kovacık Mountain foothills (Gökçebel Location), thorny cushion formation, 1695 m, N 37°13'984" – E 31°19'318", 11.vi.2009, Çinbilgel 5446; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 2083 m, N 37°16'553" – E 31°22'889", 17.vi.2010, Çinbilgel 6439; İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1241 m, N 37°14'491" – E 31°24'214", 02.vii.2010, Çinbilgel 7094, E. Medit., (Ch).
	<i>Ononis spinosa</i> L. subsp. <i>leiosperma</i> (Boiss.) Sirj.	Manavgat, Derme Plateau (towards Dermebeli Location), stony hillsides, 1827 m, N 37°13'386" – E 31°20'237", 10.vi.2008, Çinbilgel 3793; Manavgat, Kızılpınar Plateau way (Kavaklı Location), stony places, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4043; Manavgat, Kepez Plateau (İnçayırı Location), damp places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5859, (H).
	<i>Ornithopus compressus</i> L.	Manavgat, around of Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest, 702 m, N 37°09'905" – E 31°16'998", 06.vi.2011, Çinbilgel 7836, Medit., (Th).
	<i>Pisum sativum</i> L. subsp. <i>elatius</i> (M.Bieb.) Aschers. & Graebn. var. <i>elatius</i>	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3462; İbradı, Ekinlik Location surroundings, <i>Cedrus libani</i> forest (mixed with <i>Pinus nigra</i>), 1532 m, N 37°13'668" – E 31°23'510", 09.v.2010, Çinbilgel 6139, Medit., (Th).
	* <i>Securigera grandiflora</i> Lassen	Manavgat, north of Çay Neighbourhood (Dikmen Hill), <i>Pinus brutia</i> forest, 816 m, N 37°12'492" – E 31°16'225", 07.vi.2011, Çinbilgel 7891, End., Medit., (Th).
	<i>Securigera parviflora</i> (Desv.) Lassen	İbradı, between Gavur Breach and Etülgenboğazı, <i>Pinus brutia</i> forest, 1186 m, N 37°09'074" – E 31°25'116", 20.iv.2008, Çinbilgel 2985; İbradı, Söbüova Location (around of Akpınar Location), stony places, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3546; İbradı, Akmuar Location, Juniper Forest clearings, 1036 m, N 37°11'773" – E 31°24'864", 21.v.2011, Çinbilgel 7678, E. Medit., (Th).
	<i>Securigera securidaca</i> (L.) Degen & Dorfl.	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest clearings, 655 m, N 37°11'683" – E 31°16'103", 06.vi.2011, Çinbilgel 7751, (Th).
	<i>Securigera varia</i> (L.) Lassen	İbradı, Tahtalı Location, stony places, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4383; İbradı, Melik Plateau, stony places, 1387 m, N 37°15'533" – E 31°24'528", 30.vi.2010, Çinbilgel 6735; İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1241 m, N 37°14'491" – E 31°24'214", 02.vii.2010, Çinbilgel 7093, E. Medit., (H).
	<i>Spartium junceum</i> L.	Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest clearings, 1028 m, N 37°12'886" – E 31°17'080", 10.x.2009, Çinbilgel 5756; Manavgat, Burmahan Neighbourhood, Akpınar Plateau way, Yellibel Location, <i>Pinus brutia</i> forest, 896 m, N 37°09'186" – E 31°17'076", 26.iii.2011, Çinbilgel 7442; Manavgat, Çay Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°12'133" – E 31°16'452", 27.iii.2011, Çinbilgel 7481, Medit., (Ph).
	<i>Trifolium arvense</i> L. var. <i>arvense</i>	Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4103; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest, 1344 m, N 37°15'376" – E 31°24'546", 02.vii.2010, Çinbilgel 7023; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7173, (Th).

	<i>Trifolium campestre</i> Schreb. subsp. <i>campestre</i> var. <i>campestre</i>	Manavgat, Arğıça Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4126; İbradı, around of Melik Plateau, forest clearings – stony places, 1406 m, N 37°15'491" – E 31°24'478", 30.vi.2010, Çinbilgel 6697; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 622 m, N 37°11'707" – E 31°16'064", 06.vi.2011, Çinbilgel 7748, (Th).
	* <i>Trifolium caudatum</i> Boiss.	Manavgat, Kovacık Mountain (around of Dermebeli Location), dwarf bush formation, 1867 m, N 37°13'751" – E 31°20'136", 29.vi.2009, Çinbilgel 5460; İbradı, Çeşgar Plateau Breach, stony places, 1648 m, N 37°17'056" – E 31°24'397", 01.vii.2010, Çinbilgel 6798, End., (H).
	<i>Trifolium cherleri</i> L.	İbradı, Toka Plateau Breach, meadows, 1676 m, N 37°13'774" – E 31°23'168", 21.v.2011, Çinbilgel 7663; Manavgat, around of Burmahan Neighbourhood (Öküztası Location), <i>Pinus brutia</i> forest, 683 m, N 37°10'293" – E 31°16'897", 06.vi.2011, Çinbilgel 7797; Manavgat, around of Burmahan Neighbourhood (Öküztası Location), <i>Pinus brutia</i> forest, 683 m, N 37°10'293" – E 31°16'897", 06.vi.2011, Çinbilgel 7807, Medit., (Th).
	<i>Trifolium fragiferum</i> L. var. <i>fragiferum</i>	Manavgat, Akpınar Plateau (Hıncırak Location), rubble rocks, 1468 m, N 37°09'337" – E 31°20'412", 01.vi.2009, Çinbilgel 4863; Manavgat, Kovacık Mountain foothills, thorny cushion formation, 1712 m, N 37°13'900" – E 31°19'264", 09.vi.2009, Çinbilgel 5215, (H).
	<i>Trifolium grandiflorum</i> Schreb.	İbradı, Gevenli Location, <i>Pinus nigra</i> forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3513; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest clearings, 965 m, N 37°10'640" – E 31°25'709", 19.v.2011, Çinbilgel 7591; İbradı, Akmuar Location, Juniper Forest clearings, 1036 m, N 37°11'773" – E 31°24'864", 21.v.2011, Çinbilgel 7684, E. Medit., (Th).
	<i>Trifolium hirtum</i> All.	Manavgat, Arğıça Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4127; Manavgat, around of Burmahan Neighbourhood (Payır Location), <i>Pinus brutia</i> forest, 803 m, N 37°09'640" – E 31°17'067", 06.vi.2011, Çinbilgel 7851, Medit., (Th).
	<i>Trifolium lappaceum</i> L.	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7719; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7727, Medit., (Th).
	<i>Trifolium lucanicum</i> Gasp	İbradı, Gevenli Location, <i>Pinus nigra</i> forest 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3501; İbradı, Melik Plateau, dwarf bush formation, 1728 m, N 37°16'359" – E 31°24'498", 01.vii.2010, Çinbilgel 6933; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7167, Medit., (Th).
	<i>Trifolium nervulosum</i> Boiss. & Heldr.	İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> clearings, 965 m, N 37°10'640" – E 31°25'709", 19.v.2011, Çinbilgel 7595; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 990 m, N 37°10'778" – E 31°25'386", 21.v.2011, Çinbilgel 7696, E. Medit., (Th).
	<i>Trifolium nigrescens</i> Viv. subsp. <i>petrisavii</i>	Manavgat, Kovacık Mountain (Gökçebel Location), thorny cushion formation, 1709 m, N 37°14'032" – E 31°19'421", 09.vi.2009, Çinbilgel 5225; İbradı, Melik Plateau, forest clearings, 1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6960; Manavgat, around of Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest, 702 m, N 37°09'905" – E 31°16'998", 06.vi.2011, Çinbilgel 7837, (Th).
	<i>Trifolium physodes</i> Steven ex M.Bieb. var. <i>physodes</i>	Manavgat, Kızılıpınar Plateau (Kocaboğaz Location), arid meadows, 1790 m, N 37°13'949" – E 31°20'094", 29.vi.2009, Çinbilgel 5480; İbradı, Çeşgar Plateau surroundings (Çeşgar Plateau Breach), stony places, 1682 m, N 37°17'288" – E 31°24'324", 01.vii.2010 Çinbilgel 6906; İbradı, Akmuar Location, Juniper Forest, 1047 m, N 37°11'401" – E 31°25'006", 21.v.2011, Çinbilgel 7687, Medit., (H).
	<i>Trifolium pratense</i> L. var. <i>pratense</i>	Manavgat, Kızılıpınar Plateau surroundings, stony rocky places, 1798 m, N 37°13'991" – E 31°20'198", 09.vi.2008, Çinbilgel 3650; İbradı, between Melik Mountain (Hasan Hill) and Çeşgar Plateau Location, stony slopes, 1657 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4259; Manavgat, around of Burmahan Neighbourhood (Öküztası Location), <i>Pinus brutia</i> forest, 683 m, N 37°10'293" – E 31°16'897", 06.vi.2011, Çinbilgel 7803, (H).
	<i>Trifolium pamphylicum</i> Boiss. & Heldr.	Manavgat, Kızılıpınar Plateau way, <i>Pinus brutia</i> forest, 886 m, N 37°13'095" – E 31°16'406", 25.v.2008, Çinbilgel 3603a, E. Medit., (Th).
	<i>Trifolium repens</i> L. var. <i>repens</i>	Manavgat, Çağlan Breach, stony places, 1641 m, N 37°15'823" – E 31°20'867", 12.vi.2008, Çinbilgel 3996; İbradı, Melik Plateau, openings – stony places, 1387 m, N 37°15'533" – E 31°24'528", 30.vi.2010, Çinbilgel 6726; İbradı, Aşağıtoka Plateau, thorny cushion formation, 1526 m, N 37°13'831" – E 31°22'356", 20.v.2011, Çinbilgel 7655, (H).
	<i>Trifolium resupinatum</i> L. var. <i>resupinatum</i>	İbradı, Cevzlidere Location (around of Suçikan Location), streamsides, 1140 m, N 37°13'377" – E 31°24'211", 05.x.2010, Çinbilgel 7275; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest clearings, 965 m, N 37°10'640" – E 31°25'709", 19.v.2011, Çinbilgel 7590; Manavgat, Çay Neighbourhood, Eşekbeli Location, <i>Pinus nigra</i> forest, 1612 m, N 37°14'867" – E 31°19'864", 07.vi.2011, Çinbilgel 7908, (Th).
	<i>Trifolium scabrum</i> L.	Manavgat, Düşük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4073; İbradı, Melik Plateau, forest clearings,

		1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6965; Manavgat, around of Burmahan Neighbourhood (Öküzaşı Location), <i>Pinus brutia</i> forest, 683 m, N 37°10'293" – E 31°16'897", 06.vi.2011, Çinbilgel 7808, Medit., (Th).
	<i>Trifolium striatum</i> L.	İbradı, Melik Plateau, meadows, 1446 m, N 37°15'954" – E 31°24'591", 01.vii.2010, Çinbilgel 6758, (Th).
	<i>Trifolium subterraneum</i> L.	Manavgat, Genger Location, <i>Pinus brutia</i> forest, 1150 m, N 37°09'242" – E 31°17'835", 06.vi.2011, Çinbilgel 7865, (Th).
	<i>Trifolium tomentosum</i> L. var. <i>tomentosum</i>	İbradı, Eynif Plain, meadows, 941 m, N 37°08'989" – E 31°26'561", 19.v.2011, Çinbilgel 7578, (Th).
	<i>Trigonella cariensis</i> Boiss.	Manavgat, Akpınar Plateau (Yumrutaş Location), mixed <i>Cedrus libani</i> forest, 1520 m, N 37°09'840" – E 31°21'028", 19.v.2009, Çinbilgel 4787, E. Medit., (Th).
	<i>Trigonella foenum – graecum</i> L.	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3486, (Th).
	* <i>Trigonella lycica</i> Hub. – Mor.	Manavgat, Kaldırım Mountain (Hıncırak Location), mixed <i>Cedrus libani</i> forest, 1475 m, N 37°09'368" – E 31°20'410", 01.vi.2009, Çinbilgel 4894; İbradı, Aşağıtoka Plateau, stony places, 1546 m, N 37°13'972" – E 31°22'369", 16.vi.2010, Çinbilgel 6265; İbradı, Suçikan Location, <i>Pinus nigra</i> forest clearings, 1186 m, N 37°14'441" – E 31°24'305", 03.vii.2011, Çinbilgel 8026, End., E. Medit., (Th).
	<i>Trigonella monspeliaca</i> L.	Manavgat, Burmahan Neighbourhood – Kovacık Mountain way, rocky places, 1214 m, N 37°13'606" – E 31°17'996", 25.v.2008, Çinbilgel 3591/a; Manavgat, Düğnük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4089, Medit., (Th).
	<i>Trigonella spicata</i> Sibth. & Sm.	Manavgat, Burmahan Neighbourhood – Kovacık Mountain way, <i>Pinus brutia</i> forest, 886 m, N 37°13'095" – E 31°16'406", 25.v.2008, Çinbilgel 3600/a; İbradı, Eynif Plain, meadows, 941 m, N 37°08'989" – E 31°26'561", 19.v.2011, Çinbilgel 7579; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7713, E. Medit., (Th).
	<i>Vicia cassia</i> Boiss.	Manavgat, Between Çay and Burmahan Neighbourhoods, streamsides, 706 m, N 37°11'423" – E 31°16'306", 06.vi.2011, Çinbilgel 7757; Manavgat, around of Burmahan Neighbourhood (Değirmenderesi Location), damp places, 685 m, N 37°10'723" – E 31°16'850", 06.vi.2011, Çinbilgel 7788, E. Medit., (Th).
	<i>Vicia cracca</i> L. subsp. <i>stenophylla</i> Vel.	Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4833; İbradı, Ekinlik Location surroundings, <i>Cedrus libani</i> forest (mixed with <i>Pinus nigra</i>), 1532 m, N 37°13'668" – E 31°23'510", 09.v.2010, Çinbilgel 6148; Manavgat, Genger Location, <i>Pinus brutia</i> forest, 1150 m, N 37°09'242" – E 31°17'835", 06.vi.2011, Çinbilgel 7866, (H).
	<i>Vicia grandiflora</i> Scop. var. <i>grandiflora</i>	Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 773 m, N 37°09'851" – E 31°16'989", 06.vi.2011, Çinbilgel 7847, (Th).
	<i>Vicia hirsuta</i> (L.) Gray	İbradı, between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çinbilgel 2931; İbradı, Gavur Breach way – Etilgen boğazı, <i>Pinus brutia</i> forest, 977 m, N 37°09'100" – E 31°25'485", 20.iv.2008, Çinbilgel 2999, (Th).
	<i>Vicia lunata</i> Boiss. var. <i>lunata</i>	Manavgat, Yelibelen Location, <i>Pinus brutia</i> forest, 1048 m, N 37°09'140" – E 31°17'606", 06.vi.2011, Çinbilgel 7861, (Th).
	<i>Vicia lutea</i> L. var. <i>hirta</i> (Balbis) Loisel	İbradı, Akmuar Location, rock, 1024 m, N 37°11'720" – E 31°24'844", 19.v.2011, Çinbilgel 7610, (Th).
	<i>Vicia sativa</i> L. subsp. <i>nigra</i> (L.) Ehrh. var. <i>nigra</i>	İbradı, around of Yirme Hill, forest clearings, 1783 m, N 37°14'667" – E 31°23'097", 18.v.2008, Çinbilgel 3178; İbradı, Melik Plateau foothills, forest clearings – stony places, 1406 m, N 37°15'491" – E 31°24'478", 30.vi.2010, Çinbilgel 6700; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest clearings, 965 m, N 37°10'640" – E 31°25'709", 19.v.2011, Çinbilgel 7589, (Th).
	<i>Vicia villosa</i> Roth subsp. <i>eriocarpa</i> (Hausskn.) P.W.Ball	Manavgat, Burmahan Neighbourhood, Kovacık Mountain way, <i>Pinus brutia</i> forest, 886 m, N37° 13' 095" – E031° 16' 406", 25.v.2008, Çinbilgel 3601/a, (H).
Fagaceae		
	<i>Quercus cerris</i> L.	İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3517; Manavgat, Akpınar Plateau way, roadsides, 1300 m, N 37°11'110" – E 31°21'021", 03.vi.2009, Çinbilgel 5185; İbradı, around of Gevenli Location, <i>Pinus brutia</i> forest, 1147 m, N 37°12'229" – E 31°24'506", 02.vii.2010, Çinbilgel 7238, Medit., (Ph).
	<i>Quercus coccifera</i> L.	İbradı, Akmuar Location, Juniper forest, 964 m, N 37°11'201" – E 31°25'141", 18.vi.2008, Çinbilgel 4445; Manavgat, Akpınar Plateau way, damaged forest, 1306 m, N 37°08'484" – E 31°19'182", 11.x.2009, Çinbilgel 5822; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 974 m, N 37°10'600" – E 31°25'718", 26.ix.2010, Çinbilgel 7204, Medit., (Ph).
	<i>Quercus infectoria</i> Oliv. subsp. <i>veneris</i> (A.Kern.) Meikle	Manavgat, Akpınar Plateau way, <i>Pinus brutia</i> forest clearings, 1207 m, N 37°08'797" – E 31°18'139", 11.x.2009, Çinbilgel 5823, (Ph).
	<i>Quercus trojana</i> Webb subsp. <i>trojana</i>	Manavgat, Burunçalı Location, mixed <i>Cedrus libani</i> forest, 1567 m, N 37°08'813" – E 31°19'104", 02.vi.2009, Çinbilgel 4996; İbradı, Melik Plateau,

		<i>Pinus nigra</i> forest, 1511 m, N 37°15'476" – E 31°24'352", 30.vi.2010, Çinbilgel 6664; İbradı, Çeşgar Plateau, rocky places, 1625 m, N 37°17'554" – E 31°24'094", 01.vii.2010, Çinbilgel 6902, E. Medit., (Ph).
Gentianaceae		
	<i>Centaurium erythraea</i> Rafn. subsp. <i>turcicum</i> (Velen.) Melderis	İbradı, Eynif Plain, open places, 947 m, N 37°07'654" – E 31°27'562", 02.vii.2011, Çinbilgel 7966, (H).
	<i>Centaurium pulchellum</i> (Sw.) Druce	İbradı, Eynif Plain, open places, 947 m, N 37°07'654" – E 31°27'562", 02.vii.2011, Çinbilgel 7966/a, (Th).
Geraniaceae		
	<i>Erodium ciconium</i> (L.) L'Hér.	İbradı, Suçikan Location, streamsides, 1259 m, N 37°14'503" – E 31°24'173", 02.vii.2010, Çinbilgel 7066, (Th).
	<i>Erodium cicutarium</i> (L.) L'Hér. subsp. <i>cicutarium</i>	İbradı, around of Yirme Hill, forest clearings, 1783 m, N 37°14'667" – E 31°23'097", 18.v.2008, Çinbilgel 3186; Manavgat, Yalıncaadağ, dwarf bush-thorny cushion formation, 2059 m, N 37°11'417" – E 31°18'791", 11.vi.2009, Çinbilgel 5378; İbradı, Yukartoka Plateau, dwarf bush formation, 1635 m, N 37°15'828" – E 31°22'221", 17.vi.2010, Çinbilgel 6272, (Th).
	<i>Geranium columbinum</i> L.	Manavgat, around of Burmahan Neighbourhood, roadsides, 700 m, N 37°10'623" – E 31°16'847", 06.vi.2011, Çinbilgel 7793; Manavgat, Çay Neighbourhood, openings – roadsides, 690 m, N 37°12'122" – E 31°16'461", 07.vi.2011, Çinbilgel 7938, (Th).
	* <i>Geranium glaberrimum</i> Boiss. & Heldr.	Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest clearings, 1430 m, N 37°09'494" – E 31°20'838", 19.v.2009, Çinbilgel 4816; Manavgat, Kocatop Mountain foothills (Ocakyeri Location), rock, 1782 m, N 37°12'226" – E 31°18'317", 11.vi.2009, Çinbilgel 5429; İbradı, Çeşgar Plateau, rock, 1625 m, N 37°17'470" – E 31°23'695", 01.vii.2010, Çinbilgel 6838, End., E. Medit., (mt.), (Ch).
	<i>Geranium lucidum</i> L.	Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4123; İbradı, Çeşgar Plateau, rock, 1592 m, N 37°17'611" – E 31°23'752", 01.vii.2010, Çinbilgel 6864; Manavgat, Burmahan Neighbourhood, Erenli Location, streamsides, 753 m, N 37°09'844" – E 31°16'991", 26.iii.2011, Çinbilgel 7436, (Th).
	<i>Geranium molle</i> L.	Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4659; İbradı, Çukuryurt Plateau surroundings, under the forest, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6172; Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, Kocaarmut Location, <i>Pinus brutia</i> forest clearings – stony places, 709 m, N 37°12'174" – E 31°16'360", 27.iii.2011, Çinbilgel 7553, (Th).
	<i>Geranium purpureum</i> Vill.	Manavgat, Akpınar Plateau (Hıncırak Location), rubble rocks, 1468 m, N 37°09'337" – E 31°20'412", 01.vi.2009, Çinbilgel 4873; İbradı, Eynif Plain, Yarbaşı Location, on the rock, 943 m, N 37°08'421" – E 31°27'636", 28.iii.2010, Çinbilgel 5966; İbradı, Çeşgar Plateau, rock, 1625 m, N 37°17'470" – E 31°23'695", 01.vii.2010, Çinbilgel 6839, (Th).
	<i>Geranium pusillum</i> Burm.f.	İbradı, Çeşgar Plateau, dwarf bush formation, 1600 m, N 37°17'521" – E 31°23'740", 01.vii.2010, Çinbilgel 6830, (Th).
	<i>Geranium pyrenaicum</i> Burm.f.	Manavgat, Dermebeli Location (Arıtışı), on the rock and stony places, 1874 m, N 37°13'519" – E 31°20'317", 10.vi.2008, Çinbilgel 3819; Manavgat, Kızılpınar Plateau, under the <i>Cedrus libani</i> forest, 1717 m, N 37°14'110" – E 31°19'899", 09.vi.2009, Çinbilgel 5235; İbradı, Melik Plateau, forest clearings, 1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6961, (Ch).
	<i>Geranium rotundifolium</i> L.	İbradı, Suçikan Location way, stony places, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2784; Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4837; İbradı, Melik Plateau, stony places, 1491 m, N 37°15'544" – E 31°24'739", 02.vii.2010, Çinbilgel 6992, (Th).
	<i>Geranium tuberosum</i> L.	İbradı, Yukartoka Plateau (Between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3060; Manavgat, Akpınar Plateau, mixed <i>Cedrus libani</i> forest, 1485 m, N 37°09'577" – E 31°21'003", 03.v.2009, Çinbilgel 4671; Manavgat, Kaldırım Mountain, stony places, 1820 m, N 37°09'912" – E 31°20'423", 01.vi.2009, Çinbilgel 4861, Ir. – Tur., (G).
	<i>Pelargonium endlicherianum</i> Fenzl	Manavgat, Akpınar Plateau (Sumaklı Location), mixed <i>Cedrus libani</i> forest, 1517 m, N 37°08'847" – E 31°19'742", 02.vi.2009, Çinbilgel 5026; İbradı, Çeşgar Plateau, rock, 1623 m, N 37°17'463" – E 31°23'699", 01.vii.2010, Çinbilgel 6854, (H).
Hypericaceae		
	<i>Hypericum confertum</i> Choisy subsp. <i>stenobotrys</i> (Boiss.) Holmboe	Manavgat, Sulucan Hill, snow patches – meadows, 2073 m, N 37°11'515" – E 31°18'836", 11.vi.2009, Çinbilgel 5374; İbradı, Melik Plateau, dwarf bush-thorny cushion formation, 1503 m, N 37°16'190" – E 31°24'575", 01.vii.2010, Çinbilgel 6769; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest, 1347 m, N 37°15'332" – E 31°24'499", 02.vii.2010, Çinbilgel 7015, E. Medit. (mt.), (H).

	<i>Hypericum hircinum</i> L. subsp. <i>majus</i> (Aiton) N.Robson	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, streamsides, 628 m, 05.viii.2009, Çinbilgel 5735; Manavgat, Çay Neighbourhood surroundings (Değirmenderesi Bridge), streamsides, 705 m, N 37°12'138" – E 31°16'452", 07.vi.2011, Çinbilgel 7880, <i>Medit.</i> , (Ph).
	<i>Hypericum origanifolium</i> Willd. var. <i>origanifolium</i>	Manavgat, Kaldırım Mountain (Hıncırak Location), mixed <i>Cedrus libani</i> forest, 1475 m, N 37°09'368" – E 31°20'410", 01.vi.2009, Çinbilgel 4899; Manavgat, between Derme Plateau and Sürüçikan Mountain, rocky hills, 2223 m, N 37°12'479" – E 31°19'531", 30.vi.2009, Çinbilgel 5596; İbradı, Suçkan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7080, (H).
	<i>Hypericum perforatum</i> L.	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7711; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 622 m, N 37°11'707" – E 31°16'064", 06.vi.2011, Çinbilgel 7745, <i>Medit.</i> , (H).
	<i>Hypericum perforatum</i> L. subsp. <i>veronense</i> (Schränk) H.Linb.	Manavgat, around of Burmahan Neighbourhood, roadsides, 700 m, N 37°10'623" – E 31°16'847", 06.vi.2011, Çinbilgel 7792; Manavgat, around of Burmahan Neighbourhood (Payır Location), roadsides, 771 m, N 37°09'673" – E 31°17'034", 06.vi.2011, Çinbilgel 7855, (H).
Iridaceae		
	* <i>Crocus antalyensis</i> B.Mathew subsp. <i>antalyensis</i>	İbradı, Gevenli Location, under the <i>Pinus brutia</i> forest, 1102 m, N 37°12'287" – E 31°24'574", 05.iii.2011, Çinbilgel 7338; İbradı, Yaylacık Mountain, under the <i>Pinus brutia</i> forest, 1034 m, N 37°10'543" – E 31°25'356", 06.iii.2011, Çinbilgel 7380, <i>End.</i> , <i>E. Medit.</i> , (G).
	* <i>Crocus asumaniae</i> B.Mathew & T.Baytop	Manavgat, Kızılpınar Plateau (between Gökçebel Location and Katranlıboğazi Location), <i>Cedrus libani</i> – <i>Pinus nigra</i> forest clearings, 1702 m, N 37°14'046" – E 31°19'447", 10.x.2009, Çinbilgel 5790, <i>End.</i> , <i>E. Medit.</i> , (G).
	* <i>Crocus biflorus</i> Mill. subsp. <i>isauricus</i> (Siehe ex Bowles) B.Mathew	Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest and stony slopes, 1731 m, N37° 09' 303" – E031° 20' 650", 26.iii.2009, Çinbilgel 4534; İbradı, Toka Plateau Breach, <i>Cedrus libani</i> forest clearings, 1696 m, N 37°13'718" – E 31°23'256", 28.iii.2010, Çinbilgel 5906; İbradı, Yaylacık Mountain, <i>Pinus brutia</i> forest, 1034 m, N 37°10'543" – E 31°25'356", 06.iii.2011, Çinbilgel 7381, <i>End.</i> , <i>E. Medit.</i> , (G).
	<i>Crocus chrysanthus</i> (Herb.) Herb.	Manavgat, Kepez Plateau (İnçayırın Location), meadows, 1674 m, N 37°10'569" – E 31°21'292", 03.v.2009, Çinbilgel 4723; İbradı, Aşağıtoka Plateau, moist – stony places, 1521 m, N 37°13'187" – E 31°22'678", 27.iii.2010, Çinbilgel 5866; İbradı, Çukuryurt Plateau, moist stony places, 1272 m, N 37°12'077" – E 31°24'031", 05.iii.2011, Çinbilgel 7343, (G).
	<i>Crocus speciosus</i> M.Bieb. subsp. <i>speciosus</i>	Manavgat, Kızılpınar Plateau, <i>Cedrus libani</i> forest clearings, 1496 m, N 37°14'385" – E 31°19'343", 10.x.2009, Çinbilgel 5762; Manavgat, Kızılpınar Plateau (Hebil Plateau Location), <i>Cedrus libani</i> forest clearings, 1713 m, N 37°13'988" – E 31°19'329", 10.x.2009, Çinbilgel 5789; İbradı, Melik Plateau, <i>Pinus nigra</i> forest clearings – meadows, 1431 m, N 37°15'838" – E 31°24'585", 05.x.2010, Çinbilgel 7289, (G).
	<i>Gladiolus anatolicus</i> (Boiss.) Stapf	Manavgat, Kızılpınar Plateau way, Kavaklı Location, <i>Juniperus excelsa</i> forest, 1323 m, N 37°13'974" – E 31°18'405", 07.vi.2011, Çinbilgel 7922, <i>E. Medit.</i> , (G).
	<i>Gladiolus italicus</i> Mill.	İbradı, between Toka Plateau and Belova Location, slopes, 1528 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3337, (G).
	* <i>Iris schachtii</i> Markgr.	İbradı, Yirme Hill, rocky hills, 1820 m, N 37°14'717" – E 31°23'178", 18.v.2008, Çinbilgel 3237; İbradı, Toka Plateau way, Belova Location, stony – rocky places, 1528 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3326; İbradı, Toka Plateau Breach, stony slopes, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5967, <i>End.</i> , <i>Ir. – Tur.</i> , (G).
Juglandaceae		
	<i>Juglans regia</i> L.	İbradı, Suçkan Location way, streamsides, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2781; Manavgat, Akpınar Plateau, roadsides, 1416 m, N 37°09'357" – E 31°20'711", 11.x.2009, Çinbilgel 5827, (Ph).
Juncaceae		
	<i>Juncus articulatus</i> L. subsp. <i>articulatus</i>	İbradı, Melik Plateau, wet places, streamsides, 1404 m, N 37°15'734" – E 31°24'641", 01.vii.2010, Çinbilgel 6740; İbradı, Melik Plateau, streamsides, 1537 m, N 37°15'549" – E 31°24'633", 02.vii.2010, Çinbilgel 6944; İbradı, Melik Plateau, streamsides, 1388 m, N 37°15'606" – E 31°24'669", 05.x.2010, Çinbilgel 7321, <i>Euro-Sib.</i> , (H).
	<i>Juncus inflexus</i> L. subsp. <i>inflexus</i>	Manavgat, Kepez Plateau, wet places, 1636 m, N 37°10'971" – E 31°21'114", 03.vi.2009, Çinbilgel 5174; İbradı, Melik Plateau, wet places, streamsides, 1404 m, N 37°15'734" – E 31°24'641", 01.vii.2010, Çinbilgel 6739, (H).
Lamiaceae		
	* <i>Ajuga bombycina</i> Boiss.	Manavgat, Derme Plateau, rock, 1759 m, N 37°12'512" – E 31°20'675", 19.vii.2008, Çinbilgel 4485; Manavgat, Kovacık Mountain (around of Dernebeli Location), dwarf bush formation, 1867 m, N 37°13'751" – E 31°20'136", 29.vi.2009, Çinbilgel 5466; İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1147 m, N 37°12'229" – E 31°24'506", 02.vii.2010, Çinbilgel 7237, <i>End.</i> , <i>E. Medit.</i> , (H).

	<i>Ajuga chamaepitys</i> (L.) Schreb. subsp. <i>chia</i> (Schreb.) Arcang.	Manavgat, Akpınar Plateau (Sumaklı Location), mixed <i>Cedrus libani</i> forest, 1517 m, N 37°08'847" – E 31°19'742", 02.vi.2009, Çinbilgel 5031; İbradı, Çeşgar Plateau, rock, 1621 m, N 37°17'565" – E 31°24'085", 01.vii.2010, Çinbilgel 6899; İbradı, Gevenli Location, stony places, 1075 m, N 37°12'345" – E 31°24'803", 02.vii.2010, Çinbilgel 7147, (H).
	<i>Ajuga chamaepitys</i> (L.) Schreb. subsp. <i>cuneatifolia</i> (Stapf) P.H.Davis	Manavgat, Akpınar Plateau (Yumrutaş Location), mixed <i>Cedrus libani</i> forest, 1520 m, N 37°09'840" – E 31°21'028", 19.v.2009, Çinbilgel 4781; Manavgat, Kovacık Mountain (around of Dermebeli Location), dwarf bush-thorny cushion formation, 1867 m, N 37°13'751" – E 31°20'136", 29.vi.2009, Çinbilgel 5461; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 1933 m, N 37°16'534" – E 31°22'619", 17.vi.2010, Çinbilgel 6451, (H).
	<i>Ajuga chamaepitys</i> (L.) Schreb. subsp. <i>palaestina</i> (Boiss.) Bornm.	Manavgat, Kaldırım Mountain (Hıncırak Location), mixed <i>Cedrus libani</i> forest, 1475 m, N 37°09'368" – E 31°20'410", 01.vi.2009, Çinbilgel 4908; Manavgat, Burunçalı Location (Dardaş), mixed <i>Cedrus libani</i> forest, 1558 m, N 37°08'772" – E 31°19'511", 02.vi.2009, Çinbilgel 5095, E. Medit., (H).
	* <i>Ballota cristata</i> P.H.Davis	İbradı, Sürüçkan Mountain, rocky places, 1947 m, N 37°12'779" – E 31°19'826", 10.vi.2008, Çinbilgel 3782; Manavgat, Burunçalı Location, slopes, 1846 m, N 37°09'124" – E 31°18'924", 02.vi.2009, Çinbilgel 4984; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çinbilgel 6717, End., E. Medit., (Ch).
	<i>Clinopodium graveolens</i> subsp. <i>rotundifolium</i> (Pers.) Govaerts	Manavgat, Kızılıpınar Plateau surroundings (eastern of Nurboğazi Location), stony places, 1962 m, N 37°14'256" – E 31°20'649", 11.vi.2008, Çinbilgel 3849; Manavgat, Kaldırım Mountain, under the <i>Cedrus libani</i> forest, 1551 m, N 37°09'461" – E 31°20'374", 01.vi.2009, Çinbilgel 4877; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 1840 m, N 37°16'441" – E 31°22'564", 17.vi.2010, Çinbilgel 6459, (Th).
	<i>Clinopodium vulgare</i> L. subsp. <i>vulgare</i>	İbradı, Suçikan Location way, under the forest, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4423; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1470 m, N 37°15'607" – E 31°24'429", 30.vi.2010, Çinbilgel 6634; İbradı, Suçikan Location, streamsides, 1200 m, N 37°13'958" – E 31°24'208", 02.vii.2010, Çinbilgel 7115, (H).
	<i>Lamium amplexicaule</i> L. var. <i>amplexicaule</i>	Manavgat, around of Kepezbeli Location, stony places, 1485 m, N 37°09'744" – E 31°21'083", 03.v.2009, Çinbilgel 4688; İbradı, Akmur Location, stony places, 1044 m, N 37°11'687" – E 31°24'880", 28.iii.2010, Çinbilgel 5948; Manavgat, Çay Neighbourhood, Kızılıpınar Plateau way (Düğnük Plateau), <i>Pinus brutia</i> forest clearings, 1163 m, N 37°13'385" – E 31°17'716", 27.iii.2011, Çinbilgel 7506, Euro-Sib., (Th).
	* <i>Lamium eriocephalum</i> Benth.	İbradı, Melik Mountain (Çeşgar Hill), screes, 2040 m, N 37°15'610" – E 31°23'438", 18.v.2008, Çinbilgel 3313; İbradı, Melik Mountain, screes, 2210 m, N 37°15'862" – E 31°23'599", 18.v.2008, Çinbilgel 3319; Manavgat, Kaldırım Mountain, screes, 1802 m, N 37°09'879" – E 31°20'418", 01.vi.2009, Çinbilgel 4918, End., E. Medit. (mt.), (H).
	<i>Lamium garganicum</i> L. subsp. <i>striatum</i> (Sm.) Hayek var. <i>striatum</i>	İbradı, Melik Mountain (Hasan Hill), stony hillsides, 2049 m, N 37°16'896" – E 31°23'156", 18.vi.2008, Çinbilgel 4326; İbradı, around of Yirme Hill, <i>Pinus nigra</i> forest, 1550 m, N 37°14'328" – E 31°22'786", 08.v.2010, Çinbilgel 6025; İbradı, Suçikan Location, streamsides, 1259 m, N 37°14'503" – E 31°24'173", 02.vii.2010, Çinbilgel 7062; İbradı, Güzle Plateau, Juniper forest, 1306 m, N 37°09'817" – E 31°23'980", 20.iv.2008, Çinbilgel 2831; Manavgat, Burunçalı Location, slopes, 1935 m, N 37°09'133" – E 31°19'062", 02.vi.2009, Çinbilgel 5071; İbradı, Melik Mountain (Orta Hill), windy hills, 2175 m, N 37°16'298" – E 31°23'052", 17.vi.2010, Çinbilgel 6373, Medit., (H).
	<i>Lamium macrodon</i> Boiss. & Huet	İbradı, Söbüova Location (around of Akpınar Location), under the forest, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3536; İbradı, Aşağıtoka Plateau, stony places, 1521 m, N 37°14'190" – E 31°22'688", 28.iii.2010, Çinbilgel 5879; İbradı, Çukuryurt Plateau, damp places, 1229 m, N 37°12'385" – E 31°24'069", 05.iii.2011, Çinbilgel 7347, Ir. – Tur., (Th).
	* <i>Marrubium astracanicum</i> Jacq. subsp. <i>macrodon</i> (Bornm.) P.H.Davis	Manavgat, Derme Plateau, rocky places, 1740 m, N 37°12'694" – E 31°20'565", 19.vii.2008, Çinbilgel 4488; Manavgat, Sulucan Hill, rock, 2081 m, N 37°11'492" – E 31°18'857", 11.vi.2009, Çinbilgel 5387; İbradı, Yukarıtoka Plateau, dwarf bush formation, 1635 m, N 37°15'828" – E 31°22'221", 17.vi.2010, Çinbilgel 6273, End., (Ch).
	* <i>Marrubium globosum</i> Montbret & Aucher ex Benth. subsp. <i>globosum</i>	Manavgat, Susuzkır Hill, <i>Pinus nigra</i> forest, 1718 m, N 37°16'910" – E 31°21'829", 12.vi.2008, Çinbilgel 3976; İbradı, Melik Mountain (eastern hillsides of Hasan Hill), rocky slopes, 2020 m, N 37°16'897" – E 31°23'181", 18.vi.2008, Çinbilgel 4306; İbradı, Melik Plateau, forest clearings – stony places, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6569, End., Ir. – Tur., (Ch).
	* <i>Marrubium globosum</i> Montbret & Aucher ex Benth. subsp. <i>micranthum</i> (Boiss. & Heldr.) P.H.Davis	İbradı, Çeşgar Plateau, stony slopes, 1659 m, N 37°17'402" – E 31°24'250", 17.vi.2008, Çinbilgel 4229; Manavgat, Kocatop Mountain (Örencik Location), dwarf bush formation, 1929 m, N 37°11'697" – E 31°18'642", 11.vi.2009, Çinbilgel 5434; İbradı, around of Toka Plateau Breach, stony slopes, 1570 m, N 37°13'602" – E 31°23'417", 09.v.2010, Çinbilgel 6126, End., E. Medit., (Ch).
	<i>Marrubium vulgare</i> L.	İbradı, between Taşkesiği Location and Cevizlidere Location, under the forest, 1143 m, N 37°13'710" – E 31°24'146", 18.vi.2008, Çinbilgel 4373; İbradı, Melik Plateau, stony places, 1452 m, N 37°15'960" – E 31°24'578",

		01.vii.2010, Çinbilgel 6748; Manavgat, Yelibelen Location, <i>Pinus brutia</i> forest, 1048 m, N 37°09'140" – E 31°17'606", 06.vi.2011, Çinbilgel 7862, Medit., (Ch).
	<i>Melissa officinalis</i> L. subsp. <i>officinalis</i>	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, streamsides, 628 m, 05.viii.2009, Çinbilgel 5738; Manavgat, around of Burmahan Neighbourhood (Öküztası Location), <i>Pinus brutia</i> forest, 683 m, N 37°10'293" – E 31°16'897", 06.vi.2011, Çinbilgel 7798, E. Medit., (H).
	<i>Mentha aquatica</i> L.	Manavgat, Burmahan Neighbourhood, Erenli Location, streamsides, 611 m, N 37°09'918" – E 31°16'962", 26.iii.2011, Çinbilgel 7463, (H).
	<i>Mentha longifolia</i> (L.) L. subsp. <i>thyphoides</i> (Briq.) Harley	Manavgat, Akpınar Plateau (Kelsu Location), watersides, 1573 m, N 37°09'507" – E 31°20'312", 01.vi.2009, Çinbilgel 4888; İbradı, Suçikan Location, streamsides, 1251 m, N 37°14'475" – E 31°24'186", 02.vii.2010, Çinbilgel 7108; İbradı, Suçikan Location, streamsides, 1220 m, N 37°14'429" – E 31°24'260", 02.vii.2010, Çinbilgel 7113, (H).
	<i>Mentha x piperita</i> L.	Manavgat, Kızılpınar Plateau, watersides, 1805 m, N 37°14'086" – E 31°20'164", 09.vi.2008, Çinbilgel 3700; Manavgat, Derme Plateau, gardens, 1699 m, N 37°12'911" – E 31°20'427", 05.viii.2009, Çinbilgel 5710, (H).
	<i>Mentha spicata</i> L. subsp. <i>condensata</i> (Briq.) Greuter & Burdet	Manavgat, Derme Plateau (Zahmedin Plateau Location), stony – wet places, 1761 m, N 37°12'348" – E 31°20'456", 20.vii.2008, Çinbilgel 4525, Medit., (H).
	<i>Mentha spicata</i> L. subsp. <i>spicata</i>	Manavgat, Kepez Plateau (İncayırı Location), stony slopes, 1649 m, N 37°11'195" – E 31°20'964", 11.x.2009, Çinbilgel 5840; İbradı, Melik Plateau, streamsides, 1404 m, N 37°15'734" – E 31°24'641", 01.vii.2010, Çinbilgel 6742; İbradı, Cevizlidere Location, streamsides, 1140 m, N 37°13'377" – E 31°24'211", 05.x.2010, Çinbilgel 7276, (H).
	<i>Micromeria graeca</i> (L.) Benth. ex Reichb. subsp. <i>graeca</i>	İbradı, between Taşkesiği Location and Cevizlidere Location, under the forest, 1143 m, N 37°13'710" – E 31°24'146", 18.vi.2008, Çinbilgel 4374; İbradı, Melik Plateau, under the <i>Pinus nigra</i> forest, stony places, 1496 m, N 37°15'607" – E 31°24'380", 30.vi.2010, Çinbilgel 6590; İbradı, Cevizlidere Location, rocky places, 1181 m, N 37°13'782" – E 31°24'185", 02.vii.2010, Çinbilgel 7145, Medit., (H).
	<i>Micromeria myrtifolia</i> Boiss. & Hohen.	İbradı, Suçikan Location way, stony rocky places, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4412; İbradı, Melik Plateau, stony places, 1486 m, N 37°15'604" – E 31°24'398", 30.vi.2010, Çinbilgel 6598; İbradı, Cevizlidere Location, rocky places, 1181 m, N 37°13'782" – E 31°24'185", 02.vii.2010, Çinbilgel 7144, E. Medit., (Ch).
	* <i>Nepeta cadmea</i> Boiss.	İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1241 m, N 37°14'491" – E 31°24'214", 02.vii.2010, Çinbilgel 7095; İbradı, Cevizlidere Location, streamsides, 1163 m, N 37°13'696" – E 31°24'146", 02.vii.2010, Çinbilgel 7118; İbradı, around of Melik Plateau, <i>Pinus nigra</i> forest, 1255 m, N 37°14'931" – E 31°24'420", 03.vii.2011, Çinbilgel 8029, End., E. Medit., (H).
	<i>Nepeta cilicia</i> Boiss. ex Benth.	Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), rocky places, 1734 m, N 37°13'822" – E 31°19'204", 09.vi.2009, Çinbilgel 5201; İbradı, Melik Plateau, forest clearings – stony places, 1406 m, N 37°15'466" – E 31°24'469", 30.vi.2010, Çinbilgel 6682; İbradı, Suçikan Location, streamsides, 1252 m, N 37°14'487" – E 31°24'182", 02.vii.2010, Çinbilgel 7099, E. Medit., (mt.), (H).
	<i>Nepeta italica</i> L.	İbradı, Çeşgar Plateau Location, dwarf bush formation, 1617 m, N 37°17'485" – E 31°23'565", 17.vi.2008, Çinbilgel 4218; Manavgat, Kocatop Mountain (Örencik Location), dwarf bush formation, 1861 m, N 37°11'829" – E 31°18'607", 11.vi.2009, Çinbilgel 5418; İbradı, Melik Plateau, <i>Pinus nigra</i> forest – stony places, 1486 m, N 37°15'604" – E 31°24'398", 30.vi.2010, Çinbilgel 6594, Medit., (H).
	<i>Nepeta nuda</i> L. subsp. <i>albiflora</i> (Boiss.) Gams	İbradı, Toka Plateau, screes, 1537 m, N 37°14'189" – E 31°22'385", 19.v.2008, Çinbilgel 3356; Manavgat, Burunçalı Location, slopes, 1935 m, N 37°09'133" – E 31°19'062", 02.vi.2009, Çinbilgel 5070; İbradı, Çeşgar Plateau Breach, stony places, 1682 m, N 37°17'288" – E 31°24'324", 01.vii.2010, Çinbilgel 6910, (H).
	<i>Origanum bilgeri</i> P.H.Davis	İbradı, Akmuar Location, rock, 1042 m, N 37°11'701" – E 31°24'851", 02.vii.2010, Çinbilgel 7229; İbradı, Melik Plateau, stony slopes, 1374 m, N 37°15'565" – E 31°24'628", 05.v.2010, Çinbilgel 7282; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest clearings, 1486 m, N 37°16'052" – E 31°24'647", 05.x.2010, Çinbilgel 7315, End., E. Medit., (Ch).
	<i>Origanum majorana</i> L.	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, <i>Pinus brutia</i> forest, 667 m, N 37°11'995" – E 31°16'386", 10.x.2009, Çinbilgel 5750; Manavgat, Burmahan Neighbourhood, Akpınar Plateau way, roadsides, 798 m, N 37°09'483" – E 31°17'167", 11.x.2009, Çinbilgel 5813, E. Medit., (Ch).
	* <i>Origanum minutiflorum</i> O.Schwarz & P.H.Davis	İbradı, Between Gavur Breach and Etülgenboğazi, <i>Pinus brutia</i> forest, 1186 m, N 37°09'074" – E 31°25'116", 20.iv.2008, Çinbilgel 2976; İbradı, Aşağıtoka Plateau, debris, 1547 m, N 37°13'903" – E 31°22'326", 19.v.2008, Çinbilgel 3390; Manavgat, Kovacık Mountain surroundings, rocky places, 1807 m, N 37°13'996" – E 31°20'274", 25.v.2008, Çinbilgel 3652, End., E. Medit., (Ch).
	* <i>Origanum saccatum</i> P.H.Davis	İbradı, Suçikan Location way, under the forest – rocky places, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4411; Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, under the <i>Pinus brutia</i> forest, 635 m,

		N 37°11'846" – E 31°16'238", 10.x.2009, Çinbilgel 5751; İbradı, Suçikan Location, stony places, 1200 m, N 37°13'960" – E 31°24'209", 02.vii.2010, Çinbilgel 7096, End., E. Medit., (H).
	<i>Phlomis armeniaca</i> Willd.	Manavgat, between Dermebeli Location and Sürüçikan Mountain, stony – rocky slopes, 1908 m, N 37°13'516" – E 31°20'206", 10.vi.2008, Çinbilgel 3749; İbradı, Çeşgar Location, stony slopes, 1588 m, N 37°17'747" – E 31°23'791", 17.vi.2008, Çinbilgel 4201; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 1933 m, N 37°16'534" – E 31°22'619", 17.vi.2010, Çinbilgel 6449, Ir. – Tur., (Ch).
	<i>Phlomis grandiflora</i> H.S.Thompson var. <i>grandiflora</i>	İbradı, Söbüova Location (around of Akpınar Location), under the forest, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3560; Manavgat, Kaldırım Mountain (Hıncırak Location), mixed <i>Cedrus libani</i> forest, 1475 m, N 37°09'368" – E 31°20'410", 01.vi.2009, Çinbilgel 4904; İbradı, Cevzlidere Location, streamsides, 1163 m, N 37°13'696" – E 31°24'146", 02.vii.2010, Çinbilgel 7116, E. Medit., (Ph).
	* <i>Phlomis leucophracta</i> P.H.Davis & Hub.–Mor.	Manavgat, Dügünük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747" – E 31°17'105", 13.vi.2008, Çinbilgel 4090; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7732, End., E. Medit., (Ph).
	<i>Phlomis lunariifolia</i> Sm.	Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4109; Manavgat, Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest, 702 m, N 37°09'905" – E 31°16'998", 06.vi.2011, Çinbilgel 7832; Manavgat, Çay Neighbourhood, roadsides, 690 m, N 37°12'122" – E 31°16'461", 07.vi.2011, Çinbilgel 7936, E. Medit., (Ph).
	<i>Phlomis samia</i> L.	Manavgat, Kızılıpınar Plateau way, Kavaklı Location, stony places, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4048; İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1223 m, N 37°14'468" – E 31°24'259", 02.vii.2010, Çinbilgel 7050, E. Medit., (Ph).
	<i>Prunella laciniata</i> (L.) L.	Manavgat, Kovacık Mountain foothills (Gökçebe Location), thorny cushion formation, 1695 m, N 37°13'984" – E 31°19'318", 11.vi.2009, Çinbilgel 5440; İbradı, Aşağıtoka Plateau, stony places, 1560 m, N 37°12'846" – E 31°22'666", 16.vi.2010, Çinbilgel 6239; İbradı, Melik Plateau surroundings (towards Suçikan Location), <i>Pinus nigra</i> forest, 1347 m, N 37°15'332" – E 31°24'499", 02.vii.2010, Çinbilgel 7004, Euro-Sib., (H).
	<i>Prunella orientalis</i> Bornm.	İbradı, Gevenli Location, <i>Pinus brutia</i> forest clearings, 1071 m, N 37°12'300" – E 31°24'793", 03.vii.2011, Çinbilgel 8009, Medit., (H).
	<i>Prunella vulgaris</i> L.	Manavgat, Kızılıpınar Plateau way, between Akin – Karahan Locations, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çinbilgel 4144; Manavgat, Kepez Plateau (İncayırın Location), wet places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5860; İbradı, Akmuar Location, damp places, 1042 m, N 37°11'845" – E 31°24'839", 02.vii.2010, Çinbilgel 7159, Euro-Sib., (H).
	* <i>Salvia adenocaulon</i> P.H.Davis	İbradı, Çeşgar Plateau Location, stony slopes, 1572 m, N 37°17'675" – E 31°23'744", 17.vi.2008, Çinbilgel 4203; Manavgat, Kızılıpınar Plateau, thorny cushion formation, 1766 m, N 37°14' 063" – E 31°20'070", 03.viii.2009, Çinbilgel 5615; İbradı, Melik Plateau, meadows, 1446 m, N 37°15'954" – E 31°24'591", 01.vii.2010, Çinbilgel 6755, End., Medit., (H).
	* <i>Salvia cadmica</i> Boiss. var. <i>cadmica</i>	İbradı, Çeşgar Plateau, rock, 1623 m, N 37°17'463" – E 31°23'699", 01.vii.2010, Çinbilgel 6856, End., (H).
	<i>Salvia microstegia</i> Boiss. & Balansa	İbradı, Yukarıtoka Plateau (between Between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3064; Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), thorny cushion formation, 1758 m, N 37°13'773" – E 31°19'193", 09.vi.2009, Çinbilgel 5194; İbradı, Melik Mountain (Orta Hill), dwarf bush formation, 2153 m, N 37°16'186" – E 31°22'903", 17.vi.2010, Çinbilgel 6338, Ir. – Tur., (H).
	<i>Salvia sclarea</i> L.	Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4112; İbradı, Cevzlidere Location, stony places, 1200 m, N 37°13'892" – E 31°24'189", 02.vii.2010, Çinbilgel 7138, (H).
	<i>Salvia tomentosa</i> Mill.	Manavgat, Susuzkır Hill, <i>Pinus nigra</i> forest, 1718 m, N 37°16'910" – E 31°21'829", 12.vi.2008, Çinbilgel 3977; İbradı, Suçikan Location way, under the forest, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4409; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1462 m, N 37°15'665" – E 31°24'474", 30.vi.2010, Çinbilgel 6610, Medit., (Ch).
	<i>Salvia verbenaca</i> L.	İbradı, Akmuar Location, Juniper Forest clearings, 1067 m, N 37°11'986" – E 31°24'842", 05.x.2010, Çinbilgel 7323, Medit., (H).
	<i>Salvia virgata</i> Jacq.	İbradı, Boğazyurt Location, stony places, 1032 m, N 37°11'838" – E 31°24'840", 19.v.2008, Çinbilgel 3597; Manavgat, Kızılıpınar Plateau, dwarf bush formation, 1750 m, N 37°13'650" – E 31°20'051", 29.vi.2009, Çinbilgel 5512; İbradı, Akmuar Location, Juniper Forest clearings, 1073 m, N 37°11'976" – E 31°24'786", 02.vii.2010, Çinbilgel 7153, Ir. – Tur., (H).
	<i>Salvia viridis</i> L.	Manavgat, Çay Neighbourhood, roadsides – stony places, 690 m, N 37°12'122" – E 31°16'461", 07.vi.2011, Çinbilgel 7937, Medit., (Th).

	<i>Satureja cuneifolia</i> Ten.	Manavgat, Burunçalı Location (Aslanboğazi), rocky hills, 1907 m, N 37°08'951" – E 31°19'501", 02.vi.2009, Çinbilgel 5052; İbradı, Toka Plateau Breach surroundings, 1570 m, N 37°13'602" – E 31°23'417", 09.v.2010, Çinbilgel 6122; İbradı, Melik Plateau (towards Çeşgar Plateau Breach), stony places, 1448 m, N 37°15'942" – E 31°24'577", 05.x.2010, Çinbilgel 7294, Medit., (Ch).
	<i>Satureja thymbra</i> L.	Manavgat, between Çay and Burmahan Neighbourhoods, <i>Pinus brutia</i> forest, 700 m, N 37°11'519" – E 31°16'218", 06.vi.2011, Çinbilgel 7772, E. Medit., (Ch).
	* <i>Scutellaria brevibracteata</i> Stapf subsp. <i>brevibracteata</i>	Manavgat, Kızılpınar Plateau, thorny cushion formation, 1766 m, N 37°14'063" – E 31°20'070", 03.viii.2009, Çinbilgel 5610; Manavgat, Kızılpınar Plateau, dwarf bush formation, 1757 m, N 37°14'073" – E 31°19'973", 10.x.2009, Çinbilgel 5770; İbradı, Belova Location, stony places, 1519 m, N 37°13'719" – E 31°23'577", 02.vii.2010, Çinbilgel 7256, End., E. Medit., (H).
	<i>Scutellaria brevibracteata</i> Stapf subsp. <i>subvelutina</i> (Rech.f.) Greuter & Burdet	Manavgat, Kızılpınar Plateau, thorny cushion formation, 1788 m, N 37°14'441" – E 31°20'293", 03.viii.2009, Çinbilgel 5644; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çinbilgel 6706; İbradı, Suçikan Location, streamsides, 1251 m, N 37°14'475" – E 31°24'186", 02.vii.2010, Çinbilgel 7104, E. Medit., (H).
	<i>Scutellaria orientalis</i> L. subsp. <i>alpina</i> (Boiss.) O.Schwarz var. <i>alpina</i>	İbradı, Yukarıtoka Plateau (between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3079; Manavgat, Kovacık Mountain foothills (Gökçebel Location), thorny cushion formation, 1687 m, N 37°14'038" – E 31°19'336", 09.vi.2009, Çinbilgel 5186; İbradı, Yukarıtoka Plateau, <i>Pinus nigra</i> forest clearing, 1783 m, N 37°16'349" – E 31°22'313", 17.vi.2010, Çinbilgel 6465, (H).
	* <i>Scutellaria salviifolia</i> Benth.	Manavgat, Kızılpınar Plateau way, Kavaklı Location, stony places, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4047; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, stony places, 1486 m, N 37°15'604" – E 31°24'398", 30.vi.2010, Çinbilgel 6599; İbradı, Melik Plateau surroundings (towards Suçikan Location), <i>Pinus nigra</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7033, End., (H).
	* <i>Sideritis condensata</i> Boiss. & Heldr.	İbradı, Suçikan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7088; İbradı, Suçikan Location, stony places, 1221 m, N 37°14'335" – E 31°24'250", 02.vii.2010, Çinbilgel 7092; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 665 m, N 37°11'982" – E 31°16'333", 07.vi.2011, Çinbilgel 7875, End., E. Medit., (H).
	* <i>Sideritis erythrantha</i> Boiss. & Heldr. var. <i>erythrantha</i>	İbradı, Melik Mountain (Hasan Hill), stony hillsides, 1761 m, N 37°16'962" – E 31°23'396", 18.vi.2008, Çinbilgel 4267; Manavgat, Kızılpınar Plateau surroundings (Nurboğazi Location), rock, 1913 m, N 37°14'166" – E 31°20'469", 04.viii.2009, Çinbilgel 5671; İbradı, Toka Plateau way, Belova Location, stony places, 1660 m, N 37°13'649" – E 31°23'311", 02.vii.2010, Çinbilgel 7253, End., E. Medit., (Ch).
	* <i>Sideritis pisidica</i> Boiss. & Heldr.	İbradı, Melik Plateau, stony slopes, 1615 m, N 37°17'572" – E 31°24'080", 17.vi.2008, Çinbilgel 4186; İbradı, around of Çeşgar Plateau, rocky places, 1619 m, N 37°17'574" – E 31°24'084", 01.vii.2010, Çinbilgel 6818; İbradı, Çeşgar Plateau, rocky places, 1632 m, N 37°17'458" – E 31°24'009", 01.vii.2010, Çinbilgel 6942, End., E. Medit., (H).
	<i>Stachys byzantina</i> K.Koch	İbradı, Çeşgar Plateau, stony slopes, 1626 m, N 37°17'559" – E 31°23'696", 17.vi.2008, Çinbilgel 4256; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1486 m, N 37°15'604" – E 31°24'398", 30.vi.2010, Çinbilgel 6595; İbradı, Suçikan Location, stony places, 1221 m, N 37°14'335" – E 31°24'250", 02.vii.2010, Çinbilgel 7091, Euro-Sib., (H).
	* <i>Stachys cretica</i> L. subsp. <i>anatolica</i> Rech.f.	Manavgat, Kocatop Mountain (Örencik Location), dwarf bush formation, 1861 m, N 37°11'829" – E 31°18'607", 11.vi.2009, Çinbilgel 5419; İbradı, Ekinlik Location surroundings, <i>Cedrus libani</i> forest (mixed with <i>Pinus nigra</i>), 1532 m, N 37°13'668" – E 31°23'510", 09.v.2010, Çinbilgel 6145; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7190, End., Ir. – Tur., (H).
	* <i>Stachys citrina</i> Boiss. & Heldr. subsp. <i>citrina</i>	Manavgat, Kovacık Mountain surroundings, on the rock, 1887 m, N 37°14'043" – E 31°20'377", 09.vi.2008, Çinbilgel 3680; İbradı, Melik Mountain (Hasan Hill), slopes, 2000 m, N 37°17'002" – E 31°23'605", 18.vi.2008, Çinbilgel 4353; Manavgat, Kocatop Mountain, windy hills, 2080 m, N 37°12'398" – E 31°18'732", 10.vi.2009, Çinbilgel 5357, End., E. Medit., (Ch).
	<i>Stachys lavandulifolia</i> Vahl	İbradı, Yukarıtoka Plateau (between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3066; İbradı, Melik Mountain (Hasan Hill), rocky slopes, 2020 m, N 37°16'897" – E 31°23'181", 18.vi.2008, Çinbilgel 4309; Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), thorny cushion formation, 1771 m, N 37°13'824" – E 31°19'246", 09.vi.2009, Çinbilgel 5270, Ir. – Tur., (H).
	<i>Stachys woronowii</i> (Schischk. ex Grossh.) R.R.Mill.	İbradı, Aşağıtoka Plateau, debris, 1547 m, N 37°13'903" – E 31°22'326", 19.v.2008, Çinbilgel 3375; Manavgat, Kızılpınar Plateau (Katranlıboğazi Location), mixed <i>Cedrus libani</i> forest, 1705 m, N 37°14'064" – E 31°19'589", 09.vi.2009, Çinbilgel 5218; İbradı, Melik Plateau, forest clearings – stony places, 1406 m, N 37°15'466" – E 31°24'469", 30.vi.2010, Çinbilgel 6679, Ir. – Tur., (Th).

	<i>Teucrium chamaedrys</i> L. subsp. <i>chamaedrys</i>	Manavgat, Kızılıpınar Plateau, dwarf bush formation, 1747 m, N 37°14'087" – E 31°19'983", 29.vi.2009, Çinbilgel 5503; İbradı, Melik Plateau, forest clearings – stony places, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6561; İbradı, Tahtalı Location, <i>Pinus nigra</i> forest, 1275 m, N 37°12'721" – E 31°24'600", 02.vii.2010, Çinbilgel 7120, Euro-Sib., (Ch).
	<i>Teucrium chamaedrys</i> L. subsp. <i>lydium</i> O.Schwarz	Manavgat, Çağlan Breach, stony places, 1654 m, N 37°15'995" – E 31°21'198", 12.vi.2008, Çinbilgel 3939; İbradı, Gevenli Location, <i>Pinus brutia</i> forest clearings – stony places, 1071 m, N 37°12'300" – E 31°24'793", 03.vii.2011, Çinbilgel 8010, E. Medit., (H).
	<i>Teucrium lamiifolium</i> d'Urv. subsp. <i>lamiifolium</i>	Manavgat, between Kovacık Mountain and Sürüçikan Mountain, dwarf bush formation, 1965 m, N 37°12'796" – E 31°19'814", 30.vi.2009, Çinbilgel 5585; İbradı, Çeşgar Plateau surroundings, rocky places, 1619 m, N 37°17'574" – E 31°24'084", 01.vii.2010, Çinbilgel 6819, (H).
	<i>Teucrium polium</i> L. subsp. <i>polium</i>	İbradı, Melik Plateau, stony places, 1615 m, N 37°17'572" – E 31°24'080", 17.vi.2008, Çinbilgel 4179; Manavgat, between Kovacık Mountain and Dermebeli Location, dwarf bush formation, 1916 m, N 37°13'542" – E 31°20'204", 30.vi.2009, Çinbilgel 5565; İbradı, Çeşgar Plateau, rocky places, 1665 m, N 37°17'211" – E 31°24'325", 01.vii.2010, Çinbilgel 6870, (Ch).
	<i>Thymbra spicata</i> L. subsp. <i>spicata</i>	Manavgat, Çay Neighbourhood, Burmahan Neighbourhood way, under the <i>Pinus brutia</i> forest, 642 m, N 37°11'726" – E 31°16'096", 26.iii.2011, Çinbilgel 7402; between Çay and Burmahan Neighbourhoods, <i>Pinus brutia</i> forest, 700 m, N 37°11'519" – E 31°16'218", 06.vi.2011, Çinbilgel 7778, E. Medit., (Ch).
	<i>Thymus cilicicus</i> Boiss. & Balansa	İbradı, Aşağıtoka Plateau, stony slopes, 1527 m, N 37°12'804" – E 31°22'579", 16.vi.2010, Çinbilgel 6234; İbradı, Aşağıtoka Plateau, rocky hills, 1529 m, N 37°12'869" – E 31°22'570", 02.vii.2011, Çinbilgel 7984, E. Medit., (Ch).
	<i>Thymus leucotrichus</i> Hal. subsp. <i>leucotrichus</i>	Manavgat, Kocatop Mountain (Koca Hill = Köygöründüğü Location), windy hills, 2150 m, N 37°12'346" – E 31°18'633", 10.vi.2009, Çinbilgel 5345; İbradı, Melik Mountain (Orta Hill), rocky places, 1967 m, N 37°16'123" – E 31°22'618", 17.vi.2010, Çinbilgel 6311; İbradı, Akmuar Location, <i>Juniperus excelsa</i> forest, 1044 m, N 37°11'404" – E 31°24'996", 03.vii.2011, Çinbilgel 8035, E. Medit., (Ch).
	<i>Thymus longicaulis</i> C.Presl subsp. <i>chaubardii</i> (Rchb.f.) Jalas	Manavgat, Kovacık Mountain foothills, thorny cushion formation, 1851 m, N 37°13'209" – E 31°19'027", 10.vi.2009, Çinbilgel 5304; İbradı, Melik Plateau, dwarf bush formation, 1502 m, N 37°16'195" – E 31°24'573", 01.vii.2010, Çinbilgel 6935; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7184, Medit., (Ch).
	<i>Thymus longicaulis</i> C.Presl subsp. <i>longicaulis</i>	İbradı, Aşağıtoka Plateau, dwarf bush formation, 1522 m, N 37°12'779" – E 31°22'674", 02.vii.2011, Çinbilgel 7978; İbradı, Aşağıtoka Plateau, thorny cushion formation, 1526 m, N 37°12'919" – E 31°22'639", 16.vi.2010, Çinbilgel 6177, Euro-Sib., (Ch).
	<i>Thymus sipyleus</i> Boiss.	Manavgat, Kaldırım Mountain, thorny cushion formation, 1837 m, N 37°09'993" – E 31°20'451", 01.vi.2009, Çinbilgel 4923; Çandır Hill, dwarf bush formation, 1987 m, N 37°14'492" – E 31°21'151", 04.viii.2009, Çinbilgel 5683; İbradı, Melik Plateau, dwarf bush formation, 1825 m, N 37°16'842" – E 31°24'494", 01.vii.2010, Çinbilgel 6930, (Ch).
	<i>Vitex agnus – castus</i> L.	İbradı, between Akpınar Location and Suçikan Location, streamsides, 1091 m, N 37°13'075" – E 31°24'456", 18.vi.2008, Çinbilgel 4432, Medit., (Ph).
	<i>Ziziphora capitata</i> L.	Manavgat, Kızılıpınar Plateau way, Kavaklı Location, stony places, 1360 m, N 37°14'112" – E 31°18'573", 13.vi.2008, Çinbilgel 4038; Akpınar Plateau (Sumaklı Location), mixed <i>Cedrus libani</i> forest, 1517 m, N 37°08'847" – E 31°19'742", 02.vi.2009, Çinbilgel 5022; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7181, Ir. – Tur., (Th).
	<i>Ziziphora clinopodioides</i> Lam.	Manavgat, around of Kızılıpınar Plateau (Nurboğazi Location), rocky places, 1907 m, N 37°14'167" – E 31°20'459", 11.vi.2008, Çinbilgel 3843; Manavgat, around of Çandır Hill, dwarf bush formation, 1904 m, N 37°14'197" – E 31°20'828", 04.viii.2009, Çinbilgel 5660; İbradı, Toka Plateau, stony places, 1538 m, N 37°14'038" – E 31°22'783", 09.v.2010, Çinbilgel 6075, Ir. – Tur., (Ch).
Liliaceae		
	<i>Fritillaria acmopetala</i> Boiss. subsp. <i>acmopetala</i>	İbradı, Güzle Plateau, under the forest, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2806; İbradı, Gevenli Location, under the forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3522; İbradı, Gevenli Location, stony places, 1133 m, N 37°11'988" – E 31°24'420", 19.v.2011, Çinbilgel 7622, E. Medit., (G).
	* <i>Fritillaria acmopetala</i> Boiss. subsp. <i>wendelboi</i> Rix	İbradı, Yirme Hill, rocky hills, 1820 m, N 37°14'717" – E 31°23'178", 18.v.2008, Çinbilgel 3245; İbradı, hillsides of Melik Mountain (Çataldüzlülüğü Location), stony places, 1843 m, N 37°15'258" – E 31°23'363", 18.v.2008, Çinbilgel 3286, End., E. Medit., (G).
	* <i>Fritillaria crassifolia</i> Boiss. & A.Huet. subsp. <i>crassifolia</i>	İbradı, between Güzle Plateau – Yaylacık Mountain, screes, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çinbilgel 2885; İbradı, around of Yirme Hill, stony slopes, 1875 m, N 37°14'859" – E 31°23'148", 18.v.2008, Çinbilgel 3283; Manavgat, Burunçalı Location (Aslanboğazi), rocky hills, 1907 m, N 37°08'951" – E 31°19'501", 02.vi.2009, Çinbilgel 5048, End., Ir. – Tur., (G).

	<i>Fritillaria elwesii</i> Boiss.	İbradı, Suçikan Location surroundings, streamsides, 1257 m, N 37°14'782" – E 31°24'366", 02.vii.2010, Çınbilgel 7045; İbradı, Suçikan Location, streamsides, 1220 m, N 37°14'429" – E 31°24'260", 02.vii.2010, Çınbilgel 7112, E. Medit., (G).
	<i>Fritillaria pinardii</i> Boiss.	İbradı, Aşağıtoka Plateau, meadows, 1521 m, N 37°14'190" – E 31°22'688", 28.iii.2010, Çınbilgel 5880, Ir. – Tur., (G).
	* <i>Fritillaria whittallii</i> Baker	İbradı, Toka Plateau way, Belova Location, rocky places, 1723 m, N 37°13'458" – E 31°23'314", 19.v.2008, Çınbilgel 3451; Manavgat, Kepezbeli Location, stony slopes, 1641 m, N 37°10'300" – E 31°21'722", 19.v.2009, Çınbilgel 4773; Manavgat, Kaldırım Mountain (Aktünek Hill), windy hills, 2198 m, N 37°10'556" – E 31°20'329", 03.vi.2009, Çınbilgel 5128, End., E. Medit., (mt.), (G).
	<i>Gagea fibrosa</i> (Desf.) Schult. & Schult.f.	İbradı, Eynif Plain (Kışla Location), stony places, 954 m, N 37°09'724" – E 31°26'060", 19.iv.2009, Çınbilgel 4632, (G).
	<i>Gagea foliosa</i> (C.Presl) Schult. & Schult.f.	İbradı, Melik Mountain (hillsides of Çeşgar Hill), screens, 2000 m, N 37°15'593" – E 31°23'351", 18.v.2008, Çınbilgel 3310; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest and stony slopes, 1731 m, N 37°09'303" – E 31°20'650", 26.iii.2009, Çınbilgel 4537; İbradı, Çukuryurt Plateau, meadows, 1268 m, N 37°12'117" – E 31°24'023", 28.iii.2010, Çınbilgel 5929, (G).
	<i>Gagea granatellii</i> (Parl.) Parl.	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way (Düğnük Plateau), <i>Pinus brutia</i> forest clearings, 1163 m, N 37°13'385" – E 31°17'716", 27.iii.2011, Çınbilgel 7509, Medit., (G).
	<i>Gagea juliae</i> Pascher	Manavgat, Burmahan Neighbourhood, Akpınar Plateau way, Erenli Location, under the <i>Pinus brutia</i> forest, 718 m, N 37°09'958" – E 31°17'023", 26.iii.2011, Çınbilgel 7428; Manavgat, Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest clearings, 611 m, N 37°09'918" – E 31°16'962", 26.iii.2011, Çınbilgel 7464, E. Medit., (G).
	<i>Gagea villosa</i> (M.Bieb.) Sweet var. <i>villosa</i>	Manavgat, eastern hillsides of Kaldırım Mountain, dwarf bush formation, 1985 m, N 37°10'405" – E 31°20'777", 03.vi.2009, Çınbilgel 5118; İbradı, Aşağıtoka Plateau, meadows, 1521 m, N 37°14'190" – E 31°22'688", 28.iii.2010, Çınbilgel 5877; Manavgat, Çay Neighbourhood, Kızılpınar Plateau way (Düğnük Plateau), <i>Pinus brutia</i> forest clearings, 1163 m, N 37°13'385" – E 31°17'716", 27.iii.2011, Çınbilgel 7510, Medit., (G).
	<i>Tulipa armena</i> Boiss. var. <i>armena</i>	İbradı, Suçikan Location way, stony places, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çınbilgel 2755; Manavgat, Kepezbeli Location, stony slopes, 1641 m, N 37°10'300" – E 31°21'722", 19.v.2009, Çınbilgel 4770, Ir. – Tur., (G).
Linaceae		
	<i>Linum austriacum</i> L. subsp. <i>austriacum</i>	Manavgat, Kocatop Mountain foothills, dwarf bush formation, 1797 m, N 37°12'618" – E 31°18'470", 11.vi.2009, Çınbilgel 5376, (H).
	<i>Linum bienne</i> Mill.	İbradı, Toka Plateau way, 1520 m, N 37°14'174" – E 31°22'773", 08.v.2010, Çınbilgel 6035; İbradı, Aşağıtoka Plateau, dwarf bush-thorny cushion formation, 1526 m, N 37°13'831" – E 31°22'356", 20.v.2011, Çınbilgel 7659; İbradı, Aşağıtoka Plateau, wet meadows, 1514 m, N 37°13'772" – E 31°22'587", 08.vi.2011, Çınbilgel 7955, Medit., (H).
	* <i>Linum boissieri</i> Asch. & Sint. ex Boiss.	İbradı, Yukarıtoka Plateau (between Beydiğın Plateau and Göcebe Location), stony places, 1551 m, N 37°14'763" – E 31°22'308", 17.v.2008, Çınbilgel 3136; İbradı, Toka Plateau, stony places, 1520 m, N 37°14'332" – E 31°22'709", 08.v.2010, Çınbilgel 6033 & Eren; İbradı, Aşağıtoka Plateau, moist stony places, 1517 m, N 37°13'122" – E 31°22'525", 08.vi.2011, Çınbilgel 7946, End., E. Medit., (H).
	<i>Linum corymbulosum</i> Rchb.	Manavgat, around of Burmahan Neighbourhood, <i>Pinus brutia</i> forest, 695 m, N 37°11'319" – E 31°16'296", 06.vi.2011, Çınbilgel 7784, Medit., (Th).
Malvaceae		
	<i>Alcea biennis</i> Winterl	Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 680 m, N 37°10'071" – E 31°16'971", 06.vi.2011, Çınbilgel 7831/a, (H).
	<i>Malva neglecta</i> Wallr.	İbradı, Toka Plateau (Göcebe Location), rocky places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çınbilgel 3111; Manavgat, Kızılpınar Plateau, Nurboğazi Location, rocky places, 1907 m, N 37°14'167" – E 31°20'459", 11.vi.2008, Çınbilgel 3835; Manavgat, Çiğdem Hill surroundings, dwarf bush formation, 1916 m, N 37°14'086" – E 31°20'824", 04.viii.2009, Çınbilgel 5649, (H).
Moraceae		
	<i>Ficus carica</i> L. subsp. <i>carica</i>	Manavgat, Akin – Karahan Location, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çınbilgel 4140, Medit., (Ph).
	<i>Morus nigra</i> L.	(Ph), (Observation).
Myrtaceae		
	<i>Myrtus communis</i> L. subsp. <i>communis</i>	Manavgat, between Burmahan Neighbourhood and Çay Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°11'284" – E 31°16'291", 11.x.2009, Çınbilgel 5794; Manavgat, Çay Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°12'133" – E 31°16'452", 27.iii.2011, Çınbilgel 7480, Medit., (Ph).
Oleaceae		
	<i>Fontanesia phyllraeoides</i> Labill.	İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 974 m, N 37°10'600" – E 31°25'718", 26.ix.2010, Çınbilgel 7200; Manavgat, around of Burmahan

		Neighbourhood, streamsides, 693 m, N 37°11'151" – E 31°16'304", 06.vi.2011, Çinbilgel 7767, E. Medit., (Ph).
	<i>Fraxinus angustifolia</i> Vahl. subsp. <i>angustifolia</i>	İbradı, Suçikan Location, streamsides, 1259 m, N 37°14'503" – E 31°24'173", 02.vii.2010, Çinbilgel 7073; İbradı, Suçikan Location, streamsides, 1220 m, N 37°14'429" – E 31°24'260", 02.vii.2010, Çinbilgel 7111, (Ph).
	* <i>Fraxinus omus</i> L. subsp. <i>cilicica</i> (Lingelsh.) Yalt.	Manavgat, Akpınar Plateau (Doğruluk Location), <i>Cedrus libani</i> forest clearings, 1424 m, N 37°08'886" – E 31°20'013", 02.vi.2009, Çinbilgel 4973, End., E. Medit., (Ph).
	<i>Olea europaea</i> L. subsp. <i>europaea</i>	Manavgat, Kızılpınar Plateau way, Göcek Location, roadsides, 600 m, 13.vi.2008, Çinbilgel 4155, (Ph).
	<i>Phillyrea latifolia</i> L.	Manavgat, Akpınar Plateau way, damaged forest, 1306 m, N 37°08'484" – E 31°19'182", 11.x.2009, Çinbilgel 5821; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 974 m, N 37°10'600" – E 31°25'718", 26.ix.2010, Çinbilgel 7202, Medit., (Ph).
Onagraceae		
	<i>Epilobium gemmascens</i> C.A.Mey.	Manavgat, Kızılpınar Plateau, watersides, 1780 m, N 37°14'087" – E 31°20'158", 03.viii.2009, Çinbilgel 5640; Manavgat, Kızılpınar Plateau (Hebil Plateau), wet places, 1678 m, N 37°13'897" – E 31°19'263", 10.x.2009, Çinbilgel 5788; Manavgat, Kepez Plateau (İnçayırı Location), watersides – wet places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5857, (H).
	<i>Epilobium lanceolatum</i> Sebast. & Mauri	Manavgat, Eşekbeli Location, <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1750 m, N 37°15'074" – E 31°20'047", 12.vi.2008, Çinbilgel 4015; Manavgat, Kepez Plateau (İnçayırı Location), wet places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5858; İbradı, Çeşgar Plateau, rock, 1618 m, N 37°17'456" – E 31°23'710", 01.vii.2010, Çinbilgel 6845, (H).
	<i>Epilobium montanum</i> L.	İbradı, Melik Plateau, stony slopes, 1615 m, N 37°17'572" – E 31°24'080", 17.vi.2008, Çinbilgel 4182, Euro-Sib., (H).
	<i>Epilobium tetragonum</i> L. subsp. <i>tetragonum</i>	Manavgat, Kovacık Mountain foothills (Elmalı Location), moist stony places, 1615 m, N 37°13'748" – E 31°18'967", 11.vi.2009, Çinbilgel 5408, (H).
Orchidaceae		
	<i>Cephalanthera kurdica</i> Bornm. ex Kraenzl.	İbradı, Gevenli Location, under the <i>Pinus brutia</i> forest, 1108 m, N 37°12'316" – E 31°24'613", 17.v.2008, Çinbilgel 3017; İbradı, Gevenli Location, under the forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3524; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7708, Ir. – Tur., (G).
	<i>Cephalanthera rubra</i> (L.) Rich.	İbradı, Taşkesiği Location, under the <i>Pinus nigra</i> forest, 1111 m, N 37°14'142" – E 31°24'258", 18.vi.2008, Çinbilgel 4406; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1502 m, N 37°15'548" – E 31°24'365", 30.vi.2010, Çinbilgel 6655; Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 773 m, N 37°09'851" – E 31°16'989", 06.vi.2011, Çinbilgel 7846, (G).
	<i>Limodorum abortivum</i> (L.) Sw. var. <i>abortivum</i>	İbradı, between Gavur Breach and Etülgenboğazı, <i>Pinus brutia</i> forest, 1186 m, N 37°09'074" – E 31°25'116", 20.iv.2008, Çinbilgel 2973; İbradı, Gevenli Location, under the <i>Pinus brutia</i> forest, 1108 m, N 37°12'316" – E 31°24'613", 17.v.2008, Çinbilgel 3016, (G).
	* <i>Ophrys cilicica</i> Schltr.	İbradı, Suçikan Location, streamsides, 1220 m, N 37°14'429" – E 31°24'260", 02.vii.2010, Çinbilgel 7114, End., E. Medit., (G).
	<i>Ophrys phrygia</i> H.Fleischm. & Bornm.	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7704, Ir. – Tur., (G).
	<i>Orchis anatolica</i> Boiss.	İbradı, between Gavur Breach and Etülgenboğazı, <i>Pinus brutia</i> forest, 977 m, N 37°09'100" – E 31°25'485", 20.iv.2008, Çinbilgel 2986; Manavgat, Çay Neighbourhood, Eşekbeli Location, <i>Pinus nigra</i> forest, 1612 m, N 37°14'867" – E 31°19'864", 07.vi.2011, Çinbilgel 7902, E. Medit., (G).
	<i>Orchis tridentata</i> Scop.	Manavgat, Burunçalı Location (Dardaş), mixed <i>Cedrus libani</i> forest, 1558 m, N 37°08'772" – E 31°19'511", 02.vi.2009, Çinbilgel 5094, Medit., (G).
Orobanchaceae		
	<i>Orobanche alba</i> Stephan ex Willd. subsp. <i>alba</i>	İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1476 m, N 37°15'632" – E 31°24'434", 30.vi.2010, Çinbilgel 6580, (Vp).
	<i>Orobanche anatolica</i> Boiss. & Reut. ex Reut.	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest (mixed with <i>Pinus nigra</i>), 1660 m, N 37°13'683" – E 31°23'340", 09.v.2010, Çinbilgel 6061, (Vp).
	<i>Orobanche caryophyllacea</i> Sm.	Manavgat, Dermebeli Location, dwarf bush formation, 1935 m, N 37°13'656" – E 31°20'290", 29.vi.2009, Çinbilgel 5456; İbradı, Yukarıtoka Plateau, <i>Pinus nigra</i> forest clearing, 1783 m, N 37°16'349" – E 31°22'313", 17.vi.2010, Çinbilgel 6463, (Vp).
	<i>Orobanche cilicica</i> Beck	İbradı, Suçikan Location, streamsides, 1251 m, N 37°14'475" – E 31°24'186", 02.vii.2010, Çinbilgel 7106, (Vp).
	<i>Orobanche minor</i> Sm.	Manavgat, Sarıtaş Location, rock, 1998 m, N 37°15'078" – E 31°20'867", 04.viii.2009, Çinbilgel 5669, (Vp).
	<i>Orobanche mutellii</i> F.W.Schultz	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way (Kavaklı Location), <i>Juniperus excelsa</i> forest, 1323 m, N 37°13'974" – E 31°18'405", 07.vi.2011, Çinbilgel 7926, (Vp).
	<i>Orobanche nana</i> Noë ex Reut.	Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4824; İbradı, Melik Plateau, forest clearings – stony places, 1406 m, N 37°15'491" – E 31°24'478",

		30.vi.2010, Çinbilgel 6702; Manavgat, Akpınar Plateau way, <i>Pinus brutia</i> forest, 1310 m, N 37°08'495" – E 31°18'855", 06.vi.2011, Çinbilgel 7873, (Vp).
	<i>Parentucellia latifolia</i> (L.) Caruel subsp. <i>latifolia</i>	İbradı, Boğazyurt Location, stony places, 974 m, N 37°10'338" – E 31°25'772", 19.v.2011, Çinbilgel 7582; İbradı, Gevenli Location, <i>Pinus brutia</i> forest clearings, 1102 m, N 37°12'102" – E 31°24'531", 19.v.2011, Çinbilgel 7619, Medit., (Th).
Oxalidaceae		
	<i>Oxalis corniculata</i> L.	Manavgat, around of Burmahan Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°10'781" – E 31°16'754", 06.vi.2011, Çinbilgel 7780, (Th).
Paeoniaceae		
	<i>Paeonia mascula</i> (L.) Mill. subsp. <i>mascula</i>	İbradı, around of Yirme Hill (Dikenli Karlığı Location), damp places, 1814 m, N 37°14'700" – E 31°23'139", 18.v.2008, Çinbilgel 3218, (H).
Papaveraceae		
	<i>Corydalis erdelii</i> Zucc.	İbradı, Melik Mountain (Çeşgar Hill), stony slopes, 2100 m, N 37°15'610" – E 31°23'438", 18.v.2008, Çinbilgel 3316; Manavgat, Kepezbeli Location, stony slopes, 1674 m, N 37°10'329" – E 31°21'582", 03.v.2009, Çinbilgel 4703; İbradı, Toka Plateau Breach, <i>Cedrus libani</i> forest clearings, 1690 m, N 37°13'702" – E 31°23'278", 28.iii.2010, Çinbilgel 5908, (G).
	<i>Corydalis solida</i> (L.) Clairv. subsp. <i>solida</i>	İbradı, Suçkan Location way, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2763; Manavgat, Kepezbeli Location, stony slopes, 1674 m, N 37°10'329" – E 31°21'582", 03.v.2009, Çinbilgel 4704; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest – stony places, 1421 m, N 37°09'367" – E 31°20'728", 26.iii.2011, Çinbilgel 7450, (G).
	<i>Fumaria asepala</i> Boiss.	İbradı, between Güzle Plateau – Yaylacık Mountain, screes, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çinbilgel 2895; İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest – roadsides, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3470; İbradı, Melik Mountain (Çeşgar Hill), stony hills, 2264 m, N 37°16'008" – E 31°23'667", 18.vi.2010, Çinbilgel 6529, Ir. – Tur., (Th).
	<i>Glaucium leiocarpum</i> Boiss.	İbradı, Aşağıtoka Plateau, screes, 1547 m, N 37°13'903" – E 31°22'326", 19.v.2008, Çinbilgel 3387; Manavgat, Burunçalı Location, slopes, 1916 m, N 37°09'083" – E 31°19'415", 02.vi.2009, Çinbilgel 5002; İbradı, Melik Plateau, screes, 1491 m, N 37°15'544" – E 31°24'739", 02.vii.2010, Çinbilgel 6987, (H).
	* <i>Papaver pilosum</i> Sibth. & Sm. subsp. <i>sparsipilosum</i> (Boiss.) Kadereit	Manavgat, between Dermebeli and Sürüçkan Mountain, stony – rocky slopes, 1908 m, N 37°13'516" – E 31°20'206", 10.vi.2008, Çinbilgel 3748; Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), dwarf bush-thorny cushion formation, 1758 m, N 37°13'773" – E 31°19'193", 09.vi.2009, Çinbilgel 5193; İbradı, Melik Plateau, screes, 1491 m, N 37°15'544" – E 31°24'739", 02.vii.2010, Çinbilgel 6984, End., (H).
	<i>Papaver rhoeas</i> L.	İbradı, Söbüova Location, under the forest, 1057 m, N 37°12'262" – E 31°24'792", 25.v.2008, Çinbilgel 3532; Manavgat, Arğıçca Location – cutting area, <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4108; Manavgat, around of Burmahan Neighbourhood, roadsides, 700 m, N 37°10'623" – E 31°16'847", 06.vi.2011, Çinbilgel 7791, (Th).
Phyllanthaceae		
	<i>Andrachne telephioides</i> L.	Manavgat, Kaldırım Mountain (Hıncırak Location), mixed <i>Cedrus libani</i> forest, 1475 m, N 37°09'368" – E 31°20'410", 01.vi.2009, Çinbilgel 4896; İbradı, Çukuryurt Plateau surroundings, rock, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6167; İbradı, Melik Plateau surroundings (towards Suçkan Location), <i>Pinus nigra</i> forest, 1347 m, N 37°15'332" – E 31°24'499", 02.vii.2010, Çinbilgel 7012, (H).
Plantaginaceae		
	<i>Chaenorhinum minus</i> (L.) Lange subsp. <i>minus</i>	Manavgat, Düğnük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4083; Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4115, (Th).
	<i>Cochlidiosperma trichadena</i> (Jord. & Four.) Govaerts	İbradı, between Güzle Plateau – Yaylacık Mountain, damp places, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çinbilgel 2896; Manavgat, Kepezbeli Location, mixed <i>Cedrus libani</i> forest clearings, 1583 m, N 37°10'241" – E 31°21'544", 03.v.2009, Çinbilgel 4693; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest clearings, 1421 m, N 37°09'367" – E 31°20'728", 26.iii.2011, Çinbilgel 7449, Medit., (Th).
	<i>Digitalis cariensis</i> Boiss. ex Benth.	Manavgat, between Dermebeli and Sürüçkan Mountain, rocky slopes, 1917 m, N 37°12'891" – E 31°19'890", 10.vi.2008, Çinbilgel 3763; Manavgat, Dutluca Plateau, stony places, 1282 m, N 37°13'785" – E 31°18'213", 13.vi.2008, Çinbilgel 4061; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1428 m, N 37°15'422" – E 31°24'423", 30.vi.2010, Çinbilgel 6674, E. Medit., (H).
	* <i>Digitalis davisiana</i> Heywood	İbradı, between Suçkan Location and Akpınar Location, under the forest, 1068 m, N 37°12'609" – E 31°24'678", 18.vi.2008, Çinbilgel 4434; İbradı, Tahtalı Location, under the forest, 1085 m, N 37°12'611" – E 31°24'679", 02.vii.2010, Çinbilgel 7127; İbradı, Tahtalı Location, <i>Pinus brutia</i> forest, 1066 m, N 37°12'606" – E 31°24'675", 03.vii.2011, Çinbilgel 8012, End., E. Medit., (H).
	<i>Digitalis ferruginea</i> L. subsp. <i>ferruginea</i>	Manavgat, Derme Plateau, dwarf bush-thorny cushion formation, 1812 m, N 37°12'698" – E 31°20'052", 20.vii.2008, Çinbilgel 4496; İbradı, Melik Plateau,

		<i>Pinus nigra</i> forest, 1487 m, N 37°15'592" – E 31°24'391", 30.vi.2010, Çinbilgel 6647; İbradı, Melik Plateau surroundings (towards Suçikan Location), <i>Pinus nigra</i> forest, 1347 m, N 37°15'332" – E 31°24'499", 02.vii.2010, Çinbilgel 7014, Euro-Sib., (H).
	<i>Globularia trichosantha</i> Fisch. & C.A.Mey. subsp. <i>trichosantha</i>	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings (stony slopes), 1701 m, N 37°13'682" – E 31°23'231", 19.v.2008, Çinbilgel 3412; Manavgat, Kaldırım Mountain, dwarf bush-thorny cushion formation, 1837 m, N 37°09'993" – E 31°20'451", 01.vi.2009, Çinbilgel 4929; İbradı, around of Yirme Hill, <i>Pinus nigra</i> forest, 1600 m, N 37°14'120" – E 31°22'886", 08.v.2010, Çinbilgel 6010, Ir. – Tur., (H).
	<i>Kickxia elatine</i> (L.) Dumort subsp. <i>crinita</i> (Mabille) Greuter	Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4121; Manavgat, Kepez Plateau, moist stony places, 1634 m, N 37°10'611" – E 31°21'285", 11.x.2009, Çinbilgel 5862; İbradı, Akmuar Location, streamsides, 1042 m, N 37°11'719" – E 31°24'820", 02.vii.2010, Çinbilgel 7225, Medit., (Th).
	* <i>Linaria corifolia</i> Desf.	İbradı, Melik Plateau, under the <i>Pinus nigra</i> forest, 1496 m, N 37°15'607" – E 31°24'380", 30.vi.2010, Çinbilgel 6587, End., Ir. – Tur., (H).
	* <i>Linaria dumanii</i> A.Duran & Y.Menemen	İbradı, Eynif Plain, arid meadows, 949 m, N 37°08'004" – E 31°27'963", 19.v.2011, Çinbilgel 7570; İbradı, Eynif Plain, arid meadows, 947 m, N 37°07'654" – E 31°27'562", 02.vii.2011, Çinbilgel 7965, End., (Th).
	* <i>Linaria genistifolia</i> (L.) Mill. subsp. <i>confertiflora</i> (Boiss.) P.H.Davis	İbradı, Melik Plateau, pathway-stony places, 1553 m, N 37°16'606" – E 31°24'539", 01.vii.2010, Çinbilgel 6787; İbradı, Suçikan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7083, Ir. – Tur., (H).
	<i>Linaria pelisseriana</i> (L.) Mill.	Manavgat, around of Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest, 702 m, N 37°09'905" – E 31°16'998", 06.vi.2011, Çinbilgel 7834, Medit., (Th).
	<i>Linaria simplex</i> DC.	İbradı, Yukarıtoka Plateau (between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3065; İbradı, Melik Plateau, stony slopes, 1644 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4176, Medit., (Th).
	<i>Plantago holosteum</i> Scop.	Manavgat, Kızılıpınar Plateau, stony places, 1750 m, N 37°14'071" – E 31°19'981", 19.vii.2008, Çinbilgel 4455; Manavgat, Kovacık Mountain (Gökçebel Location), thorny cushion formation, 1709 m, N 37°14'032" – E 31°19'421", 09.vi.2009, Çinbilgel 5224; İbradı, Aşağıtoka Plateau, meadows, 1513 m, N 37°13'794" – E 31°22'474", 16.vi.2010, Çinbilgel 6250, Medit., (H).
	<i>Plantago lagopus</i> L.	İbradı, Boğazyurt Location, stony places, 974 m, N 37°10'338" – E 31°25'772", 19.v.2011, Çinbilgel 7584; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest clearings, 965 m, N 37°10'640" – E 31°25'709", 19.v.2011, Çinbilgel 7593; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1006 m, N 37°11'054" – E 31°25'243", 03.vii.2011, Çinbilgel 8041, Medit., (H).
	<i>Plantago lanceolata</i> L.	Manavgat, Arğıçca Breach Location (burning area), <i>Pinus brutia</i> forest clearings, 1090 m, N 37°13'042" – E 31°17'108", 13.vi.2008, Çinbilgel 4100; İbradı, Çeşgar Plateau Breach, stony places, 1682 m, N 37°17'288" – E 31°24'324", 01.vii.2010, Çinbilgel 6913; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7729, (H).
	<i>Plantago major</i> L. subsp. <i>intermedia</i> (Gilib.) Lange	Manavgat, Kızılıpınar Plateau, thorny cushion formation, 1861 m, N 37°14'688" – E 31°20'385", 03.viii.2009, Çinbilgel 5632, (H).
	<i>Veronica anagallis aquatica</i> L.	Manavgat, Derme Plateau, Kovantaşı Location, wet places, 1804 m, N 37°11'937" – E 31°20'141", 20.vii.2008, Çinbilgel 4501; Manavgat, around of Dermebeli Location, under the rock, 1902 m, N 37°13'711" – E 31°20'283", 29.vi.2009, Çinbilgel 5520; İbradı, Akmuar Location, wet places, 1042 m, N 37°11'845" – E 31°24'839", 02.vii.2010, Çinbilgel 7158, (H).
	<i>Veronica arvensis</i> L.	İbradı, Güzle Plateau way, Gavur Breach Location, forest, 1274 m, N 37°09'174" – E 31°24'799", 20.iv.2008, Çinbilgel 2951, Euro-Sib., (Th).
	<i>Veronica biloba</i> L.	Manavgat, Akpınar Plateau, stony places, 1403 m, N 37°09'414" – E 31°20'759", 19.v.2009, Çinbilgel 4751, Ir. – Tur., (Th).
	<i>Veronica bozakmanii</i> M.A.Fisch.	İbradı, Güzle Plateau (Mezarlı Location), moist stony places, 1287 m, N 37°10'193" – E 31°23'650", 18.iv.2009, Çinbilgel 4565; İbradı, Eynif Plain (Yarbaşı Location), stony places, 959 m, N 37°08'603" – E 31°27'459", 28.iii.2010, Çinbilgel 5962; İbradı, Aşağıtoka Plateau, stony places, 1555 m, N 37°13'162" – E 31°22'666", 20.v.2011, Çinbilgel 7652, Ir. – Tur., (Th).
	* <i>Veronica cuneifolia</i> D.Don subsp. <i>cuneifolia</i>	İbradı, Suçikan Location way, stony places, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2774; Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4827; İbradı, Toka Plateau Breach surroundings, <i>Cedrus libani</i> forest clearings – stony places, 1570 m, N 37°13'602" – E 31°23'417", 09.v.2010, Çinbilgel 6130, End., E. Medit., (H).
	* <i>Veronica cuneifolia</i> D.Don subsp. <i>isaurica</i> P.H.Davis	İbradı, Between Güzle Plateau – Yaylacık Mountain, screes, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çinbilgel 2893; Manavgat, Kocatop Mountain (Kırtağacıboğazı Location), rock, 1974 m, N 37°12'819" – E 31°18'917", 10.vi.2009, Çinbilgel 5293; İbradı, Melik Mountain (southern hillsides of Hasan Hill), dwarf bush formation, 2073 m, N 37°16'539" – E 31°23'096", 17.vi.2010, Çinbilgel 6433, End., E. Medit., (mt.), (H).

	<i>Veronica cymbalaria</i> Bodard	İbradı, southeast slopes of Yaylacık Mountain, under the <i>Cedrus libani</i> forest, 1356 m, N 37°09'929" – E 31°25'029", 19.iv.2009, Çinbilgel 4588; İbradı, Çukuryurt Plateau surroundings, rocky places, 1237 m, N 37°12'041" – E 31°24'160", 09.v.2010, Çinbilgel 6171; Manavgat, Çay Neighbourhood surroundings (Dikmen Hill), <i>Pinus brutia</i> forest, 816 m, N 37°12'492" – E 31°16'225", 07.vi.2011, Çinbilgel 7886, <i>Medit., (Th)</i> .
	* <i>Veronica elmaliensis</i> M.A.Fisch.	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings, 1701 m, N 37°13'682" – E 31°23'231", 19.v.2008, Çinbilgel 3397; İbradı, Çeşgar Plateau, on the rock, 1642 m, N 37°17'662" – E 31°23'989", 17.vi.2008, Çinbilgel 4195; Manavgat, Kocatop Mountain foothills, dwarf bush formation, 1836 m, N 37°12'052" – E 31°18'401", 11.vi.2009, Çinbilgel 5426, <i>End., E. Medit., (H)</i> .
	<i>Veronica hederifolia</i> L.	Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4643; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 980 m, N 37°10'766" – E 31°25'403", 05.iii.2011, Çinbilgel 7330; Manavgat, Eşekbeli Location, <i>Pinus nigra</i> forest (mixed with <i>Cedrus libani</i>), 1684 m, N 37°15'059" – E 31°19'785", 27.iii.2011, Çinbilgel 7524, <i>(Th)</i> .
Platanaceae		
	<i>Platanus orientalis</i> L.	İbradı, Suçikan Location way, streamsides, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2726; İbradı, Between Akpınar Location and Suçikan Location, streamsides, 1091 m, N 37°13'075" – E 31°24'456", 18.vi.2008, Çinbilgel 4433; Manavgat, Burmahan Neighbourhood, around of Erenli Location, <i>Pinus brutia</i> forest, 723 m, N 37°09'738" – E 31°16'983", 26.iii.2011, Çinbilgel 7462, <i>(Ph)</i> .
Plumbaginaceae		
	<i>Acantholimon acerosum</i> (Willd.) Boiss. subsp. <i>acerosum</i> var. <i>acerosum</i>	Manavgat, around of Çağlan Breach, moving rock and stony places, 1665 m, N 37°15'851" – E 31°21'000", 12.vi.2008, Çinbilgel 3929; Manavgat, Kızılpınar Plateau, thorny cushion formation, 1766 m, N 37°14'063" – E 31°20'070", 03.viii.2009, Çinbilgel 5609; İbradı, Çeşgar Plateau Breach, stony places, 1682 m, N 37°17'288" – E 31°24'324", 01.vii.2010, Çinbilgel 6907, <i>Ir. – Tur., (Ch)</i> .
	<i>Acantholimon ulicinum</i> (Willd. ex Schult.) Boiss. var. <i>ulicinum</i>	Manavgat, Çiğdem Hill, windy hills, 1962 m, N 37°15'366" – E 31°21'036", 11.vi.2008, Çinbilgel 3924; Manavgat, Kovacık Mountain (around of Uzunçaşır Location), rock, 1960 m, N 37°13'622" – E 31°20'149", 29.vi.2009, Çinbilgel 5511; İbradı, Melik Mountain (Çeşgar Hill), windy hills, 2263 m, N 37°16'085" – E 31°23'707", 18.vi.2010, Çinbilgel 6514, <i>E. Medit., (Ch)</i> .
	<i>Plumbago europaea</i> L.	İbradı, Toka Plateau way, Belova Location, stony places, 1519 m, N 37°13'719" – E 31°23'577", 02.vii.2010, Çinbilgel 7255, <i>Euro-Sib., (H)</i> .
Poaceae		
	<i>Aegilops biuncialis</i> Vis.	İbradı, Söbüova Location (around of Akpınar Location), roadsides, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3553; İbradı, between Akpınar Location and Suçikan Location, stony places, 1091 m, N 37° 13' 075" – E 031° 24' 456", 18.vi.2008, Çinbilgel 4428; İbradı, Boğazyurt Location, stony places, 974 m, N 37°10'338" – E 31°25'772", 19.v.2011, Çinbilgel 7583, <i>(Th)</i> .
	<i>Aegilops triuncialis</i> L. subsp. <i>triuncialis</i>	İbradı, Taşkesiği Location, <i>Pinus nigra</i> forest – roadsides, 1308 m, N 37°15'191" – E 31°24'495", 18.vi.2008, Çinbilgel 4367; Manavgat, Kızılpınar Plateau way, Kavaklı Location, <i>Juniperus excelsa</i> forest, 1374 m, N 37°14'135" – E 31°18'658", 07.vi.2011, Çinbilgel 7921, <i>(Th)</i> .
	<i>Aegilops umbellulata</i> Zhuk.	İbradı, Melik Plateau, streamsides, 1404 m, N 37°15'734" – E 31°24'641", 01.vii.2010, Çinbilgel 6741; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7706; Manavgat, around of Burmahan Neighbourhood (Öküztaşı Location), <i>Pinus brutia</i> forest, 683 m, N 37°10'293" – E 31°16'897", 06.vi.2011, Çinbilgel 7800, <i>Ir. – Tur., (Th)</i> .
	<i>Agrostis stolonifera</i> L.	Manavgat, around of Burmahan Neighbourhood (Değirmenderesi Location), damp places, 685 m, N 37°10'723" – E 31°16'850", 06.vi.2011, Çinbilgel 7787; Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 749 m, N 37°09'726" – E 31°17'001", 06.vi.2011, Çinbilgel 7813, <i>Euro-Sib., (H)</i> .
	<i>Aira elegantissima</i> Schur subsp. <i>elegantissima</i>	Manavgat, Kepez Plateau, damp places, 1636 m, N 37°10'971" – E 31°21'114", 03.vi.2009, Çinbilgel 5164; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1462 m, N 37°15'665" – E 31°24'474", 30.vi.2010, Çinbilgel 6605, <i>Medit., (Th)</i> .
	<i>Alopecurus arundinaceus</i> Poir.	İbradı, Toka Plateau, meadows, 1544 m, N 37°14'435" – E 31°22'650", 18.v.2008, Çinbilgel 3157; Manavgat, Sarıtaş Location surroundings, damp places, 1757 m, N 37°15'221" – E 31°20'452", 12.vi.2008, Çinbilgel 4011; İbradı, Toka Plateau, meadows, 1500 m, N 37°14'367" – E 31°22'605", 08.v.2010, Çinbilgel 6041, <i>Euro-Sib., (H)</i> .
	* <i>Alopecurus lanatus</i> Sibth. & Sm.	İbradı, Toka Plateau, moist stony places, 1500 m, N 37°14'332" – E 31°22'569", 08.v.2010, Çinbilgel 6051, <i>End., E. Medit., (H)</i> .
	<i>Alopecurus textilis</i> Boiss. subsp. <i>textilis</i>	Manavgat, Kaldırım Mountain, windy hills, 2065 m, N 37°10'436" – E 31°20'671", 03.vi.2009, Çinbilgel 5162; Manavgat, Kocatop Mountain (Koca Hill = Köygördüğü Location), windy hills, 2150 m, N 37°12'346" – E 31°18'633", 10.vi.2009, Çinbilgel 5343, <i>Ir. – Tur., (H)</i> .

	<i>Alopecurus vaginatus</i> (Willd.) Kunth	İbradı, hillsides of Melik Mountain, stony slopes, 1955 m, N 37°15'593" – E 31°23'351", 18.v.2008, Çinbilgel 3301; İbradı, Melik Mountain (Orta Hill), windy hills, 2187 m, N 37°16'210" – E 31°22'985", 17.vi.2010, Çinbilgel 6387; İbradı, Toka Plateau Breach, meadows, 1647 m, N 37°13'777" – E 31°23'191", 21.v.2011, Çinbilgel 7661, (H).
	<i>Apera intermedia</i> Hack.	Manavgat, Kızılpinar Plateau way, Akin – Karahan Location, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çinbilgel 4147; İbradı, Aşağıtoka Plateau, stony places, 1515 m, N 37°13'851" – E 31°22'437", 16.vi.2010, Çinbilgel 6269; Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 749 m, N 37°09'726" – E 31°17'001", 06.vi.2011, Çinbilgel 7813, Ir. – Tur., (Th).
	<i>Arrhenatherum elatius</i> (L.) P.Beauv. ex J.Presl & C.Presl subsp. <i>elatius</i>	İbradı, Aşağıtoka Plateau, debris, 1547 m, N 37°13'903" – E 31°22'326", 19.v.2008, Çinbilgel 3370, Euro-Sib., (H).
	<i>Arundo donax</i> L.	Manavgat, Between Burmahan Neighbourhood and Çay Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°11'284" – E 31°16'291", 11.x.2009, Çinbilgel 5795, (H).
	<i>Avena sterilis</i> L. subsp. <i>ludoviciana</i> (Durieu) Gillet & Magne	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7733; Manavgat, Çay Neighbourhood surroundings (Dikmen Hill), <i>Pinus brutia</i> forest, 816 m, N 37°12'492" – E 31°16'225", 07.vi.2011, Çinbilgel 7895; Manavgat, Çay Neighbourhood, roadsides, 690 m, N 37°12'122" – E 31°16'461", 07.vi.2011, Çinbilgel 7934, (Th).
	<i>Avena sterilis</i> L. subsp. <i>sterilis</i>	Manavgat, Kızılpinar Plateau way, between Akin – Karahan Locations, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çinbilgel 4151, (Th).
	<i>Brachypodium distachyon</i> (L.) P.Beauv.	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7716; Manavgat, Çay Neighbourhood surroundings (Dikmen Hill), <i>Pinus brutia</i> forest, 816 m, N 37°12'492" – E 31°16'225", 07.vi.2011, Çinbilgel 7889, 4911/a, Medit., (Th).
	<i>Brachypodium sylvaticum</i> (Huds.) P.Beauv.	İbradı, Gevenli Location, streamsides, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3519; İbradı, Suçikan Location surroundings, streamsides, 1257 m, N 37°14'782" – E 31°24'366", 02.vii.2010, Çinbilgel 7046; İbradı, Suçikan Location, streamsides, 1252 m, N 37°14'487" – E 31°24'182", 02.vii.2010, Çinbilgel 7102, Euro-Sib., (H).
	<i>Briza humilis</i> M.Bieb.	İbradı, Gevenli Location, under the <i>Pinus brutia</i> forest, 1108 m, N 37°12'316" – E 31°24'613", 17.v.2008, Çinbilgel 3013; Manavgat, Yelibelen Location, <i>Pinus brutia</i> forest, 952 m, N 37°09'197" – E 31°17'364", 06.vi.2011, Çinbilgel 7858, (Th).
	<i>Bromus diandrus</i> Roth	İbradı, Söbüova Location (around of Akpınar Location), stony places, 1057 m, N 37°12'262" – E 31°24'792", 19.v.2008, Çinbilgel 3562, (Th).
	<i>Bromus japonicus</i> Thunb. subsp. <i>japonicus</i>	Manavgat, Kızılpinar Plateau, mixed <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1718 m, N 37°14'103" – E 31°19'803", 09.vi.2009, Çinbilgel 5242; İbradı, Çeşgar Plateau, dwarf bush formation, 1633 m, N 37°17'430" – E 31°23'654", 01.vii.2010, Çinbilgel 6874; İbradı, Aşağıtoka Plateau, rocky hills, 1529 m, N 37°12'869" – E 31°22'570", 02.vii.2011, Çinbilgel 7983, (Th)
	<i>Bromus scoparius</i> L.	Manavgat, Kızılpinar Plateau, under the <i>Cedrus libani</i> forest, 1717 m, N 37°14'110" – E 31°19'899", 09.vi.2009, Çinbilgel 5239; İbradı, Toka Plateau, meadows, 1500 m, N 37°14'367" – E 31°22'605", 08.v.2010, Çinbilgel 6036; İbradı, Aşağıtoka Plateau, meadows, 1514 m, N 37°13'772" – E 31°22'587", 08.vi.2011, Çinbilgel 7956, (Th).
	<i>Bromus sterilis</i> L.	Manavgat, Burunçalı Location, mixed <i>Cedrus libani</i> forest, 1567 m, N 37°08'813" – E 31°19'104", 02.vi.2009, Çinbilgel 4998; İbradı, Melik Plateau, screes, 1491 m, N 37°15'544" – E 31°24'739", 02.vii.2010, Çinbilgel 6989; Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 680 m, N 37°10'071" – E 31°16'971", 06.vi.2011, Çinbilgel 7828, (Th).
	<i>Bromus tectorum</i> L.	Manavgat, Akpınar Plateau, stony places, 1403 m, N 37°09'414" – E 31°20'759", 19.v.2009, Çinbilgel 4753; İbradı, Toka Plateau Breach, stony slopes, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5968; İbradı, Cevizlidere Location, stony places, 1200 m, N 37°13'892" – E 31°24'189", 02.vii.2010, Çinbilgel 7135, (Th).
	<i>Bromus tomentellus</i> Boiss. subsp. <i>tomentellus</i>	Manavgat, Kovacık Mountain foothills (Gökçebel Location), dwarf bush-thorny cushion formation, 1725 m, N 37°13'999" – E 31°19'407", 09.vi.2009, Çinbilgel 5214; İbradı, Toka Plateau, stony slopes, 1520 m, N 37°14'038" – E 31°22'783", 09.v.2010, Çinbilgel 6086; İbradı, Aşağıtoka Plateau, meadows, 1523 m, N 37°14'183" – E 31°22'738", 20.v.2011, Çinbilgel 7637, Ir. – Tur., (H).
	<i>Catapodium rigidum</i> (L.) C.E.Hubb. subsp. <i>rigidum</i> var. <i>rigidum</i>	İbradı, Cevizlidere Location, stony places, 1149 m, N 37°13'475" – E 31°24'190", 02.vii.2010, Çinbilgel 7141, (Th).
	<i>Chrysopogon gryllus</i> (L.) Trin. subsp. <i>gryllus</i>	Manavgat, between Çay and Burmahan Neighbourhoods, <i>Pinus brutia</i> forest, 700 m, N 37°11'519" – E 31°16'218", 06.vi.2011, Çinbilgel 7777; İbradı, Eynif Plain, arid meadows, 947 m, N 37°07'654" – E 31°27'562", 02.vii.2011, Çinbilgel 7968, (H).

	<i>Cynodon dactylon</i> (L.) Pers. var. <i>villosus</i> Regel	İbradı, Melik Plateau, stony places, 1387 m, N 37°15'533" – E 31°24'528", 30.vi.2010, Çinbilgel 6732, (H).
	<i>Cynosurus echinatus</i> L.	İbradı, Melik Plateau, forest clearings – stony places, 1406 m, N 37°15'466" – E 31°24'469", 30.vi.2010, Çinbilgel 6675; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 990 m, N 37°10'778" – E 31°25'386", 21.v.2011, Çinbilgel 7697; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 983 m, N 37°10'973" – E 31°25'293", 03.vii.2011, Çinbilgel 8047, Medit., (Th).
	<i>Dactylis glomerata</i> L. subsp. <i>hispanica</i> (Roth) Nyman	Manavgat, Akpınar Plateau (Kelsu Location), under the forest, 1591 m, N 37°09'534" – E 31°20'305", 01.vi.2009, Çinbilgel 4843; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1462 m, N 37°15'665" – E 31°24'474", 30.vi.2010, Çinbilgel 6611; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 622 m, N 37°11'707" – E 31°16'064", 06.vi.2011, Çinbilgel 7749, (H).
	<i>Echinaria capitata</i> (L.) Desf.	Manavgat, Düğnük Plateau, stony places, 1165 m, N 37°13'430" – E 31°17'747", 13.vi.2008, Çinbilgel 4071, (Th)
	<i>Elymus hispidus</i> (Opiz) Melderis subsp. <i>podperae</i> (Nábelek) Melderis	İbradı, Melik Mountain (Hasan Hill), stony slopes, 1833 m, N 37°16'969" – E 31°23'331", 18.vi.2008, Çinbilgel 4277; İbradı, Melik Plateau, forest clearings – stony places, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6565, Ir. – Tur., (H).
	<i>Elymus panormitanus</i> (Parl.) Tzvelev	Manavgat, Çiğdem Hill, rocky stony hillsides, 2015 m, N 37°15'429" – E 31°21'200", 11.vi.2008, Çinbilgel 3921; İbradı, Yukarıtoka Plateau, stony places, 1879 m, N 37°15'975" – E 31°22'558", 17.vi.2010, Çinbilgel 6290; İbradı, Melik Plateau surroundings (towards Suçikan Location), <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7028, Medit., (H).
	<i>Eremopoa capillaris</i> R.R.Mill.	Manavgat, Sürüçkan Mountain, stony places, 2107 m, N 37°12'335" – E 31°19'690", 30.vi.2009, Çinbilgel 5560; İbradı, Melik Mountain (Orta Hill), dwarf bush formation, 2153 m, N 37°16'186" – E 31°22'903", 17.vi.2010, Çinbilgel 6333; Manavgat, around of Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest, 702 m, N 37°09'905" – E 31°16'998", 06.vi.2011, Çinbilgel 7842, E. Medit., (Th).
	<i>Festuca jeanpertii</i> (St.–Yves) F.Markgr. subsp. <i>jeanpertii</i>	İbradı, Toka Plateau (Göçebe Location), rocky places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çinbilgel 3121; Manavgat, around of Derme, dwarf bush formation, 2223 m, N 37°12'479" – E 31°19'531", 30.vi.2009, Çinbilgel 5593; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 2083 m, N 37°16'553" – E 31°22'889", 17.vi.2010, Çinbilgel 6442, E. Medit., (H).
	<i>Festuca valesiaca</i> Schleich. ex Gaudin	İbradı, Toka Plateau (Göçebe Location), rocky places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çinbilgel 3118; Manavgat, Burunçalı Location, mixed <i>Cedrus libani</i> forest, 1567 m, N 37°08'813" – E 31°19'104", 02.vi.2009, Çinbilgel 4986; İbradı, Melik Plateau, dwarf bush formation, 1502 m, N 37°16'195" – E 31°24'573", 01.vii.2010, Çinbilgel 6936, (H).
	<i>Hordeum bulbosum</i> L.	İbradı, Yukarıtoka Plateau (between Beydiğın Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3034; Manavgat, Kepez Plateau, 1636 m, N 37°10'971" – E 31°21'114", 03.vi.2009, Çinbilgel 5171; İbradı, Toka Plateau way, <i>Pinus nigra</i> forest, 1662 m, N 37°13'925" – E 31°22'981", 08.v.2010, Çinbilgel 6008, (H).
	<i>Hordeum murinum</i> L. subsp. <i>glaucum</i> (Steud.) Tzvelev	İbradı, Çeşgar Plateau, stony places, 1588 m, N 37°17'747" – E 31°23'791", 17.vi.2008, Çinbilgel 4198; Manavgat, Akpınar Plateau, stony places, 1403 m, N 37°09'414" – E 31°20'759", 19.v.2009, Çinbilgel 4752; İbradı, Melik Plateau, stony places, 1452 m, N 37°15'960" – E 31°24'578", 01.vii.2010, Çinbilgel 6751, (Th)
	<i>Hordeum murinum</i> L. subsp. <i>murinum</i>	Manavgat, Çağlan Breach, stony places, 1641 m, N 37°15'823" – E 31°20'867", 12.vi.2008, Çinbilgel 4001; Manavgat, Kızılınar Plateau, rocky stony places, 1720 m, N 37°14'073" – E 31°19'695", 09.vi.2009, Çinbilgel 5277; İbradı, Toka Plateau (Beydiğın Plateau), stony places, 1518 m, N 37°14'403" – E 31°22'504", 03.vii.2011, Çinbilgel 7993, (Th)
	<i>Hyparrhenia hirta</i> (L.) Stapf	İbradı, around of Çeşgar Plateau Breach, stony places, 1641 m, N 37°17'441" – E 31°24'212", 01.vii.2010, Çinbilgel 6817, (H).
	<i>Koeleria pyramidata</i> (Lam.) P.Beauv.	Manavgat, Dermebeli Location (Aritaşı), on the rock and stony places, 1874 m, N 37°13'519" – E 31°20'317", 10.vi.2008, Çinbilgel 3824; Manavgat, Burunçalı Location, mixed <i>Cedrus libani</i> forest, 1567 m, N 37°08'813" – E 31°19'104", 02.vi.2009, Çinbilgel 4989; İbradı, Melik Mountain (Çeşgar Hill), windy hills, 2260 m, N 37°16'093" – E 31°23'712", 18.vi.2010, Çinbilgel 6510, Euro-Sib., (H).
	<i>Lolium rigidum</i> Gaudich. var. <i>rigidum</i>	Manavgat, Kızılınar Plateau way, between Akin – Karahan Locations, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çinbilgel 4150; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7172; Manavgat, around of Burmahan Neighbourhood, roadsides, 700 m, N 37°10'623" – E 31°16'847", 06.vi.2011, Çinbilgel 7790, (Th).
	<i>Melica ciliata</i> L. subsp. <i>ciliata</i>	İbradı, Suçikan Location way, stony rocky places, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4421; İbradı, Melik Plateau, forest clearings – stony places, 1406 m, N 37°15'466" – E 31°24'469", 30.vi.2010, Çinbilgel 6681; İbradı, Suçikan Location, streamsides, 1252 m, N 37°14'487" – E 31°24'182", 02.vii.2010, Çinbilgel 7098, (H).

	<i>Melica minuta</i> L.	Manavgat, Kovacık Mountain surroundings, stony – rocky slopes, 1807 m, N 37°13'996" – E 31°20'274", 09.vi.2008, Çinbilgel 3658; Manavgat, Kovacık Mountain surroundings, on the rock, 1887 m, N 37°14'043" – E 31°20'377", 09.vi.2008, Çinbilgel 3698; Manavgat, between Dermebeli Location and Sürüçikan Mountain, stony slopes, 1914 m, N 37°13'471" – E 31°20'139", 10.vi.2008, Çinbilgel 3734, <i>Medit.</i> , (H).
	<i>Melica persica</i> Kunth. subsp. <i>inaequiglumis</i> (Boiss.) Bor	İbradı, Çeşgar Plateau, stony slopes, 1572 m, N 37°17'675" – E 31°23'744", 17.vi.2008, Çinbilgel 4205; İbradı, Çeşgar Plateau Breach, stony places, 1682 m, N 37°17'288" – E 31°24'324", 01.vii.2010, Çinbilgel 6909; İbradı, Melik Plateau, dwarf bush formation, 1665 m, N 37°17'156" – E 31°24'327", 01.vii.2010, Çinbilgel 6924, (H).
	<i>Milium vernale</i> M.Bieb. subsp. <i>montianum</i> (Parl.) Trab. ex Emb. & Maire	İbradı, Toka Plateau, <i>Pinus nigra</i> forest clearings, 1544 m, N 37°14'435" – E 31°22'650", 18.v.2008, Çinbilgel 3169; İbradı, around of Yirme Hill, <i>Pinus nigra</i> forest, 1550 m, N 37°14'328" – E 31°22'786", 08.v.2010, Çinbilgel 6022; İbradı, Suçikan Location, streamsides, 1259 m, N 37°14'503" – E 31°24'173", 02.vii.2010, Çinbilgel 7069, <i>Medit.</i> , (Th).
	<i>Narduroides salzmannii</i> (Boiss.) Rouy	<i>Medit.</i> , (Th), (Observation).
	<i>Oryzopsis holciformis</i> (M.Bieb.) Hack. var. <i>holciformis</i>	İbradı, Suçikan Location way, under the forest, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4417; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1502 m, N 37°15'548" – E 31°24'365", 30.vi.2010, Çinbilgel 6654; İbradı, around of Çukuryurt Plateau, <i>Pinus nigra</i> forest, 1227 m, N 37°12'049" – E 31°24'169", 03.vii.2011, Çinbilgel 8004, (H).
	<i>Oryzopsis miliacea</i> (L.) Asch. & Schweinf. subsp. <i>thomasi</i> (Duby) K.Richt.	Manavgat, Çay Neighbourhood surroundings (Değirmenderesi Bridge), streamsides, 705 m, N 37°12'138" – E 31°16'452", 07.vi.2011, Çinbilgel 7883, (H).
	<i>Phleum montanum</i> K.Koch subsp. <i>serrulatum</i> (Boiss.) Doğan	Manavgat, Derme Plateau (Babacık Plateau Location), dwarf bush formation, 1784 m, N 37°12'727" – E 31°20'180", 30.vi.2009, Çinbilgel 5551, <i>E. Medit.</i> , (H).
	<i>Pheum pratense</i> L.	Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 703 m, N 37°10'159" – E 31°17'019", 06.vi.2011, Çinbilgel 7821, Euro-Sib., (H).
	<i>Phleum subulatum</i> (Savi) Asch. & Graebn. subsp. <i>subulatum</i>	Manavgat, Arğıça Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4129; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çinbilgel 6714; Manavgat, around of Burmahan Neighbourhood (Payır Location), <i>Pinus brutia</i> forest, 803 m, N 37°09'640" – E 31°17'067", 06.vi.2011, Çinbilgel 7849, (Th).
	<i>Poa angustifolia</i> L.	İbradı, southern hillsides of Yaylacık Mountain, stony places, 1541 m, N 37°10'262" – E 31°24'049", 18.iv.2009, Çinbilgel 4576; Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), rocky places, 1734 m, N 37°13'822" – E 31°19'204", 09.vi.2009, Çinbilgel 5200; İbradı, Çukuryurt Plateau surroundings, <i>Pinus nigra</i> forest, 1357 m, N 37°12'461" – E 31°23'914", 09.v.2010, Çinbilgel 6160, (H).
	<i>Poa annua</i> L.	İbradı, Suçikan Location way, streamsides, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2787; Manavgat, Kepez Plateau, wet meadows, 1624 m, N 37°11'085" – E 31°21'009", 19.v.2009, Çinbilgel 4793; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7742, (Th).
	<i>Poa bulbosa</i> L.	İbradı, between Güzle Plateau – Yaylacık Mountain, moist places, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çinbilgel 2892; Manavgat, Kepezbeli Location, stony places, 1644 m, N 37°10'331" – E 31°21'666", 19.v.2009, Çinbilgel 4739; İbradı, Toka Plateau Breach, stony slopes, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5980, (H).
	<i>Poa nemoralis</i> L.	Manavgat, Dutluca Plateau, under the rock – stony places, 1282 m, N 37°13'785" – E 31°18'213", 13.vi.2008, Çinbilgel 4062; İbradı, Çeşgar Plateau surroundings, rock, 1628 m, N 37°17'545" – E 31°24'098", 01.vii.2010, Çinbilgel 6904, (H).
	<i>Poa pratensis</i> L.	Manavgat, Kepez Plateau, meadows, 1628 m, N 37°11'110" – E 31°21'021", 03.vi.2009, Çinbilgel 5179, (H).
	<i>Poa timoleontis</i> Heldr. ex Boiss.	Manavgat, around of Sartaş Location, stony hillsides, 2080 m, N 37°14'845" – E 31°20'924", 11.vi.2008, Çinbilgel 3902; Manavgat, Kepez Plateau, stony places, 1636 m, N 37°10'971" – E 31°21'114", 03.vi.2009, Çinbilgel 5166; İbradı, Toka Plateau, stony places, 1538 m, N 37°14'038" – E 31°22'783", 09.v.2010, Çinbilgel 6081, <i>E. Medit.</i> , (H).
	<i>Poa trivialis</i> L.	Manavgat, Eşekbeli Location, <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1750 m, N 37°15'074" – E 31°20'047", 12.vi.2008, Çinbilgel 4017; Manavgat, Kepez Plateau (İnçayırı Location), damp places, 1638 m, N 37°11'549" – E 31°20'684", 11.x.2009, Çinbilgel 5855; Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 680 m, N 37°10'071" – E 31°16'971", 06.vi.2011, Çinbilgel 7826, (H).
	<i>Polypogon viridis</i> (Gouan) Breistr.	Manavgat, Kızılpinar Plateau way, between Akin – Karahan Locations, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çinbilgel 4147, Euro-Sib., (H).

	<i>Sesleria alba</i> Sm.	İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7186, (H).
	<i>Stipa arabica</i> Trin. & Rupr.	İbradı, Melik Mountain (the eastern slopes of Hasan Hill), stony slopes, 1833 m, N 37°16'969" – E 31°23'331", 18.vi.2008, Çinbilgel 4272, Ir. – Tur., (H).
	<i>Stipa bromoides</i> (L.) Dörf.	İbradı, Suçikan Location way, under the forest – stony rocky places, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4422; İbradı, Melik Plateau, stony places, 1486 m, N 37°15'604" – E 31°24'398", 30.vi.2010, Çinbilgel 6596; İbradı, Tahtalı Location, <i>Pinus nigra</i> forest, 1275 m, N 37°12'721" – E 31°24'600", 02.vii.2010, Çinbilgel 7121, Medit., (H).
	<i>Stipa ehrenbergiana</i> Trin. & Rupr.	İbradı, Toka Plateau, around of Erenler Hill, stony places, 1574 m, N 37°15'010" – E 31°22'335", 17.v.2008, Çinbilgel 3151; İbradı, Melik Plateau, stony slopes, 1644 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4168; Manavgat, Derme Plateau (Babacık Plateau Location), dwarf bush formation, 1784 m, N 37°12'727" – E 31°20'180", 30.vi.2009, Çinbilgel 5549, Ir. – Tur., (H).
	<i>Stipa holosericea</i> Trin.	Manavgat, Çiğdem Hill, windy hills, 1962 m, N 37°15'366" – E 31°21'036", 11.vi.2008, Çinbilgel 3925; İbradı, Toka Plateau Breach, stony slopes, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5978; İbradı, Aşağıtoka Plateau, thorny cushion formation, 1523 m, N 37°12'927" – E 31°22'506", 16.vi.2010, Çinbilgel 6201, Ir. – Tur., (H).
	<i>Stipa lessingiana</i> Trin. & Rupr.	İbradı, Aşağıtoka Plateau, stony places, 1530 m, N 37°12'883" – E 31°22'500", 16.vi.2010, Çinbilgel 6211; İbradı, Melik Plateau, under the forest, 1469 m, N 37°15'618" – E 31°24'438", 30.vi.2010, Çinbilgel 6585, (H).
	<i>Stipa pulcherrima</i> K.Koch subsp. <i>epilosa</i> (Martinovsky) Tzvelev	İbradı, Aşağıtoka Plateau, debris, 1547 m, N 37°13'903" – E 31°22'326", 19.v.2008, Çinbilgel 3369, (H).
	<i>Triticum aestivum</i> L.	Manavgat, Arğıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4137, (Th).
	<i>Zea mays</i> L. subsp. <i>mays</i>	Manavgat, Derme Plateau, gardens, 1803 m, N 37°13'054" – E 31°20'219", 30.vi.2009, Çinbilgel 5571; Manavgat, Derme Plateau, crop fields, 1699 m, N 37°12'911" – E 31°20'427", 05.viii.2009, Çinbilgel 5711, (Th).
	<i>Zingiber pispidica</i> (Boiss.) Tutin	Manavgat, Kepez Plateau, meadows, 1628 m, N 37°11'110" – E 31°21'021", 03.vi.2009, Çinbilgel 5178; İbradı, Toka Plateau, 1500 m, N 37°14'367" – E 31°22'605", 08.v.2010, Çinbilgel 6042; İbradı, Melik Plateau, forest clearings – stony places, 1406 m, N 37°15'466" – E 31°24'469", 30.vi.2010, Çinbilgel 6676, Ir. – Tur., (Th).
Polygalaceae		
	* <i>Polygala pruinosa</i> Boiss. subsp. <i>megaptera</i> Cullen	İbradı, Melik Mountain (Hasan Hill), stony slopes, 2136 m, N 37°16'753" – E 31°23'122", 18.vi.2008, Çinbilgel 4335; Manavgat, Kocatop Mountain (Koca Hill = Köygöründüğü Location), windy hills, 2150 m, N 37°12'346" – E 31°18'633", 10.vi.2009, Çinbilgel 5344; İbradı, Melik Mountain (Orta Hill), rocky places, 1967 m, N 37°16'123" – E 31°22'618", 17.vi.2010, Çinbilgel 6312, End., (H).
	<i>Polygala pruinosa</i> Boiss. subsp. <i>pruinosa</i>	Manavgat, Kızılıpınar Plateau, dwarf bush, 1747 m, N 37°14'087" – E 31°19'983", 29.vi.2009, Çinbilgel 5502, (H).
Polygonaceae		
	<i>Atraphaxis billardieri</i> Jaub. & Spach var. <i>billardieri</i>	İbradı, Çeşgar Plateau, on the rock, 1620 m, N 37°17'210" – E 31°24'327", 17.vi.2008, Çinbilgel 4252; Manavgat, Sürüçikan Mountain, dwarf bush-thorny cushion formation, 2107 m, N 37°12'335" – E 31°19'690", 30.vi.2009, Çinbilgel 5555, Ir. – Tur., (Ch).
	<i>Polygonum arenastrum</i> Boreau	Manavgat, Derme Plateau (Zahmedin Plateau Location), 1761 m, N 37°12'348" – E 31°20'456", 20.vii.2008, Çinbilgel 4521; Manavgat, Kepez Plateau (İncayırı Location), stony places, 1624 m, N 37°11'525" – E 31°20'883", 11.x.2009, Çinbilgel 5838; İbradı, Melik Plateau, stony places, 1536 m, N 37°16'555" – E 31°24'517", 05.x.2010, Çinbilgel 7302, (Th).
	<i>Polygonum aviculare</i> L.	Manavgat, Eşekbeli Location, <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1750 m, N 37°15'074" – E 31°20'047", 12.vi.2008, Çinbilgel 4019; Manavgat, Derme Plateau (Zahmedin Plateau Location), stony places, 1761 m, N 37°12'348" – E 31°20'456", 20.vii.2008, Çinbilgel 4522; İbradı, Cevizlidere (around of Suçikan Location), streamsides, 1140 m, N 37°13'377" – E 31°24'211", 05.x.2010, Çinbilgel 7277, (Th).
	<i>Polygonum bellardii</i> All.	Manavgat, Kepez Plateau, meadows, 1631 m, N 37°11'208" – E 31°21'007", 11.x.2009, Çinbilgel 5843; İbradı, Melik Plateau, stony paths, 1502 m, N 37°16'157" – E 31°24'543", 01.vii.2010, Çinbilgel 6782; İbradı, Melik Plateau, forest clearings, 1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6957, (Th).
	<i>Polygonum cognatum</i> Meissn.	İbradı, Toka Plateau, stony places, 1528 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3321; Manavgat, Kovacık Mountain foothills (Hebil Plateau), arid meadows, 1766 m, N 37°13'788" – E 31°19'208", 09.vi.2009, Çinbilgel 5206; İbradı, Yukarıtoka Plateau, dwarf bush-thorny cushion formation, 1615 m, N 37°15'329" – E 31°22'423", 18.vi.2010, Çinbilgel 6541, (H).
	* <i>Polygonum karacae</i> Ziel. & Borat.	Manavgat, Kızılıpınar Plateau, dwarf bush-thorny cushion formation, 1757 m, N 37° 14'073" – E 031°19'973", 10.x.2009, Çinbilgel 5771; İbradı, Melik

		Mountain foothills, dwarf bush-thorny cushion formation, 1599 m, N 37°15'290" – E 31°22'419", 18.vi.2010, Çinbilgel 6471; İbradı, Aşağıtoka Plateau, stony places, 1511 m, N 37°13'446" – E 31°22'387", 02.vii.2010, Çinbilgel 7265, End., E. Medit. (mt.), (H).
	<i>Polygonum salebrosum</i> Coode & Cullen	Manavgat, Derme Plateau, meadows, 1805 m, N 37°13'179" – E 31°20'211", 10.x.2009, Çinbilgel 5781; İbradı, Melik Plateau, meadows, 1374 m, N 37°15'565" – E 31°24'628", 05.x.2010, Çinbilgel 7279; İbradı, Melik Plateau, stony places, 1443 m, N 37°15'917" – E 31°24'576", 05.x.2010, Çinbilgel 7293, Medit., (H).
	<i>Rumex acetosella</i> L.	İbradı, Toka Plateau Breach, stony places, 1693 m, N 37°13'752" – E 31°23'233", 08.v.2010, Çinbilgel 5983; İbradı, around of Çeşgar Plateau Breach, stony places, 1672 m, N 37°17'262" – E 31°24'324", 01.vii.2010, Çinbilgel 6809; İbradı, Aşağıtoka Plateau, moist stony places, 1534 m, N 37°12'906" – E 31°22'673", 02.vii.2011, Çinbilgel 7988, (Th).
	<i>Rumex crispus</i> L.	İbradı, Güzle Plateau, forest, 1293 m, N 37°10'330" – E 31°23'601", 20.iv.2008, Çinbilgel 2877, (H).
	<i>Rumex pulcher</i> L. subsp. <i>divaricatus</i> (L.) Murbeck	İbradı, around of Suçikan Location (Tahtalı Location), roadsides, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4382; İbradı, Çeşgar Plateau, dwarf bush-thorny cushion formation, 1580 m, N 37°17'516" – E 31°23'880", 01.vii.2010, Çinbilgel 6882; İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1223 m, N 37°14'468" – E 31°24'259", 02.vii.2010, Çinbilgel 7057, (H).
	<i>Rumex scutatus</i> L.	Manavgat, around of Çağlan Breach, stony places, 1654 m, N 37°15'995" – E 31°21'198", 12.vi.2008, Çinbilgel 3935; Manavgat, Kovacık Mountain foothills (Hebil Plateau oluşu), stony places, 1677 m, N 37°13'893" – E 31°19'274", 09.vi.2009, Çinbilgel 5216, (H).
	<i>Rumex tuberosus</i> L. subsp. <i>tuberosus</i>	İbradı, Yukarıtoka Plateau (Between Beydiğın Plateau and Göcebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3031; Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 749 m, N 37°09'726" – E 31°17'001", 06.vi.2011, Çinbilgel 7817, (H).
Primulaceae		
	<i>Anagallis arvensis</i> L. var. <i>arvensis</i>	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7740, (Th).
	<i>Androsace villosa</i> L.	Manavgat, Çandır Hill, stony hills, 2037 m, N 37°16'626" – E 31°21'029", 11.vi.2008, Çinbilgel 3883, Euro-Sib., (H).
	* <i>Cyclamen cilicicum</i> Boiss. & Heldr.	Manavgat, Burmahan Neighbourhood, Erenli Location, streamsides, 713 m, N 37°10'048" – E 31°16'974", 11.x.2009, Çinbilgel 5810; Manavgat, Akpınar Plateau, under the <i>Cedrus libani</i> forest, 1587 m, N 37°10'202" – E 31°21'505", 11.x.2009, Çinbilgel 5828; İbradı, Suçikan Location, under the forest, N 37°13'646" – E 31°24'123", 02.vii.2010, Çinbilgel 7247, End., E. Medit., (G).
	<i>Lysimachia atropurpurea</i> L.	Manavgat, around of Burmahan Neighbourhood, Erenli Location, streamsides, 703 m, N 37°10'159" – E 31°17'019", 06.vi.2011, Çinbilgel 7822, E. Medit., (Th).
Ranunculaceae		
	<i>Anemone blanda</i> Schott & Kotschy	İbradı, Güzle Plateau – Hillsides of Yaylacık Mountain, bozuk <i>Pinus nigra</i> forest and damp places, 1430 m, N 37°10'227" – E 31°23'913", 20.iv.2008, Çinbilgel 2881; İbradı, Aşağıtoka Plateau, moist – stony places, 1521 m, N 37°13'187" – E 31°22'678", 27.iii.2010, Çinbilgel 5865; İbradı, around of Suçikan Location, moist – stony places, 1160 m, N 37°14'079" – E 31°24'218", 05.iii.2011, Çinbilgel 7359, (G).
	<i>Clematis cirrhosa</i> L.	Manavgat, around of Burmahan Neighbourhood, streamsides, 693 m, N 37°11'151" – E 31°16'304", 06.vi.2011, Çinbilgel 7766, Medit., (Ph).
	<i>Clematis vitalba</i> L.	Manavgat, Burmahan Neighbourhood, Akpınar Plateau way, forest openings, 699 m, N 37°10'800" – E 31°16'736", 26.iii.2011, Çinbilgel 7420; Manavgat, Kızılpınar Plateau way, Putraklı Location, 734 m, N 37°12'249" – E 31°16'454", 27.iii.2011, Çinbilgel 7566, (Ph).
	<i>Eranthis hyemalis</i> (L.) Salisb.	İbradı, Suçikan Location way, damp places, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2770; Manavgat, between Kepez Plateau and hillsides of Kaldırım Mountain, slopes, 1657 m, N 37°10'485" – E 31°21'360", 03.v.2009, Çinbilgel 4711; İbradı, Aşağıtoka Plateau, moist – stony places, 1521 m, N 37°13'187" – E 31°22'678", 27.iii.2010, Çinbilgel 5867, (G).
	<i>Ranunculus arvensis</i> L.	Manavgat, Between Çay and Burmahan Neighbourhoods, streamsides, 706 m, N 37°11'423" – E 31°16'306", 06.vi.2011, Çinbilgel 7760; Manavgat, around of Burmahan Neighbourhood, stony roadsides, 704 m, N 37°11'401" – E 31°16'295", 06.vi.2011, Çinbilgel 7763, (Th).
	<i>Ranunculus cf. buhsei</i> Boiss.	Manavgat, Kaldırım Mountain (GökHill Location), 2065 m, N 37°10'750" – E 31°20'220", 03.vi.2009, Çinbilgel 5114, Hyrcano–Euksin (Euro-Sib.) (mt.), (H), (Det: Özçelik).
	<i>Ranunculus cadmicus</i> Boiss.	İbradı, around of Yirme Hill (towards Melik Mountain), stony places, 1875 m, N 37°14'859" – E 31°23'148", 18.v.2008, Çinbilgel 3282; İbradı, Gevenli Location, rock, 1145 m, N 37°11'947" – E 31°24'445", 28.iii.2010, Çinbilgel 5939, (H), (Det: Özçelik).
	<i>Ranunculus chius</i> DC.	Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7709, (Th).

	<i>Ranunculus cuneatus</i> Boiss.	İbradı, Yukarıtoka Plateau (Göçebe Location – Beydiğın Plateau), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çınbilgel 3034; İbradı, Toka Plateau way, <i>Pinus nigra</i> forest, 1662 m, N 37°13'925" – E 31°22'981", 08.v.2010, Çınbilgel 5996; Manavgat, Çay Neighbourhood, Eşekbeli Location, <i>Pinus nigra</i> forest, 1612 m, N 37°14'867" – E 31°19'864", 07.vi.2011, Çınbilgel 7904, (H).
	<i>Ranunculus damascenus</i> Boiss. & Gaill.	İbradı, Toka Plateau way, Belova Location, under the <i>Cedrus libani</i> forest – stony places, 1698 m, N 37°13'706" – E 31°23'257", 19.v.2008, Çınbilgel 3427; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest clearings, 1430 m, N 37°09'494" – E 31°20'838", 19.v.2009, Çınbilgel 4812; Manavgat, Kovacık Mountain foothills (Gökçebel Location), dwarf bush-thorny cushion formation, 1695 m, N 37°13'984" – E 31°19'318", 11.vi.2009, Çınbilgel 5438, Ir. – Tur., (H), (Conf. Özçelik).
	* <i>Ranunculus demissus</i> DC. var. <i>major</i> Boiss.	İbradı, Melik Mountain (Çeşgar Hill), windy stony slopes, 2100 m, N 37°15'610" – E 31°23'438", 18.v.2008, Çınbilgel 3315; Manavgat, Kocatop Mountain, snow patches, 2080 m, N 37°12'599" – E 31°18'884", 10.vi.2009, Çınbilgel 5279; İbradı, Toka Plateau Breach, meadows, 1676 m, N 37°13'774" – E 31°23'168", 21.v.2011, Çınbilgel 7668, End., (H), (Det: Özçelik).
	<i>Ranunculus ficaria</i> L. subsp. <i>calthifolius</i> (Rchb.) Arcang.	İbradı, Akpınar (Akmuar) Location, moist stony places, 1062 m, N 37°11'985" – E 31°24'815", 05.iii.2011, Çınbilgel 7370; Manavgat, Akpınar Plateau way, (Genger Cemetery), degraded <i>Pinus brutia</i> forest forest, 1077 m, N 37°09'026" – E 31°17'742", 26.iii.2011, Çınbilgel 7444; Manavgat, Çay Neighbourhood, Kızılıpınar Plateau way (Düğnük Plateau), <i>Pinus brutia</i> clearings – streamsides, 1172 m, N 37°13'352" – E 31°17'760", 27.iii.2011, Çınbilgel 7512, (H).
	<i>Ranunculus ficaria</i> L. subsp. <i>ficariiformis</i> Rouy & Foucaud	İbradı, Suçikan Location way, streamsides, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çınbilgel 2731; Manavgat, Akpınar Plateau way, <i>Pinus brutia</i> forest, 900 m, N 37°09'068" – E 31°17'826", 26.iii.2009, Çınbilgel 4554; İbradı, Çukuryurt Plateau, meadows, 1268 m, N 37°12'117" – E 31°24'023", 28.iii.2010, Çınbilgel 5923, (H).
	<i>Ranunculus ficarioides</i> Bory & Chaub.	Manavgat, Eşekbeli Location, <i>Pinus nigra</i> forest (mixed with <i>Cedrus libani</i>), 1684 m, N 37°15'059" – E 31°19'785", 27.iii.2011, Çınbilgel 7532, (H).
	<i>Ranunculus illyricus</i> L. subsp. <i>illyricus</i>	İbradı, Yukarıtoka Plateau (between Göçebe Location and Beydiğın Plateau), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çınbilgel 3092, (H), (Det: Özçelik).
	<i>Ranunculus marginatus</i> d'Urv.	İbradı, Akmuar Location, streamsides, 1032 m, N 37°11'838" – E 31°24'840", 19.viii.2008, Çınbilgel 3574, (Th).
	<i>Ranunculus muricatus</i> L.	İbradı, Boğazyurt Location, watersides, 1032 m, N 37°11'838" – E 31°24'840", 19.v.2008, Çınbilgel 3613, (Th).
	<i>Ranunculus oxyspermus</i> Willd.	İbradı, between Güzle Plateau – Yaylacık Mountain, damp places, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çınbilgel 2901, (H), (Conf. Özçelik).
	<i>Ranunculus polyanthemus</i> L.	Manavgat, around of Sarıtaş Location, damp places, 1757 m, N 37°15'221" – E 31°20'452", 12.vi.2008, Çınbilgel 4010; İbradı, Aşağıtoka Plateau, wet meadows, 1502 m, N 37°14'273" – E 31°22'580", 27.iii.2010, Çınbilgel 5873; İbradı, Aşağıtoka Plateau, wet meadows, 1520 m, N 37°14'220" – E 31°22'415", 20.v.2011, Çınbilgel 7643, (H), (Det: Özçelik).
	<i>Ranunculus sericeus</i> Banks & Sol.	Manavgat, Kepez Plateau, wet meadows, 1624 m, N 37°11'085" – E 31°21'009", 19.v.2009, Çınbilgel 4794, Ir. – Tur., (H).
	<i>Thalictrum orientale</i> Boiss.	İbradı, Suçikan Location way, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çınbilgel 2782; İbradı, Gevenli Location, rock 1143 m, N 37°12'008" – E 31°24'420", 28.iii.2010, Çınbilgel 5945; İbradı, Akmuar Location, rock, 1024 m, N 37°11'720" – E 31°24'844", 19.v.2011, Çınbilgel 7607, E. Medit., (H).
Rhamnaceae		
	<i>Paliurus spina-christi</i> P.Mill.	İbradı, Akmuar Location, Juniper Forest, 964 m, N 37°11'201" – E 31°25'141", 18.vi.2008, Çınbilgel 4446; Manavgat, Çay Neighbourhood, Kızılıpınar Plateau way, <i>Pinus brutia</i> forest, 628 m, N 37°11'815" – E 31°16'212", 10.x.2009, Çınbilgel 5745, (Ph).
	<i>Rhamnus lycioides</i> L. subsp. <i>graeca</i> (Boiss. & Reut.) Tutin	Manavgat, around of Kovacık Mountain (Kocayokuş Location), rock, 1887 m, N 37°14'043" – E 31°20'377", 09.vi.2008, Çınbilgel 3679; Manavgat, Kızılıpınar Plateau, rock, 1875 m, N 37°14'249" E 31°20'465", 03.viii.2009, Çınbilgel 5619; İbradı, Melik Mountain (Orta Hill), rocky places, 1967 m, N 37°16'123" – E 31°22'618", 17.vi.2010, Çınbilgel 6310, E. Medit., (Ph).
	* <i>Rhamnus thymifolia</i> Bornm.	Manavgat, Dermebeli Location (Aritaşı), rock, 1885 m, N 37°13'490" – E 31°20'297", 19.vii.2008, Çınbilgel 4464; Manavgat, Kaldırım Mountain, rocky places, 1820 m, N 37°09'912" – E 31°20'423", 01.vi.2009, Çınbilgel 4858, End., (Ph).
Rosaceae		
	<i>Agrimonia eupatoria</i> L. subsp. <i>asiatica</i> (Juz.) Skalicky	Manavgat, Çay Neighbourhood, openings – roadsides, 690 m, N 37°12'122" – E 31°16'461", 07.vi.2011, Çınbilgel 7941; İbradı, Tahtalı Location, <i>Pinus brutia</i> forest, 1066 m, N 37°12'606" – E 31°24'675", 03.vii.2011, Çınbilgel 8015, (H).
	* <i>Amelanchier parviflora</i> Boiss. var. <i>dentata</i> Browicz	İbradı, around of Yirme Hill, rocky hills, 1820 m, N 37°14'717" – E 31°23'178", 18.v.2008, Çınbilgel 3248; Manavgat, Derme Plateau (Yörtikyurdu Location), rocky places, 1843 m, N 37°11'506" – E 31°20'304", 20.vii.2008, Çınbilgel

		4507; around of Kocatop Mountain (Kürtağacıboğazı Location), 2006 m, N 37°12'817" – E 31°18'915", 10.vi.2009, Çınbilgel 5291, End., E. Medit., (Ph).
	<i>*Amelanchier parviflora</i> Boiss. var. <i>parviflora</i>	Manavgat, between Çandır Hill and Çiğdem Hill, stony slopes, 2024 m, N 37°14'656" – E 31°20'981", 11.vi.2008, Çınbilgel 3886, End., E. Medit., (Ph).
	<i>Cerasus prostrata</i> (Labill.) Ser. var. <i>prostrata</i>	İbradı, Yirme Hill, rocky places, 1820 m, N 37°14'717" – E 31°23'178", 18.v.2008, Çınbilgel 3233; Manavgat, Kaldırım Mountain (Aktünek Hill), windy hills, 2208 m, N 37°10'562" – E 31°20'327", 03.vi.2009, Çınbilgel 5105; İbradı, around of Çeşgar Plateau Breach, stony places, 1672 m, N 37°17'262" – E 31°24'324", 01.vii.2010, Çınbilgel 6806, Medit., (Ch).
	<i>Cotoneaster nummularius</i> Fisch. & C.A.Mey.	Manavgat, Kaldırım Mountain, slopes, 4943 m, N 37°09'635" – E 31°20'292", 01.vi.2009, Çınbilgel 4945; İbradı, Melik Mountain (Orta Hill), stony places, 2084 m, N 37°16'414" – E 31°23'077", 17.vi.2010, Çınbilgel 6404; İbradı, Melik Plateau, stony places, 1539 m, N 37°16'498" – E 31°24'484", 05.x.2010, Çınbilgel 7307, (Ph).
	<i>Crataegus monogyna</i> Jacq. var. <i>monogyna</i>	İbradı, between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çınbilgel 2940; İbradı, Akpınar Location, <i>Pinus nigra</i> forest clearings, 1047 m, N 37°11'563" – E 31°24'882", 02.vii.2010, Çınbilgel 7161; İbradı, Akmuar Location, streamsides, 1044 m, N 37°11'719" – E 31°24'820", 02.vii.2010, Çınbilgel 7216; Manavgat, Sartaş Location surroundings, stony hillsides, 2074 m, N 37°15'015" – E 31°20'984", 11.vi.2008, Çınbilgel 3909; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çınbilgel 7187; İbradı, around of Gevenli Location, <i>Pinus brutia</i> forest, 1092 m, N 37°12'670" – E 31°24'617", 02.vii.2010, Çınbilgel 7239, (Ph).
	<i>Crataegus orientalis</i> Pall. ex M.Bieb. subsp. <i>orientalis</i>	Manavgat, Dutluca Plateau, under the rock, stony places, 1282 m, N 37°13'785" – E 31°18'213", 13.vi.2008, Çınbilgel 4057; Manavgat, between Kovacık Mountain and Dermebeli Location, dwarf bush formation, 1916 m, N 37°13'542" – E 31°20'204", 30.vi.2009, Çınbilgel 5562; İbradı, Melik Plateau, sloping openings, 1448 m, N 37°15'952" – E 31°24'583", 05.x.2010, Çınbilgel 7296, (Ph).
	<i>Crataegus x sinaica</i>	İbradı, Eynif Plain, Yarbaşı Location, rocky places, 943 m, N 37°08'421" – E 31°27'636", 28.iii.2010, Çınbilgel 5965, Ir. – Tur., (Ph).
	<i>Geum heterocarpum</i> Boiss.	İbradı, around of Yirme Hill, stony places 1805 m, N 37°14'690" – E 31°23'119", 18.v.2008, Çınbilgel 3263; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 1933 m, N 37°16'534" – E 31°22'619", 17.vi.2010, Çınbilgel 6452, (H).
	<i>Geum urbanum</i> L.	İbradı, Çeşgar Plateau, stony slopes, 1659 m, N 37°17'402" – E 31°24'250", 17.vi.2008, Çınbilgel 4227; İbradı, Yukarıtoka Plateau, dwarf bush formation, 1635 m, N 37°15'828" – E 31°22'221", 17.vi.2010, Çınbilgel 6274; İbradı, Suçikan Location, streamsides, 1259 m, N 37°14'503" – E 31°24'173", 02.vii.2010, Çınbilgel 7071, Euro-Sib., (H).
	<i>Malus sylvestris</i> (L.) Mill.	İbradı, Suçikan Location, open places, 1219 m, N 37°14'431" – E 31°24'272", 02.vii.2010, Çınbilgel 7059, (Ph).
	<i>Potentilla kotschyana</i> Fenzl	Manavgat, southern hillsides of Kaldırım Mountain, under the <i>Cedrus libani</i> forest, 1551 m, N 37°09'461" – E 31°20'374", 01.vi.2009, Çınbilgel 4881; Manavgat, Kızılıpınar Plateau surroundings (Nurboğazı Location), rock, 1913 m, N 37°14'166" – E 31°20'469", 04.viii.2009, Çınbilgel 5673; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 2073 m, N 37°16'539" – E 31°23'096", 17.vi.2010, Çınbilgel 6431, E. Medit., (H).
	<i>Potentilla recta</i> L.	İbradı, Aşağıtoka Plateau, thorny cushion formation, 1526 m, N 37°12'919" – E 31°22'639", 16.vi.2010, Çınbilgel 6179; İbradı, Melik Plateau, stony places, 1406 m, N 37°15'507" – E 31°24'484", 30.vi.2010, Çınbilgel 6704; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çınbilgel 7170, (H).
	<i>Potentilla reptans</i> L.	Manavgat, Burunçalı Location, debris, 1857 m, N 37°09'063" – E 31°19'473", 02.vi.2009, Çınbilgel 5013; İbradı, Melik Plateau, damp places, 1440 m, N 37°15'862" – E 31°24'555", 01.vii.2010, Çınbilgel 6746; İbradı, Melik Plateau surroundings, streamsides, 1340 m, N 37°15'336" – E 31°24'546", 02.vii.2010, Çınbilgel 7024, (H).
	<i>Potentilla speciosa</i> Willd.	İbradı, Akmuar Location, rock, 1042 m, N 37°11'701" – E 31°24'851", 02.vii.2010, Çınbilgel 7226, (H).
	<i>Prunus divaricata</i> Ledeb. var. <i>divaricata</i>	İbradı, southeast slopes of Yaylacık Mountain, <i>Cedrus libani</i> forest clearings, 1263 m, N 37°09'958" – E 31°25'227", 19.iv.2009, Çınbilgel 4608; İbradı, Eynif Plain, Yarbaşı Location, rocky places, 943 m, N 37°08'421" – E 31°27'636", 28.iii.2010, Çınbilgel 5964; İbradı, Suçikan Location surroundings, roadsides, 1274 m, N 37°14'870" – E 31°24'365", 02.vii.2010, Çınbilgel 7044, (Ph).
	<i>Prunus x domestica</i> L.	İbradı, Çeşgar Plateau, rock, 1625 m, N 37°17'470" – E 31°23'695", 01.vii.2010, Çınbilgel 6844, (Ph).
	<i>Pyrus elaeagnifolia</i> Pall. subsp. <i>elaegnifolia</i>	Manavgat, Kızılıpınar Plateau, thorny cushion formation, 1766 m, N 37°14'063" – E 31°20'070", 03.viii.2009, Çınbilgel 5606; İbradı, Suçikan Location, <i>Pinus nigra</i> forest, 1223 m, N 37°14'468" – E 31°24'259", 02.vii.2010, Çınbilgel 7055; İbradı, Suçikan Location, forest clearings, N 37°13'646" – E 31°24'123", 02.vii.2010, Çınbilgel 7246, (Ph).

	<i>Pyrus syriaca</i> Boiss. var. <i>microphylla</i> Zoh. ex Browicz	İbradı, Akmuar Location, open places, 1033 m, N 37°12'323" – E 31°24'784", 18.vi.2008, Çinbilgel 4435; İbradı, Eynif Plain, Yarbaşı Location, rocky places, 943 m, N 37°08'421" – E 31°27'636", 28.iii.2010, Çinbilgel 5963, (Ph).
	<i>Pyrus syriaca</i> Boiss. var. <i>syriaca</i>	İbradı, Tahtalı Location (lower of Cevizlidere Location), <i>Pinus nigra</i> forest, 1097 m, N 37°12'817" – E 31°24'591", 02.vii.2010, Çinbilgel 7123, (Ph).
	<i>Rosa canina</i> L.	İbradı, between Güzle Plateau and Gavur Breach, <i>Pinus nigra</i> forest, 1252 m, N 37°09'556" – E 31°24'264", 20.iv.2008, Çinbilgel 2924; Manavgat, Akpınar Plateau (Doğruluk Location), <i>Cedrus libani</i> forest clearings, 1424 m, N 37°08'886" – E 31°20'013", 02.vi.2009, Çinbilgel 4974; İbradı, Cevizlidere Location, streamsides, 1163 m, N 37°13'696" – E 31°24'146", 02.vii.2010, Çinbilgel 7117, (Ph).
	<i>Rosa dumalis</i> Bechst. subsp. <i>antalyensis</i> (Manden.) Ö.Nilsson	İbradı, Tahtalı Location, streamsides, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4399, End., (Ph), (Det: Özçelik).
	<i>Rosa iberica</i> Stev.	Manavgat, Derme Plateau (Zahmedin Plateau Location), dwarf bush-thorny cushion formation, 1761 m, N 37°12'348" – E 31°20'456", 20.vii.2008, Çinbilgel 4526, (Ch), (Det: Özçelik).
	<i>Rosa pulverulenta</i> M.Bieb.	Manavgat, Kaldırım Mountain, screens, 1816 m, N 37°09'893" – E 31°20'398", 01.vi.2009, Çinbilgel 4957; Manavgat, Kızılpınar Plateau, dwarf bush-thorny cushion formation, 1754 m, N 37°14'090" – E 31°19'969", 04.viii.2009, Çinbilgel 5690; İbradı, Melik Plateau, dwarf bush-thorny cushion formation, 1398 m, N 37°15'665" – E 31°24'622", 05.x.2010, Çinbilgel 7287; (Syn. R. sicula Tratt.): Manavgat, Kızılpınar Plateau, dwarf bush-thorny cushion formation, 1782 m, N 37°14'186" – E 31°20'257", 03.viii.2009, Çinbilgel 5600; Manavgat, Kızılpınar Plateau, dwarf bush-thorny cushion formation, 1757 m, N 37°14'073" – E 31°19'973", 10.x.2009, Çinbilgel 5769, (Ch), (Conf. Özçelik).
	<i>Rubus canescens</i> DC. var. <i>canescens</i>	İbradı, Tahtalı Location, under the forest, 1206 m, N 37°14'386" – E 31°24'289", 18.vi.2008, Çinbilgel 4401; İbradı, Ekinlik Location surroundings, <i>Cedrus libani</i> forest (mixed with <i>Pinus nigra</i>), 1532 m, N 37°13'668" – E 31°23'510", 09.v.2010, Çinbilgel 6143; İbradı, Toka Plateau Breach, under the <i>Cedrus libani</i> forest, N 37°15'406" – E 31°22'669", 18.vi.2010, Çinbilgel 6549, Euro-Sib., (Ch).
	<i>Rubus sanctus</i> Schreb.	İbradı, Toka Plateau way, Belova Location, pathways, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3466; İbradı, Akmuar Location, streamsides, 1044 m, N 37°11'719" – E 31°24'820", 02.vii.2010, Çinbilgel 7217, (Ph).
	<i>Sanguisorba minor</i> L. subsp. <i>balearica</i> (Bourg. ex Nyman) Muñoz Garm. & C.Navarro	Manavgat, southern hillsides of Kaldırım Mountain, under the <i>Cedrus libani</i> forest, 1551 m, N 37°09'461" – E 31°20'374", 01.vi.2009, Çinbilgel 4886; İbradı, Toka Plateau, stony slopes, 1520 m, N 37°14'332" – E 31°22'709", 08.v.2010, Çinbilgel 6028; İbradı, Melik Plateau surroundings, <i>Pinus nigra</i> forest, 1347 m, N 37°15'332" – E 31°24'499", 02.vii.2010, Çinbilgel 7006, (H).
	<i>Sanguisorba verrucosa</i> (G.Don) Ces.	İbradı, Toka Plateau, stony places, 1528 m, N 37°14'496" – E 31°22'553", 19.v.2008, Çinbilgel 3324; Manavgat, Derme Plateau (around of Dermebeli Location), stony hillsides, 1827 m, N 37°13'386" – E 31°20'237", 10.vi.2008, Çinbilgel 3797; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7175, (H).
	<i>Sorbus umbellata</i> Fritsch	Manavgat, Kovacık Mountain surroundings, on the rock, 1887 m, N 37°14'043" – E 31°20'377", 09.vi.2008, Çinbilgel 3678; Manavgat, Kızılpınar Plateau, rock, 1846 m, N 37°14'402" – E 31°20'479", 03.viii.2009, Çinbilgel 5642; İbradı, Çeşgar Plateau, rock, 1621 m, N 37°17'565" – E 31°24'085", 01.vii.2010, Çinbilgel 6900, (Ph).
Rubiaceae		
	<i>Asperula arvensis</i> L.	İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest, 1462 m, N 37°13'685" – E 31°23'484", 19.v.2008, Çinbilgel 3479; Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4832; İbradı, Toka Plateau Breach surroundings, 1570 m, N 37°13'602" – E 31°23'417", 09.v.2010, Çinbilgel 6137, Medit., (Th).
	* <i>Asperula serotina</i> (Boiss. & Heldr.) Ehrend.	Manavgat, Kaldırım Mountain, rock, 1628 m, N 37°09'639" – E 31°20'286", 01.vi.2009, Çinbilgel 4851; Manavgat, Kalbur Gözenekleri Location, rock, 1788 m, N 37°13'259" – E 31°20'585", 05.viii.2009, Çinbilgel 5699; İbradı, Akmuar Location, rock, 1042 m, N 37°11'701" – E 31°24'851", 02.vii.2010, Çinbilgel 7228, End., E. Medit., (Ch).
	<i>Asperula setosa</i> Jaub. & Spach	İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3502; İbradı, Melik Plateau, forest clearings – stony places, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6570; İbradı, Gevenli Location, <i>Pinus brutia</i> forest clearings, 1071 m, N 37°12'300" – E 31°24'793", 03.vii.2011, Çinbilgel 8008, Ir. – Tur., (Th).
	* <i>Asperula stricta</i> Boiss. subsp. <i>elmaliensis</i> Schönb.–Tem.	İbradı, Melik Mountain (Hasan Hill), stony hillsides, 1761 m, N 37°16'962" – E 31°23'396", 18.vi.2008, Çinbilgel 4269, End., E. Medit., (H).
	* <i>Asperula stricta</i> Boiss. subsp. <i>latibracteata</i> (Boiss.) Ehrend.	Manavgat, Çandır Hill yamacı, dwarf bush formation, 1890 m, N 37°14'333" – E 31°20'924", 04.viii.2009, Çinbilgel 5677; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 2083 m, N 37°16'553" – E 31°22'889", 17.vi.2010,

		Çinbilgel 6437; İbradı, Çeşgar Plateau Breach, stony places, 1618 m, N 37°16'826" – E 31°24'504", 01.vii.2010, Çinbilgel 6794, End., Ir. – Tur., (H).
	* <i>Asperula stricta</i> Boiss. subsp. <i>monticola</i> Ehrend.	Manavgat, Sarıtaş Location, stony places, 2084 m, N 37°15'212" – E 31°20'984", 11.vi.2008, Çinbilgel 3907, End., E. Medit. (mt.), (H).
	<i>Crucianella latifolia</i> L.	İbradı, Gevenli Location, <i>Pinus nigra</i> forest, 1110 m, N 37°12'010" – E 31°24'417", 19.v.2008, Çinbilgel 3508; İbradı, Suçikan Location way, stony rocky places, 1095 m, N 37°13'791" – E 31°24'185", 18.vi.2008, Çinbilgel 4407; İbradı, Melik Plateau surroundings (towards Suçikan Location), <i>Pinus nigra</i> – <i>Cedrus libani</i> forest, 1289 m, N 37°14'953" – E 31°24'400", 02.vii.2010, Çinbilgel 7034, Medit., (Th).
	<i>Crucianella macrostachya</i> Boiss.	Manavgat, Argıçca Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4116; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1462 m, N 37°15'665" – E 31°24'474", 30.vi.2010, Çinbilgel 6613; Manavgat, around of Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest, 702 m, N 37°09'905" – E 31°16'998", 06.vi.2011, Çinbilgel 7840, E. Medit., (Th).
	<i>Cruciata taurica</i> (Pall. ex Willd.) Ehrend.	İbradı, Toka Plateau (Göçebe Location), rocky places, 1106 m, N 37°14'834" – E 31°22'486", 17.v.2008, Çinbilgel 3114; Manavgat, Kepezbeli Location, stony slopes, 1616 m, N 37°10'421" – E 31°21'504", 19.v.2009, Çinbilgel 4804; Manavgat, Kalbur Gözenekleri Location, rocky places, 1788 m, N 37°13'259" – E 31°20'585", 05.viii.2009, Çinbilgel 5694, Ir. – Tur., (H).
	<i>Galium consanguineum</i> Boiss.	İbradı, Suçikan Location, streamsides, 1259 m, N 37°14'503" – E 31°24'173", 02.vii.2010, Çinbilgel 7075, Ir. – Tur., (H).
	<i>Galium floribundum</i> Sm. subsp. <i>floribundum</i>	İbradı, between Güzle Plateau – Yaylacık Mountain, damp places, 1458 m, N 37°10'205" – E 31°23'970", 20.iv.2008, Çinbilgel 2903; Manavgat, Dermebeli Location (Arıtaşı), on the rock and stony places, 1874 m, N 37°13'519" – E 31°20'317", 10.vi.2008, Çinbilgel 3808, (Th).
	<i>Galium incanum</i> Sm. subsp. <i>elatus</i> (Boiss.) Ehrend.	Manavgat, Kepezbeli Location, stony slopes, 1633 m, N 37°10'512" – E 31°21'377", 19.v.2009, Çinbilgel 4798; Manavgat, Kaldırım Mountain (Kızlaroturacağı Location), stony places, 1680 m, N 37°10'883" – E 31°20'961", 03.vi.2009, Çinbilgel 5138; İbradı, Melik Plateau, <i>Pinus nigra</i> forest, 1502 m, N 37°15'548" – E 31°24'365", 30.vi.2010, Çinbilgel 6661, Ir. – Tur., (H).
	<i>Galium murale</i> (L.) All.	Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4830; İbradı, Melik Mountain (Orta Hill), windy hills, 2065 m, N 37°16'532" – E 31°23'131", 17.vi.2010, Çinbilgel 6409; Manavgat, Çukuryurt Plateau, forest clearings, 1293 m, N 37°12'076" – E 31°23'996", 19.v.2011, Çinbilgel 7626, Medit., (Th).
	<i>Galium peplidifolium</i> Boiss.	Manavgat, Akpınar Plateau (Yumrutaş Location), mixed under the <i>Cedrus libani</i> forest, 1520 m, N 37°09'840" – E 31°21'028", 19.v.2009, Çinbilgel 4792; İbradı, Melik Plateau, under the forest, 1446 m, N 37°15'954" – E 31°24'591", 01.vii.2010, Çinbilgel 6753; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 990 m, N 37°10'778" – E 31°25'386", 21.v.2011, Çinbilgel 7692, E. Medit., (Th).
	* <i>Galium sorgerae</i> Ehrend. & Schönb.–Tem.	Manavgat, Kocatop Mountain, windy hills, 2174 m, N 37°12'467" – E 31°18'805", 10.vi.2009, Çinbilgel 5328; Manavgat, Kocatop Mountain, windy hills, 2080 m, N 37°12'398" – E 31°18'732", 10.vi.2009, Çinbilgel 5353; İbradı, Melik Mountain (Orta Hill), rocky places, 1967 m, N 37°16'123" – E 31°22'618", 17.vi.2010, Çinbilgel 6314, End., (H).
	<i>Galium spurium</i> L. subsp. <i>spurium</i>	Manavgat, hillsides of Kaldırım Mountain, rocky places, 1577 m, N 37°09'499" – E 31°20'336", 01.vi.2009, Çinbilgel 4848; İbradı, Melik Plateau, screes, 1491 m, N 37°15'544" – E 31°24'739", 02.vii.2010, Çinbilgel 6995; Manavgat, around of Burmahan Neighbourhood, stony places – roadsides, 704 m, N 37°11'401" – E 31°16'295", 06.vi.2011, Çinbilgel 7764, Euro-Sib., (Th).
	<i>Galium verticillatum</i> Danthoine ex Lam.	Manavgat, Dermebeli Location (Arıtaşı), on the rock and stony places, 1874 m, N 37°13'519" – E 31°20'317", 10.vi.2008, Çinbilgel 3823; Manavgat, Dutluca Plateau, under the rock – stony places, 1282 m, N 37°13'785" – E 31°18'213", 13.vi.2008, Çinbilgel 4065; İbradı, Melik Plateau, meadows, 1446 m, N 37°15'954" – E 31°24'591", 01.vii.2010, Çinbilgel 6761, Medit., (Th).
	<i>Galium verum</i> L. subsp. <i>glabrescens</i> Ehrend.	İbradı, Çeşgar Plateau Breach, dwarf bush formation, 1673 m, N 37°17'288" – E 31°24'324", 01.vii.2010, Çinbilgel 6917, Ir. – Tur., (H).
	<i>Galium verum</i> L. subsp. <i>verum</i>	Manavgat, Susuzkır Hill surroundings, stony places, 1666 m, N 37°16'797" – E 31°21'755", 12.vi.2008, Çinbilgel 3992; Manavgat, Kızılıpınar Plateau (Kocaboğaz Location), arid meadows, 1790 m, N 37°13'949" – E 31°20'094", 29.vi.2009, Çinbilgel 5471; İbradı, Melik Plateau, forest clearings, 1429 m, N 37°15'631" – E 31°24'512", 30.vi.2010, Çinbilgel 6560, Euro-Sib., (H).
	<i>Plocama calabrica</i> (L.f.) M.Backlund & Thulin	İbradı, Suçikan Location, stony places, 1200 m, N 37°13'960" – E 31°24'209", 02.vii.2010, Çinbilgel 7097; Manavgat, Çay Neighbourhood surroundings, <i>Pinus brutia</i> forest, 641 m, N 37°11'714" – E 31°16'060", 06.vi.2011, Çinbilgel 7718; Manavgat, around of Burmahan Neighbourhood (between Burmahan Neighbourhood and Akpınar Plateau), <i>Pinus brutia</i> forest, 846 m, N 37°09'314" – E 31°17'049", 06.vi.2011, Çinbilgel 7857, Medit., (Ch).
	<i>Sherardia arvensis</i> L.	Manavgat, Çay Neighbourhood, <i>Pinus brutia</i> forest, 688 m, N 37°12'133" – E 31°16'452", 27.iii.2011, Çinbilgel 7493; Manavgat, Çay Neighbourhood, Kızılıpınar Plateau way, Kocaarmut Location, <i>Pinus brutia</i> forest clearings –

		stony places, 709 m, N 37°12'174" – E 31°16'360", 27.iii.2011, Çinbilgel 7550, Medit., (Th).
Rutaceae		
	<i>*Haplophyllum pumiliforme</i> Hub.-Mor. & Reese	Manavgat, Çay Neighbourhood, Kavaklı Location, <i>Juniperus excelsa</i> forest, 1374 m, N 37°14'135" – E 31°18'658", 07.vi.2011, Çinbilgel 7919, End., (H).
	<i>Ruta buxbaumii</i> Poir.	İbradı, Suçikan Location, <i>Pinus nigra</i> forest clearings, 1186 m, N 37°14'441" – E 31°24'305", 03.vii.2011, Çinbilgel 8023, (H).
Salicaceae		
	<i>Populus nigra</i> L. subsp. <i>nigra</i>	Manavgat, Çay Neighbourhood, Kızılpınar Plateau way, streamsides, 628 m, 05.viii.2009, Çinbilgel 5740; Manavgat, Akpınar Plateau, roadsides, 1416 m, N 37°09'357" – E 31°20'711", 11.x.2009, Çinbilgel 5826, Euro-Sib., (Ph).
	<i>Salix alba</i> L. subsp. <i>alba</i>	İbradı, Melik Plateau surroundings (towards Suçikan Location), streamsides, 1295 m, N 37°15'056" – E 31°24'473", 02.vii.2010, Çinbilgel 7039, Euro-Sib., (Ph).
Santalaceae		
	<i>Arceuthobium oxycedri</i> (DC.) M.Bieb.	İbradı, Güzle Plateau, Juniper Forest, 1306 m, N 37°09'817" – E 31°23'980", 20.iv.2008, Çinbilgel 2836; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest clearings, 1430 m, N 37°09'494" – E 31°20'838", 19.v.2009, Çinbilgel 4814; İbradı, Çeşgar Plateau, rock, 1623 m, N 37°17'463" – E 31°23'699", 01.vii.2010, Çinbilgel 6855, (on <i>Juniperus drupacea</i>), Euro-Sib., (Vp). (Conf. Göktürk).
	<i>Osyris alba</i> L.	Medit., (Ch), (Observation).
	<i>Thesium billardieri</i> Boiss.	Manavgat, Kovacık Mountain foothills (Gökçebel Location), thorny cushion formation, 1695 m, N 37°13'984" – E 31°19'318", 11.vi.2009, Çinbilgel 5443; İbradı, Yukarıtoka Plateau – Melik Mountain (Orta Hill) foothills, dwarf bush formation, 1879 m, N 37°15'975" – E 31°22'558", 17.vi.2010, Çinbilgel 6287; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1005 m, N 37°11'155" – E 31°25'201", 02.vii.2010, Çinbilgel 7177, Ir. – Tur., (H).
	<i>Viscum album</i> L. subsp. <i>abietis</i> (Wiesb.) Abromerit	İbradı, Yirme Hill (Dikenli Karlıği Location), 1814 m, N 37°14'700" – E 31°23'139", 18.v.2008, Çinbilgel 3229, (on <i>Abies cilicica</i> subsp. <i>isaurica</i>), (Vp).
	<i>Viscum album</i> L. subsp. <i>austriacum</i> (Wiesb.) Vollman	Manavgat, Kızılpınar Plateau, <i>Cedrus libani</i> – <i>Pinus nigra</i> forest, on <i>Pinus nigra</i> , 1631 m, N 37°14'434" – E 31°19'791", 10.x.2009, Çinbilgel 5764, (Vp).
Sapindaceae		
	<i>*Acer hyrcanum</i> Fisch. & C.A.Mey. subsp. <i>sphaerocaryum</i> Yalt.	İbradı, around of Yirme Hill, rocky places, 1805 m, N 37°14'690" – E 31°23'119", 18.v.2008, Çinbilgel 3255; Manavgat, Burunçalı Location (Aslanboğazi), rocky places, 1817 m, N 37°09'027" – E 31°19'488", 02.vi.2009, Çinbilgel 5041; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 1933 m, N 37°16'534" – E 31°22'619", 17.vi.2010, Çinbilgel 6455, End., E. Medit., (Ph).
	<i>Acer monspessulanum</i> L. subsp. <i>monspessulanum</i>	Manavgat, Derme Plateau, rocky hills, 1759 m, N 37°12'512" – E 31°20'675", 19.vii.2008, Çinbilgel 4486; Manavgat, Kalbur Gözenekleri Location, rock crevices, 1751 m, N 37°13'227" – E 31°20'581", 05.viii.2009, Çinbilgel 5707; İbradı, Çeşgar Plateau, rocky places, 1665 m, N 37°17'211" – E 31°24'325", 01.vii.2010, Çinbilgel 6872, (Ph).
Saxifragaceae		
	<i>Saxifraga hederacea</i> L. var. <i>hederacea</i>	Manavgat, Burmahan Neighbourhood, Akpınar way (Öküztaşı Location), rock crevices, 719 m, N 37°10'310" – E 31°16'912", 26.iii.2011, Çinbilgel 7427, (Th).
	<i>Saxifraga tridactylites</i> L.	İbradı, Güzle Plateau, Juniper Forest, 1306 m, N 37°09'817" – E 31°23'980", 20.iv.2008, Çinbilgel 2840; Manavgat, Kovacık Mountain foothills (Hebil Plateau Location), rocky places, 1734 m, N 37°13'822" – E 31°19'204", 09.vi.2009, Çinbilgel 5198; Manavgat, Burmahan Neighbourhood, Erenli Location, on the rock, 611 m, N 37°09'918" – E 31°16'962", 26.iii.2011, Çinbilgel 7466, Medit. (mt.), (Th).
Scrophulariaceae		
	<i>*Scrophularia cryptophila</i> Boiss. & Heldr.	İbradı, around of Yirme Hill (Dikenli Karlıği Location), damp places, 1814 m, N 37°14'700" – E 31°23'139", 18.v.2008, Çinbilgel 3221; İbradı, Çeşgar Plateau, on the rock, 1642 m, N 37°17'662" – E 31°23'989", 17.vi.2008, Çinbilgel 4194; Manavgat, Sulucan Hill, rocky places, 2071 m, N 37°11'496" – E 31°18'847", 11.vi.2009, Çinbilgel 5366, End., E. Medit., (Ch).
	<i>Scrophularia lucida</i> L.	Manavgat, Akpınar Plateau (Hıncırak Location), rubble rocks, 1468 m, N 37°09'337" – E 31°20'412", 01.vi.2009, Çinbilgel 4866; Manavgat, Kovacık Mountain (Uzunçaşır Location), 1937 m, N 37°13'650" – E 31°20'051", 29.vi.2009, Çinbilgel 5450; Manavgat, around of Burmahan Neighbourhood (Payır Location), <i>Pinus brutia</i> forest, 803 m, N 37°09'640" – E 31°17'067", 06.vi.2011, Çinbilgel 7852, Medit., (Ch).
	<i>Scrophularia myriophylla</i> Boiss. & Heldr.	Manavgat, Kovacık Mountain surroundings, rock, 1887 m, N 37°14'043" – E 31°20'377", 09.vi.2008, Çinbilgel 3689; Manavgat, Kaldırım Mountain, screes, 1816 m, N 37°09'893" – E 31°20'398", 01.vi.2009, Çinbilgel 4956; İbradı, Melik Mountain (Orta Hill), windy hills, 2053 m, N 37°16'182" – E 31°22'761", 17.vi.2010, Çinbilgel 6325, E. Medit., (Ch).
	<i>Scrophularia rimarum</i> Bornm.	Manavgat, Kaldırım Mountain (Topukboğazi Location), rock debris, 1737 m, N 37°09'749" – E 31°20'297", 01.vi.2009, Çinbilgel 4909; Manavgat, around

		of Kocatop Mountain (Kürtağacıboğazı Location), 2006 m, N 37°12'817" – E 31°18'915", 10.vi.2009, Çinbilgel 5288; İbradı, Toka Plateau way, Belova Location, rock, 1500 m, N 37°13'716" – E 31°23'564", 02.vii.2011, Çinbilgel 7969, (Ch).
	<i>Scrophularia scopolii</i> Hoppe ex Pers. var. <i>scopolii</i>	Manavgat, Kızılıpınar Plateau, <i>Cedrus libani</i> forest, 1735 m, N 37°14'127" – E 31°19'929", 25.v.2008, Çinbilgel 3560/a; Manavgat, Kepez Plateau (İnçayırı Location), under the rock, 1637 m, N 37°11'073" – E 31°21'011", 11.x.2009, Çinbilgel 5845, (Ch).
	* <i>Verbascum adenocarpum</i> Hub.–Mor.	Manavgat, Akpınar Plateau, Yumrutaş Location, <i>Cedrus libani</i> forest clearings, 1514 m, N 37°09'991" – E 31°21'166", 19.v.2009, Çinbilgel 4732, End., Ir. – Tur., (H).
	* <i>Verbascum glomerulosum</i> Hub.–Mor.	Manavgat, Kepezbeli Location, <i>Cedrus libani</i> forest, 1640 m, N 37°10'269" – E 31°21'453", 03.vi.2009, Çinbilgel 5155; İbradı, Melik Plateau, screes, 1491 m, N 37°15'544" – E 31°24'739", 02.vii.2010, Çinbilgel 6690; İbradı, around of Çukuryurt Plateau, <i>Pinus nigra</i> forest, 1227 m, N 37°12'049" – E 31°24'169", 03.vii.2011, Çinbilgel 8001, End., E. Medit., (H).
	<i>Verbascum lasianthum</i> Boiss. ex Benth.	İbradı, Akmuar Location, stony places, 1033 m, N 37°12'323" – E 31°24'784", 18.vi.2008, Çinbilgel 4440; İbradı, Eynif Plain, arid meadows, 945 m, N 37°08'943" – E 31°26'971", 26.ix.2010, Çinbilgel 7193; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 1006 m, N 37°11'054" – E 31°25'243", 03.vii.2011, Çinbilgel 8038, (H).
	* <i>Verbascum myriocarpum</i> Boiss. & Heldr.	İbradı, Çeşgar Plateau, rock, 1625 m, N 37°17'470" – E 31°23'695", 01.vii.2010, Çinbilgel 6834, End., E. Medit., (H).
	* <i>Verbascum nudatum</i> Murb. var. <i>spatulatum</i> Hub.–Mor.	Manavgat, between Çay and Burmahan Neighbourhoods, <i>Pinus brutia</i> forest, 700 m, N 37°11'519" – E 31°16'218", 06.vi.2011, Çinbilgel 7775, End., E. Medit., (H).
	<i>Verbascum splendidum</i> Boiss.	İbradı, Çeşgar Plateau, dwarf bush formation, 1617 m, N 37°17'485" – E 31°23'565", 17.vi.2008, Çinbilgel 4219; İbradı, Melik Mountain (Hasan Hill), dwarf bush formation, 1933 m, N 37°16'534" – E 31°22'619", 17.vi.2010, Çinbilgel 6448; İbradı, Toka Plateau way, Belova Location, <i>Cedrus libani</i> forest clearings, 1568 m, N 37°13'549" – E 31°23'433", 03.vii.2011, Çinbilgel 7995, E. Medit., (H).
Smilacaceae		
	<i>Smilax aspera</i> L.	Manavgat, Burmahan Neighbourhood surroundings, Akpınar Plateau way, damaged places, 699 m, N 37°10'800" – E 31°16'736", 26.iii.2011, Çinbilgel 7411, (Ph).
	<i>Smilax excelsa</i> L.	Manavgat, Çay Neighbourhood, <i>Pinus brutia</i> forest – streamsides, 688 m, N 37°12'133" – E 31°16'452", 27.iii.2011, Çinbilgel 7476, Euksin (Euro-Sib.), (Ph).
Solanaceae		
	<i>Hyoscyamus niger</i> L.	Manavgat, Akpınar Plateau, stony places, 1403 m, N 37°09'414" – E 31°20'759", 19.v.2009, Çinbilgel 4749; İbradı, Melik Plateau, forest clearings, 1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6958, (Th).
	<i>Solanum decipiens</i> Opiz.	İbradı, Melik Plateau, forest clearings, 1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6955; İbradı, Akmuar Location, streamsides, 1042 m, N 37°11'701" – E 31°24'851", 02.vii.2010, Çinbilgel 7234; İbradı, Belova Location, stony places, 1611 m, N 37°13'501" – E 31°23'451", 02.vii.2010, Çinbilgel 7251, (Th).
Styracaceae		
	<i>Styrax officinalis</i> L.	İbradı, Çukuryurt Plateau surroundings, <i>Pinus nigra</i> forest, 1413 m, N 37°12'686" – E 31°23'903", 09.v.2010, Çinbilgel 6158, (Ph).
Thymelaeaceae		
	<i>Daphne gnidioides</i> Jaub. & Spach	İbradı, Güzle Plateau (towards Çobanuçtuğu Location), forest, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2809; İbradı, southeast slopes of Yaylacık Mountain, <i>Cedrus libani</i> forest clearings, 1263 m, N 37°09'958" – E 31°25'227", 19.iv.2009, Çinbilgel 4609; İbradı, Boğazyurt Location, <i>Juniperus excelsa</i> forest, 991 m, N 37°10'804" – E 31°25'375", 02.vii.2010, Çinbilgel 7206, E. Medit., (Ph).
	<i>Daphne oleoides</i> Schreb. subsp. <i>oleoides</i>	İbradı, Yukartoka Plateau (between Beydiğin Plateau and Göçebe Location), rocky places, 1543 m, N 37°14'553" – E 31°22'450", 17.v.2008, Çinbilgel 3067, (Ch).
	<i>Daphne sericea</i> Vahl. subsp. <i>sericea</i>	Manavgat, Akpınar Plateau, under the <i>Cedrus libani</i> forest, 1480 m, N 37°09'593" – E 31°21'076", 19.v.2009, Çinbilgel 4810; İbradı, Cevizlidere Location, stony places, 1177 m, N 37°13'424" – E 31°24'206", 05.iii.2011, Çinbilgel 7403, E. Medit., (Ph).
Ulmaceae		
	<i>Ulmus glabra</i> Huds.	İbradı, Yirme Hill (Dikenli Karlığı Location), rocky places, 1814 m, N 37°14'700" – E 31°23'139", 18.v.2008, Çinbilgel 3219; Manavgat, Derme Plateau, rocky places, 1740 m, N 37°12'694" – E 31°20'565", 19.vii.2008, Çinbilgel 4491; İbradı, between Yukartoka Plateau and Melik Mountain foothills, dwarf bush formation, 1791 m, N 37°15'442" – E 31°23'012", 18.vi.2010, Çinbilgel 6545, Euro-Sib., (Ph).
Urticaceae		

	<i>Parietaria lusitanica</i> L.	Manavgat, Arğıça Location (cutting area), <i>Pinus brutia</i> forest, 1054 m, N 37°12'995" – E 31°17'105", 13.vi.2008, Çinbilgel 4124, Medit., (Th).
	<i>Urtica dioica</i> L. subsp. <i>dioica</i>	İbradı, Yirme Hill (Dikenli Location), <i>Cedrus libani</i> forest clearings, 1783 m, N 37°14'667" – E 31°23'097", 18.v.2008, Çinbilgel 3194; İbradı, Çeşgar Plateau Location, stony slopes, 1572 m, N 37°17'675" – E 31°23'744", 17.vi.2008, Çinbilgel 4209; İbradı, Melik Plateau, forest clearings, 1398 m, N 37°15'575" – E 31°24'690", 02.vii.2010, Çinbilgel 6964, Euro-Sib., (H).
Verbenaceae		
	<i>Verbena officinalis</i> L. var. <i>officinalis</i>	Manavgat, Burmahan Neighbourhood, Erenli Location, <i>Pinus brutia</i> forest – watersides, 729 m, N 37°09'926" – E 31°17'026", 11.x.2009, Çinbilgel 5804; İbradı, Suçikan Location, forest clearings, 1222 m, N 37°14'460" – E 31°24'254", 02.vii.2010, Çinbilgel 7079; İbradı, Akmuar Location, streamsides, 1042 m, N 37°11'701" – E 31°24'851", 02.vii.2010, Çinbilgel 7233, (H).
Violaceae		
	<i>Viola heldreichiana</i> Boiss.	İbradı, Suçikan Location way, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008 Çinbilgel 2747; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest, 1496 m, N 37°09'444" – E 31°20'567", 02.v.2009, Çinbilgel 4646; İbradı, Melik Mountain surroundings, dwarf bush, 1735 m, N 37°15'390" – E 31°22'574", 18.vi.2010, Çinbilgel 6481, E. Medit., (Th).
	<i>Viola kitaibeliana</i> Roem. & Schult.	İbradı, Güzle Plateau – hillsides of Yaylacık Mountain, damaged <i>Pinus nigra</i> forest and meadows, damp places, 1430 m, N 37°10'227" – E 31°23'913", 20.iv.2008, Çinbilgel 2883; İbradı, Melik Mountain surroundings, rocky – stony places, 1875 m, N 37°14'859" – E 31°23'148", 18.v.2008, Çinbilgel 3278; İbradı, Toka Plateau Breach surroundings, 1570 m, N 37°13'602" – E 31°23'417", 09.v.2010, Çinbilgel 6131, (Th).
	<i>Viola occulta</i> Lehm.	Manavgat, Burunçalı Location, slopes, 1883 m, N 37°09'116" – E 31°18'988", 02.vi.2009, Çinbilgel 5055, (Th).
	<i>Viola odorata</i> L.	İbradı, southern slopes of Yaylacık Mountain, stony places, 1541 m, N 37°10'262" – E 31°24'049", 18.iv.2009, Çinbilgel 4572; İbradı, Toka Plateau way, <i>Pinus nigra</i> forest, 1662 m, N 37°13'925" – E 31°22'981", 08.v.2010, Çinbilgel 6006; Manavgat, Çay Neighborhood, Kızılpınar Plateau way (Düğnük Plateau), degraded <i>Pinus brutia</i> forest, 1163 m, N 37°13'385" – E 31°17'716", 27.iii.2011, Çinbilgel 7508, (H).
	<i>Viola sieheana</i> W.Becker	İbradı, Suçikan Location way, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2773; İbradı, Toka Plateau (Beydiğin Plateau), <i>Pinus nigra</i> forest clearing, stony places, 1544 m, N 37°14'435" – E 31°22'650", 18.v.2008, Çinbilgel 3159; İbradı, Gevenli Location, <i>Pinus brutia</i> forest, 1047 m, N 37°12'249" – E 31°24'781", 19.v.2011, Çinbilgel 7612, (H).
Vitaceae		
	<i>Ampelopsis orientalis</i> (Lam.) Planch.	Manavgat, Dutluca Plateau, under the rock – stony places, 1282 m, N 37°13'785" – E 31°18'213", 13.vi.2008, Çinbilgel 4054; Manavgat, Akpınar Plateau (Deller Location), mixed <i>Cedrus libani</i> forest, 1461 m, N 37°09'305" – E 31°20'361", 01.vi.2009, Çinbilgel 4829; İbradı, Melik Plateau, streamsides, 1537 m, N 37°15'549" – E 31°24'633", 02.vii.2010, Çinbilgel 6952, E. Medit., (Ph).
	<i>Vitis vinifera</i> L.	Manavgat, Kızılpınar Plateau way, between Akin and Karahan Locations, <i>Pinus brutia</i> forest, 740 m, N 37°13'257" – E 31°15'440", 13.vi.2008, Çinbilgel 4139; Manavgat, between Çay and Burmahan Neighbourhoods, streamsides, 706 m, N 37°11'423" – E 31°16'306", 06.vi.2011, Çinbilgel 7759, (Ph).
Xanthorrhoeaceae		
	<i>Asphodeline lutea</i> (L.) Rchb.	İbradı, Suçikan Location way, stony places, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2756; İbradı, Güzle Plateau surroundings, forest clearings, 1267 m, N 37°09'621" – E 31°24'201", 20.iv.2008, Çinbilgel 2812; Manavgat, Akpınar Plateau, <i>Cedrus libani</i> forest clearings, 1430 m, N 37°09'494" – E 31°20'838", 19.v.2009, Çinbilgel 4815, Medit., (G).
	<i>Eremurus spectabilis</i> M.Bieb.	İbradı, Suçikan Location way, damp places, 1202 m, N 37°14'424" – E 31°24'268", 19.iv.2008, Çinbilgel 2738; İbradı, Yirme Hill (Dikenli Karlığı Location), damp places, 1814 m, N 37°14'700" – E 31°23'139", 18.v.2008, Çinbilgel 3223, Ir. – Tur., (G).

Abbreviations used in floristic list are as follows: End.: Endemic, E.: East, Medit.: Mediterranean element, Ir-Tur.: Irano-Turanian element, Euro.-Sib.: Euro-Siberian element, mt: mountain, (Ph): Phanerophyte, (Ch): Chamaephyte, (H): Hemicryptophyte, (Th): Therophyte, (G): Geophyte, (Vp): Vascular Parasite.